

DETERMINATION OF STARCH BY IODINE COLORIMETRY

K. A. VAINIO

Department of Animal Husbandry, University of Helsinki

Received January 5, 1968

In 1814 de CLAUBRY (ref. BRAUTLECHT, 1953, p. 372) reported that the addition of iodine to starch solutions yields a blue coloration. Later investigations (see comprehensive survey of SAMEG, 1927) have demonstrated the specificity and exceptional sensitiveness of this colour reaction. Nevertheless, competent methods for starch determination based on this principle have not been published before the works of PALOHEIMO (1930, p. 150; 1948, p. 109; PALOHEIMO and PALOHEIMO 1931, p. 391). The main reason for the discrimination of the iodine colorimetry principle in starch determination lies evidently in the fact that the colour tone which the iodine gives to a starch solution depends both on the starch concentration and the iodine concentration of the solution so that a definite kind of colour is attained only when both concentrations are definite. In addition the colour tone depends on the temperature of the solution. These circumstances make it difficult to use ordinary photometers in iodine colorimetric starch determinations. In addition it is noticeable that when boiled in water only that part of the starch in the ground sample which comes from the broken cells occurs as colloid suspension. The starch of unbroken cells gelatinizes but remains inside the cell walls.

In the method of Paloheimo the starch is converted into dextrans which escape even the unbroken cells. The determination is performed in the following way which, with some minor modifications, also the present author has used.

1 g of the material to be analysed is ground with some water in a mortar (even when the material has beforehand been ground with a mill). The sample is then rinsed into a 600- ml beaker which has a mark at 400 ml. About 350 ml distilled water is added and the mixture is brought to boil. 20 ml of 1 N sulphuric acid and boiling water are added up to the mark. The acid normality is now 0.05. The mixture is boiled for 30 minutes by compensating the loss of evaporation with boiling water, and filtered through a rough filter. The filtering with washing takes about 3 minutes. The filtrate is poured into a 500- ml volumetric flask, the flask is rapidly cooled to room temperature and make up to the volume. This is solution BI.

Solution AI is made in the same manner as BI but of 500 mg of pure starch, if possible of the same kind as in the analysis sample. Filtration is not necessary but the solution in the volumetric flask must stay hot 3 minutes before cooling.

A 5-% solution of KJ is saturated with iodine. For the colorimetric use 1 volume of this solution is diluted with 3 volumes of water.

25 ml of solution AI is transferred into a 500-ml volumetric flask and about 400 ml of water are poured upon it. 5 ml of the iodine solution is added under simultaneous shaking. The flask is immediately made up to the volume with water and shaken. This is solution AII. Solution BII is made of solution BI in the same manner but the pipetted amount of BI must be large enough to ensure to BII a somewhat greater starch concentration than there is in AII. Thus the colour of BII must be deeper than that of AII. A diluting solution, solution C, for the colorimetric measurement is obtained when 5 ml of iodine solution is diluted with water up to 500 ml.

For comparator 17 cm high optical cells with bottom measures of 40 mm (optical depth) and 50 mm are used. Two cells, A and B, are needed. The comparator is home made. Cell A is filled with solution AII and in cell B 100 ml of solution BII is poured. BII in cell B is now diluted with solution C until B has reached the colour of A. As the iodine concentration of solution C is the same as in solutions AII and BII the iodine concentration in cell B remains unchanged during the dilution. When the colour in A and B is the same one may conclude that also the starch concentration is the same. The concentration in A is known and when the liquid volume in B is measured the amount of starch which this cell contains can be calculated. Further, the amounts of starch in the whole BII solution and finally in the BI solution can be calculated. If the starch preparation used for solution AI contains 90 % starch, the solution AII contains 0.045 mg of starch per 1 ml.

If the starch used for solution A I is not botanically of the same kind as in the material to be analysed, the result must be corrected using an equivalency coefficient. According to Paloheimo 1 g of dry wheat starch is colorimetrically equivalent to

0.82 g of potato starch	1.01 g of oat starch
0.97 » » barley »	1.09 » » rice »
1.01 » » rye »	

Experiments and results

Comparator and comparison of the coloured solutions. Paloheimo has used a home made comparator without any lenses and prisms. With this apparatus he has attained very reliable results. The only difficulty is caused by the fact that the two fields of vision which are to be compared are not closely joined to each other. The present author has used a comparator with prisms and lenses (Fig. 1) which creates a circular field of vision in which one half gets the light through the optical cell A and the other through cell B. This comparator was composed of parts of a Pulfrich photometer no longer in use in the department. The intensity of light can be adjusted by the adjusting drums of the Pulfrich photometer, or by putting frosted glass plates before or behind the optical cells, or by adjusting the current in the lamp. A proper comparator can also be ordered from any skilled optician. When diluting solution BII in the optical cell the author has added the solution C through a burette.




Fig. 1. Schematic picture of the comparator. L lamp, M mirror, C optical cell, O ocular.

I n t e n s i t y o f b o i l i n g. As in preparing solutions AI and BI (cf. p. 61 and p. 60) the starch is converted into dextrins, it is important that the two solutions are boiled in the same temperature. Differences in atmospheric pressure do not interfere here as both solutions are boiled simultaneously or successively. It is important however, that the two solutions are boiled with about the same intensity. The author observed that when the temperature measured from the middle part of the liquid in the beaker was 100.5° during very intensive boiling, it was 98.5° if the boiling was gentle. When solution AI (cf. p. 61) was prepared by gentle boiling and solution BI, made of 500 mg of potato starch, by very intensive boiling, only 85 % of the starch was recovered. Experiments with wheat starch gave approximately the same results. Experiments with barley meal gave 53.2 % starch by gentle boiling but 49.0 % by intensive boiling. These experiments were made in extreme conditions. From the figures in Table 1 it can be concluded that in routine work there is no difficulty in preparing the solutions AI and BI in similar conditions.

In all determinations made for this study magnetic mixing has been used. When water is being added into the beaker the mixing should be interrupted in order to make it possible to note the position of the liquid surface.

A c i d i t y d u r i n g t h e b o i l i n g. The experiments of PALOHEIMO (1931, p. 395) show that the method is not very sensitive to small variations in the normality of the acid solution when preparing the BI solutions. Normal foods are able to cause only insignificant deviations from the normality figure of 0.05. However, with food mixtures into which e.g. chalk is added one must first determine the neutralizing power of the sample and calculate how much 1-N H_2SO_4 is to be added to obtain a 0.05-N solution. If the sample contains acid it should be extracted beforehand with 80-% ethanol.

C o l o r i m e t r i c a l s e n s i t i v e n e s s o f s t a r c h s o l u t i o n s. 500 mg of wheat starch was weighed and solution AI (cf. p. 61) was prepared. As the purity of the starch preparation was 90 %, the solution contains 0.9 mg starch per ml. A series of diluted solutions was made by using 100 ml volumetric flasks into which the following amounts of AI were transferred: 1 ml, 0.5 ml, 0.3 ml, 0.2 ml, 0.1 ml, 0.05 ml and 0 ml. In each flask 1 ml of iodine solution (for colorimetric use, p. 61) was added, and the flasks were made up to the volume. Using the comparator the colour of the solutions was compared with the colour of the 0-ml solution. This comparison was performed both in 20° and 3° . The iodine concentration of the solutions is the same as that in the solutions AII and BII in the method of Paloheimo.

A similar series was prepared and studied with potato starch the purity of which was 86.5 %. Two further series were studied, one with wheat starch and the other with potato starch, in the preparing of which no acid was used but only 1/2 hours' boiling in water. These series were studied only in 20°.

The purpose of these experiments was to investigate the utmost concentration limits of starch which still cause a visible change in the colour of the iodine solution. These minimum concentrations were:

1. Wheat starch. Boiled in water	20°	0.18 mg/100 ml
2. » » » » acid	20°	0.27 »
3. » » » » »	3°	0.09 »
4. Potato » » » water	20°	0.18 »
5. » » » » acid	20°	0.09 »
6. » » » » »	3°	0.05 — 0.09 »

Influence of cellulose, ethanol, sugars and proteins. In some cases starch may be determined of the residue on a filter paper. It is therefore important to know whether it is possible to prepare solution BI (cf. p. 60) by boiling the residue together with the paper. In order to study this point a BI solution was prepared using 100 mg of wheat starch and a Whatman No. 4 filter paper. The paper (Ø 15 cm, 1,5 g) was torn to pieces. All the starch was recovered by iodine colorimetry.

If the material to be analysed is to be extracted with ethanol one should know to what extent the ethanol is to be removed from the material, by evaporation or by washing with water, before the starch determination. For studying this question 1 g of barley meal was boiled in 80-% ethanol for 4 hours. After filtration through a Whatman No. 4 paper (Ø 15 cm) the residue with the filter paper was allowed to stand for some minutes and was then boiled as usual when preparing solution BI. Colorimetrically 537 mg of starch per 1 g of barley dry matter were recovered as the mean of two determinations. When the starch was determined from barley meal without an ethanol extraction the corresponding figure was 530. Evidently the ethanol content in solution B I was so small that it could not interfere with the hydrolysis of the starch, and in solution BII so low that it was not able to depress significantly the iodine absorption of the dextrine solution. However, when the concentration of starch in BI is low, a greater amount of this solution should be taken for solution BII. In such a case it is advisable to leave the solution to boil for about 30 minutes before the addition of the acid.

When the BI solution was prepared from 100 mg of wheat starch with 1 g of saccharose added, the starch was entirely recovered. As the 1/2 hours' boiling in 0.05-N H₂SO₄-solution is sufficient for the inversion of saccharose one may conclude that even glucose and fructose do not have a disturbing influence on colorimetric starch determination. It was also proved that maltose and lactose have no such influence.

As some proteins and simpler nitrogenous compounds are able to bind iodine it seemed reasonable to examine whether such substances could possibly interfere with the starch determination. For that purpose a BI solution was prepared by using 100 mg of wheat starch and 20 ml of skim milk. The amount of milk used contained about 650 mg of protein. All the starch was recovered. The result was the same when 250 mg of gelatine was used.

However, when the amount of gelatine was 1 g the solution BII turned a greenish colour and the colorimetric comparison was very difficult. The same result was obtained with a casein preparation intended for food mixtures of laboratory animals. Evidently this casein was partially hydrolysed. Likewise, when about 3 g of meat poor in fat was boiled with 100 mg of starch the greenish colour appeared and the colorimetric comparison was not possible.

However, judging by the figures in Table 1 plant proteins seem not to have a disturbing influence upon iodine colorimetric starch determination.

Iodine colorimetric method compared with the amyloglucosidase method. In this department SALO (1968, p. 41) has worked out a very reliable method in which the starch is converted with amyloglucosidase into glucose and

Table 1. Starch percentages obtained by the iodine colorimetric method compared with those obtained by the amyloglucosidase method of Salo (% of dry matter).

	Colorimetric method			Amyloglucosidase method
	a	b	mean	dase method mean
1. Wheat kernels	57.7	57.3	57.5	58.2
2. Rye »	54.5	54.7	54.6	57.0
3. Barley »	53.7	53.1	53.4	53.6
4. Oat »	48.5	48.1	48.3	48.7
5. Wheat bran	22.2	21.9	22.1	21.5
6. Peas	40.2	41.3	40.8	40.2
7. Peanut cake	8.0	8.1	8.1	7.1
8. Soy meal (extracted)	1.8	1.8	1.8	3.3
9. Potatoes, peeled	59.6	60.0	59.8	59.1
10. Swedes, peeled	0.7	0.8	0.8	1.4
11. Carrots	3.0	3.1	3.1	3.8
12. Celery	9.9	10.3	10.1	10.3
13. Luzern	2.9	2.9	2.9	3.1
14. Rootstocks of horsetail (Eq. palustre)	9.3	9.8	9.6	10.9

determined as glucoseanhydride. The figures in Table 1 obtained with the amyloglucosidase method are results of Salo's work. The present author has made colorimetric estimations from the same samples. It appears that the difference between the parallel results is only in one case higher than 0.5 pct-units. The difference between the colorimetric and amyloglucosidase mean values is only in 4 cases out of 14 higher than one pct-unit, and the amyloglucosidase value is in 10 cases out of 14 higher than the colorimetric value.

In the determinations 15 ml of solution AI (cf. p. 61) was usually used for preparing solution A II. For foods 8, 10, 11 and 13, the corresponding volume was 10 ml. As to the food 12, the colour tone in the optical cell B (cf. p. 61) did not become the same as in cell A, 10 ml of the iodine solution instead of 5 ml was used in preparing the solutions AI, BI and C. When estimating the starch from foods 7, 8, 10, 11, 13 and 14, an 1 g sample was first extracted by boiling in 80-% ethanol in order to reduce the amount of coloured substances. This procedure may have been unnecessary in some of the cases.

Starch determination from faeces. Sometimes faeces, especially those of swine, contain starch included in food particles which have escaped mastication. It is consequently important that also this starch is estimated. For this reason the author has examined whether some faeces constituents might interfere in the colorimetric starch determination. 100 mg of wheat starch was mixed into 5 g of fresh cow faeces and the starch determined from the mixture. All added starch was recovered. The result was the same when 50 mg of starch was mixed into 6 g of faeces. In these cases the mixture was extracted beforehand with 80-% ethanol.

Summary

In the iodine colorimetric method of Paloheimo gently dextrinized solutions are prepared of pure starch and of the analysis sample. One of the optical cells (A) of the comparator is provided with a solution made of pure starch and the other (B) with the solution to be analysed. Both solutions have the same iodine concentration. The solution in B must have a intensive colour than that in A. Solution B is then diluted with an iodine water solution of the same iodine concentration as in the solutions A and B. When these solutions have attained the same colour it is concluded that also the starch concentration is the same and the starch content of the sample can be calculated.

The results obtained by this method are compared with those obtained with the amyloglucosidase method of Salo. Table 1 shows that the two methods give very similar results.

Different circumstances which might possibly interfere with the colorimetric starch determinations are studied. — It was observed that attention must be paid to the intensity of boiling when the 0.05-N H_2SO_4 dextrinizing solutions are boiled. If the intensity is very different in the comparison solution and the solution to be analysed, considerable errors may occur. — If the sample contains added chalk the neutralizing power of the sample should be determined beforehand and the normality of the solution adjusted to 0.05. If the sample contains acid it should be extracted beforehand with 80-% ethanol. — Cellulose and sugars have no influence on the results, nor have plant proteins or proteins of milk. However, if greater amounts of protein were added, a casein preparation intended for laboratory animals showed an obvious disturbing effect, as did gelatin and meat protein. — Faeces did not appear to have an interfering influence in colorimetric starch determination.

The iodine colorimetric sensitiveness of starch solutions was also studied. It appeared that 0.18 mg of dextrinized potato starch already deepened the colour of 100 ml dilute iodine solution in room temperature. For wheat starch the corresponding minimum concentration was 0.27 mg/100 ml. In 3° the concentration limit was even lower, 0.05—0.09 mg/100 ml.

In all the above mentioned studies the author has used as comparator essential parts of a Pulfrich photometer. A proper comparator (Fig. 1) can also be made by any skilled optician.

REFERENCES

- BRAUTLECHT, C. A. 1953. Starch, its sources, production and uses. New York. 1—408.
 PALOHEIMO, L. 1930. Zur Verwendbarkeit des jodkolorimetrischen Prinzips bei Stärkebestimmungen. *Bioch. Z.* 222: 150—172.

- PALOHEIMO, L. 1948. Determination of starch according to the principle of iodine colorimetry. *J. Sci. Agric. Soc. Finland* 20: 109—113.
- PALOHEIMO, L. und PALOHEIMO, I. 1931. Beiträge zur Jodkolorimetrie der Stärke nach der Methode von Paloheimo. *Bioch. Z.* 238: 391—400.
- SALO, M. - L. and SALMI, M. 1968. Determination of starch by the amyloglucosidase method. *J. Sci. Agric. Soc. Finland* 40: 38—45.
- SAMEC, M. 1927. *Kolloidchemie der Stärke*. Dresden und Leipzig. 1—509.

SELOSTUS

JODIKOLORIMETRINEN TÄRKKELYSMÄÄRITYS

K. A. VAINIO

Kotieläintieteen laitos, Helsingin yliopisto

Paloheimon jodikolorimetrisessä tärkkelysmääritysmenelmässä valmistetaan 0.05-N-rikkihapolla heikosti dekstrinoidut liuokset puhtaasta tärkkelyksestä ja analyysinäytteestä (liuokset AI ja BI). Näistä liuoksista tehdään laimennokset, joihin kumpaankin järjestetään sama jodikonsentraatio: saadaan liuokset AII ja BII. Laimennus on suoritettava siten, että BII tulee tummemmaksi kuin AII. Kolorimetrilaseja tarvitaan kaksi, A ja B. Lasiin A otetaan värinäytteeksi liuosta AII. Lasiin B kaadetaan määrätillavuuksu liuosta BII, joka sitten tärkkelyksen suhteen laimennetaan liuoksella C. Liuos C ei sisällä tärkkelystä, mutta sen jodiväkevvyys on sama kuin liuoksen AII ja BII. Liuosta BII laimennetaan asteettain, kunnes värihavuus kolorimetrilaseissa tulee samaksi. Liuos C:n kulutuksen perusteella voidaan nyt laskea BII:n ja siitä edelleen ko. näytteen tärkkelyspitoisuus.

Kirjoittaja on eräissä suhteissa muuntanut Paloheimon menetelmää. Niinpä on värivertailussa käytetty Pulfrich-fotometrin tarkoitukseen sopivia osia (Kuva 1.) ja liuos C lisätään byretistä. Edelleen on tutkittu tärkkelyksen jodireaktion herkkyyttä, kiehumisvoimakkuuden ja happoväkevyyden vaikutusta dekstrinointikeitossa sekä eräiden tutkittavissa näytteissä mahdollisesti esiintyvien aineiden vaikutuksia. Lopuksi on jodikolorimetrisellä menetelmällä saatuja tuloksia verrattu Salon amyloglukosidaasimenetelmällä saatuihin.

Tärkkelyksen jodireaktio osoittautui erittäin herkäksi. Niinpä 0.18 mg perunan tärkkelystä riitti aikaansaamaan värimuutoksen 100 ml:ssa laimeata jodiliuosta. — Kiehumisvoimakkuudella todettiin olevan varteenotettava vaikutus. — Jos tutkittava aines sisältää CaCO_3 -lisäyksen, on näytteen neutraloiva vaikutus etukäteen tutkittava ja dekstrinointihappamuus järjestettävä 0.05-normaaliseksi. Jos näyte sisältää happoja, on se edeltävästi uutettava 80-% etanolilla. — Suuretkaan määrät filteripaperia, sokereita, kasvivalkuaista tai maidon valkuaista eivät vaikuta häiritsevästi. Sen sijaan preparoidulla kaseiinilla, liivateella ja lihavalvakuaisella on häiritsevä vaikutus. Lehmän sonnassa ei osoittautunut olevan aineita, jotka haittaisivat tärkkelyksen määrittämistä.

Salon amyloglukosidaasimenetelmä antaa ilmeisesti erittäin luotettavia tuloksia. Tästä syystä tutkittiin jodikolorimetrisen menetelmän pätevyyttä vertaamalla sillä saatuja tuloksia Salon menetelmällä samoista näytteistä saatuihin. Taulukosta 1 nähdään, että näillä kahdella tavalla saadut tulokset ovat keskenään varsin yhtäpitäviä.

Kirjoittajan suorittamat tutkimukset osoittavat, että jodikolorimetrisen menetelmä soveltuu erittäin hyvin tärkkelyksen määrittämiseen. Lisäksi tämä menetelmä on varsin joutuisa. Ilmankin prismoilla ja linseillä varustettua kolorimetriä, kotitekoista kolorimetrikameraa käyttäen, saadaan tuloksia, joiden tarkkuus on useihin tarkoituksiin riittävä.