

ESTIMATION OF CARCASS CHARACTERISTICS IN YOUNG BEEF CATTLE

Selostus: Nuoren lihanaudan teurasominaisuuksien arvioimisesta

HILKKA RUOHOMÄKI

Agricultural Research Centre, Institute of Animal Breeding.
01301 Vantaa 30



TABLE OF CONTENTS

	Page
Abstract	389
INTRODUCTION	389
PURPOSE OF PRESENT STUDY	390
MATERIAL AND METHODS	391
Material	391
Estimated traits and estimators	392
Presentation of the results	393
Statistical analysis	393
ESTIMATED TRAITS	394
Live weight	394
Carcass weight	394
Quantities and percentages of carcass meat, fat and bone	394
Partition of different types of tissue	394
Means of estimated traits by experiment	396
Ist ESTIMATOR: live measurements	398
Measuring technique	398
Means of live measurements by experiment	399
Stepwise regression analysis, results and discussion	402
Estimation of live weight, carcass weight and carcass composition with live measurements	
IIInd ESTIMATOR: initial weight and daily gain	409
Initial weight	409
Daily gain during experiment gr./day	409
Means of initial weights and daily gains by experiment	409
Stepwise regression analysis, results and discussion	410
Estimation of carcass weight and carcass composition with initial weight and daily gain	
IIIrd ESTIMATOR: carcass quality scores	411
Evaluation technique	411
Means of carcass quality scores by experiment	413
Stepwise regression analysis, results and discussion	413
Estimation of carcass composition with carcass quality scores	
IVth ESTIMATOR: live weight, carcass weight, dressing-%, by-products and cutting results of flanks	419
Dressing-%	419
By-products	420
Dissection of flanks	420
Means of dressing-%, by-products and cutting results of flanks by experiment	420
Stepwise regression analysis, results and discussion	420
Estimation of carcass composition with live weight, carcass weight, dressing-%, by-products and cutting results of flanks	

Vth ESTIMATOR: cutting results of shanks	424
Dissection of shanks	424
Means of cutting results of shanks by experiment	425
Stepwise regression analysis, results and discussion	426
Estimation of carcass composition with cutting results of shanks	
VIth ESTIMATOR: cutting results of the fore and hind quarters	430
Means of cutting results of the fore and hind quarters by experiment	430
Stepwise regression analysis, results and discussion	431
Estimation of carcass composition with carcass weight and cutting results of the fore and hind quarters	
VIIth ESTIMATOR: detailed cutting results of the carcasses	436
Results and discussion	436
Estimation of carcass composition with detailed cutting results	
SUMMARY	440
REFERENCES	441
SELOSTUS	444

RUOHOMÄKI, H. 1975. **Estimation of carcass characteristics in young beef cattle.**
J. Scient. Agric. Soc. Finl. 47: 385—444.

Abstract. In the present study the main objectives were to determine whether sufficiently accurate estimation of live weight, carcass weight and carcass composition with live measurements is possible, and whether an accurate estimation of carcass composition with any other procedure than total carcass dissection is possible. The material consisted of 205 1-year old steers and bulls and 174 bull calves. Within experiment each trait of each animal was corrected for age, breed and feeding with a correction factor obtained with the least squares method. In the stepwise regression analysis the results for both age groups were analysed separately. From live measurements the best estimators in both age groups were width of chest, heart girth and natural length. The most reliable predictor of meat quantity was carcass weight, R^2 -value was 96 % for both age groups. In estimation of meat percentage the R^2 -value obtained for the $\frac{1}{2}$ -year olds with cutting results of fore shanks was 56 %, the respective value for the 1-year olds was considerably lower. With 1-year olds the carcass weight and some by-products gave the highest estimations for the quantity and percentage of fat, R^2 were 62 % and 50 % respectively, and for the $\frac{1}{2}$ -year olds the kidney fats and the cutting results of flanks gave the highest estimations, R^2 were 70 % and 59 % respectively. For the quantity and percentage of bone the R^2 for the 1-year olds obtained with cutting results of shanks were 68 % and 55 % respectively, and for the $\frac{1}{2}$ -year olds 82 % and 70 % respectively. When the cutting results of the fore and hind quarters were used as estimators the R^2 obtained for all the traits were higher in both age groups than with any other estimator, R^2 obtained with carcass quality scores remained lower.

Introduction

Methods to increase meat production and improve carcass quality have been objects of intensive research since World War II. This is true especially in the developed countries where meat consumption together with requirements on meat quality have increased by both the consumers and the industry.

The main purpose of research is to discover methods for the economical production of carcasses with large deposits of high-quality meat, sufficiently fats and as little as possible uneatable tissues such as bone and tendons.

Increase in weight and size, i.e. growth, and feed consumption are the only information available during the growth of beef cattle. On the economic point of view these are the most important factors in beef cattle production. Under the prevailing circumstances, however, it is not always possible to verify the weight and growth by any other means but measuring, because only a few farmers have an animal scale at their disposal.

No sufficiently accurate information can be obtained of carcass weight and composition — quantities and percentages of meat, fat and bone — of an animal during its growth. The quantities of meat, fat and bone can only be determined through dissection of the whole carcass and partition of different types of tissue. The accuracy required by scientific research can only be obtained if the different types of tissue are separated anatomically, where, for example, the intermuscular fasciae and fat can be removed from the muscle.

As partition of carcasses into different tissues is very expensive and time-consuming even in research, other methods to determine carcass composition are being examined. Estimation of carcass composition has always been started from visual evaluation and live measurements, continued to carcass measurements and quality scores, up to the inspection of different types of tissue in some small parts of a carcass.

As fat content strongly affects the rations between different types of tissue in a carcass, several techniques and devices have been developed for measuring fat, of which the ultrasonic measuring device is the most delicate and the most widely used one. In measuring the thickness of back fat in pigs some very prominent results have been obtained. In cattle, however, no equally reliable results have been obtained especially when the fat cover has been thin.

The use of different techniques and devices is no further discussed here, but it can be observed that the further the measurement and study technique develops, the more reliable information on live animal and carcass composition is obtained.

Purpose of present study

Although foreign results on the subject matter are relatively abundantly available, they are in their original form mostly unapplicable for Finnish conditions. Materials consist of foreign breeds, feeding is different, carcass classification methods differ from each other and from ours, etc. The greatest handicaps for comparison are, however, in the different cutting methods and quality requirements which vary between countries and seasons and which production finally has to adapt to.

The purpose of the present study is to determine whether adequately accurate estimation of live weight, carcass weight and carcass composition with live measurements on young beef cattle is possible, and whether carcass composition — quantities and percentages of meat, fat and bone — can be estimated without decreasing the commercial value of the carcass at all or only slightly.

Material and methods

Material

The material comprised results of 10 separate experiments. Eight of them were conducted at the S. W. Finland Agricultural Experiment Station in 1963–70, and two at the North Ostrobothnia Agricultural Experiment Station in 1969–70. In experiments I–VI there were 205 about 1-year old steers and bulls, and in the experiments VII–X 174 bull calves reared till age of 165 days. In the present study the former are called 1-year olds and the latter $\frac{1}{2}$ -year olds.

The experiments were planned and supervised by the Institutes of Animal Breeding and Animal Husbandry of the Agricultural Research Centre.

In the experiments for the 1-year olds the initial feeding was roughly taking equal, but thereafter the animals were arranged to several feeding groups, except in experiment I. The differences in feeding were mainly based on the unequal portions of concentrates, the varying levels of nitrogen content on pasture and in green silage fodder, and on the various methods the silage fodder had been prepared. In experiments VII–IX feeding remained uniform, in experiment X the effect of different portions of animal protein on the development of an animal was also studied.

The calves were delivered for the experiments by the local A. I. Society and co-operative slaughterhouse. The attempt was to obtain the calves at 1–2 weeks' age which, however, did not always succeed especially with the crossbreds.

In the experiments I and V the calves were born in spring and slaughtered the following spring. In the other experiments with 1-year olds the calves were born in autumn and slaughtered the following autumn at the end of pasture season, except in experiment IV where half of the animals stayed in the cow stable on indoor feeding over the summer. Calves in the experiments with $\frac{1}{2}$ -years olds were all born in autumn.

The purpose was that all the animals were sent for slaughter at the same age within experiment. Age differences were levelled by gradating the slaughters. The slaughter age for $\frac{1}{2}$ -year olds was predetermined to 165 days and that of the others to about one year.

Rearing till one year was justifiable especially in experiments IV, V and VI where there were progeny groups of phenotype tested bulls which had been tested at rearing stations at the age of one year. In some experiments also other criteria could have been considered for determining the moment of slaughter. Some of the most noteworthy being slaughter at same weight or at the same fatness degree, like in the study by LINDHÉ and HENNINGSSON (1968). However, an accurate determination of fatness degree would have been impossible. On the other hand the circumstances favoured slaughter at same age, as the number of stalls available in the cow stable was quite restricted, and the calves for the following experiments could only be taken in after the animals in the previous experiments had been removed. In addition carcass dissection would have spread over a much longer period which would have been quite difficult to arrange in the slaughterhouses.

The distribution of material by breed and experiment is presented in Tables 1 and 2.

Table 1. Distribution of material by experiment and breed. 1-year olds.

	S. W. Finland Experiment Station				N. Ostrobothnia Experiment Station		No. of animals
	I	II	III	IV	V	VI	
	<i>Breed</i>						
Ay.Ay	6	12	12	47	—	32	109
Fc.Fc	6	12	12	—	16	—	46
Ch.Ay	6	—	—	—	—	—	6
Ch.Fc	6	—	11	—	—	—	17
Fr.Fc	—	—	12	—	15	—	27
Total	24	24	47	47	31	32	205
Initial age, days	11	9	12	11	23	22	14.5
Deviation »	4.5	3.3	3.4	4.0	16.7	7.5	—
Slaughter age, days	340	356	366	357	338	339	351.3
Deviation »	2.4	10.1	3.7	6.2	23.0	10.6	—
Feeding group	1	2	5	2	6	4	—

In experiment II the animals were castrated at approx. 8 months and in experiment III at approx. 4 months.

Breeds: Ay = Ayrshire, Fc = Finncattle, Ch = Charolais, Fr = Friesian.

Table 2. Distribution of material by experiment and breed. $\frac{1}{2}$ -year olds.

	S. W. Finland Experiment Station				No. of animals
	VII	VIII	IX	X	
<i>Breed</i>					
Ay.Ay	6	14	10	14	44
Fc.Fc	6	14	10	14	44
Ch.Ay	6	14	10	13	43
Ch.Fc	6	14	10	13	43
Total	24	56	40	54	174
Initial age, days	15	11	10	8	10.4
Deviation »	6.6	5.8	8.8	2.2	—
Slaughter age, days	165	166	166	165	165.6
Deviation »	2.2	2.5	3.1	6.8	—
Feeding group	1	1	1	4	—

Breeds: Ay = Ayrshire, Fc = Finncattle, Ch = Charolais, Fr = Friesian.

Estimated traits and estimators

Estimated traits:

1. Live weight
2. Carcass weight
3. Quantities and percentages of carcass meat, fat and bone.

Estimators:

- I Live measurements
- II Initial weight and daily gain during experiment
- III Carcass quality scores
- IV Live weight, carcass weight, dressing-%, by-products and, only for the 1/2-years olds, cutting results of flanks
- V Cutting results of shanks
- VI Cutting results of fore and hind quarters
- VII Detailed cutting results of the carcasses

Presentation of the results

Because there are eight estimated traits and seven trait groups act as estimators the results are presented in the following order:

1. Definition and mean results of the estimated traits
2. Results obtained by each estimator (I–VII) are presented separately in the following order:
 - a) Definition and mean results of the traits
 - b) results of the stepwise regression analysis and discussion.

Statistical analysis

In order to minimize the errors caused by spurious correlations between traits, all the traits of each animal were corrected within experiment for age, feeding and breed.

In order to estimate the correction factors and to examine the statistical significance of the variation caused by different factors — age, feeding and breed — the results of each experiment were first analysed with the least squares method (HARVEY 1966).

In the analysis the slaughter age is the independent variant and the breeds and feeding groups form the levels. With the regression coefficient each trait was corrected for age thus obtained to correspond the average age in the experiment. In the same way each trait was corrected for breed (or sire) and feeding deviation within experiment.

All the corrections were made for each trait irrespective of whether the variation caused by the above mentioned factors was statistically significant or not.

Corrections were made according to the following equation:

$$y_0 = x_0 - b(a_0 - \bar{a}) - x_1 - x_2,$$

- where:
- y_0 = corrected value of a trait
 - x_0 = original, uncorrected value of a trait
 - b = regression coefficient of age for the trait
 - a_0 = slaughter age
 - \bar{a} = mean slaughter age in the experiment
 - x_1 = breed (or sire) deviation for the trait
 - x_2 = feeding deviation for the trait

In the stepwise regression analysis corrected values were used with which the best estimators (groups I—VII, page 393) to estimate the traits — live weight, carcass weight and carcass composition.

The steps were followed till the standard error of estimate started to increase, but only those steps (traits) are retained in the results which had a statistically significant value for the partial regression coefficient (b).

With stepwise regression analysis the correlations between different traits (r), multiple correlations (R), partial regression coefficients (b) and standardised partial regression coefficients (beta) could be calculated.

In discussion on the results it has to be considered that they were not analysed by breeds but the difference between breeds and also feedings are eliminated as carefully as possible with the correction factors obtained with the least squares method. It is impossible to check whether all the corrections are correct. Especially the effect of age on different traits within a certain time period can not be determined because the deviation of age has been relatively large as in experiment V.

Estimated traits

Live weight

In most experiments the live weight has been registered at the experiment station on the previous evening or the morning of transport to slaughterhouse. If the animals had been weighed on two or three consequent days before slaughter, mean of these has been used in the results. In all experiments the animals were transported to the slaughterhouse in the morning and slaughtered the same day. The animals were not on fast before slaughter.

Carcass weight

At the slaughterhouse the carcasses are weighed immediately after slaughter. An automatic scale reduces from this so called warm carcass weight 3 % to compensate for possible evaporation and drying losses. The weight thus obtained is carcass weight (= cold carcass weight) according to which the producer price is determined. All by-products, including kidney fats which in some countries remain in the carcass, are removed.

Quantities and percentages of carcass meat, fat and bone

Partition of different types of tissue

Carcass weight is registered before dissection. In average, the registered weight lies between warm and cold carcass weight, the first ones dissected closer to the warm carcass weight. The percentages for the cutting results were calculated from this so called actual weight. If calculated from carcass weight the first ones would have received cutting gain, and the latter ones cutting loss. In the results, however, only cold carcass weight is presented because presenting several weights for one carcass would disturb further treatment of

the results, especially as the percentage for by-products and the dressing-% are calculated from carcass weight according to present practice.

Dissection was performed according to the routine, the so called commercial cutting scheme, practised by the co-operative slaughterhouses. Carcasses from the S.W. Finland Experiment Station were dissected at the S.W. Finland Co-operative Slaughterhouse (LSO) in Turku and the ones from N. Ostrobothnian Experiment Station in slaughterhouse Lihakunta in Oulu.

Carcass cutting scheme is presented in Figure 1.

Carcasses in experiment VII and VIII were not dissected. In experiments I, IX and X the carcasses were dissected according to Figure 1., where the carcasses were cut into fore and hind quarters between the seventh and the eighth rib. Hereafter the fore quarters was partitioned into briskets, fore back, shoulders, neck and fore shanks, and the hind quarters into the rounds with hind shanks, back, flanks and short plates. All parts were cut into meat, fat, bone and tendon selections. The outer and inner fillets¹⁾ and steaks separately including and excluding bones were weighed.


Carcasses in experiments II—VI were also cut into fore and hind quarters as shown in Figure 1, but the quarters were partitioned no further but cut separately into tissue selections. However, in the hind quarters the outer fillets and the steaks including and excluding bones were weighed separately.

In experiments II—VI the fore and hind shank on the right hand side, and in experiments I, IX and X both fore shanks but neither hind shank were removed from the carcass. (An accurate description of shank dissection is presented in Chapter V.)

Although the cutting scheme is the same within the Co-operative Slaughterhouse Organisation, some differences appear in dissection technique. Except for shanks, dissection of the main cuts from the carcasses was the same in both slaughterhouses. In LSO meat was classified into Ist and IInd grades where the Ist one consisted almost entirely of meat and the IInd one included also some fats and fasciae. In Lihakunta, however, Ist, IInd and IIIRD grades were dissected.

In the present results all meat selections are combined making no distinction between grades.

Fig. 1.
CUTTING SCHEME OF
A NEAT CARCASS



¹⁾ outer fillet = *Musculus longissimus dorsi* — MLD

Means of estimated traits by experiment

The means and standard deviations of live weight, carcass weight and carcass cutting results are presented by experiment in Tables 3, 4 and 5. In case all the animals represent the same breed or have a similar diet, no remark about the statistical significance of breed or feeding is made. Only in those experiments where sons of several sires are tested and the difference between sires has been statistically significant, a remark is made. In tables 3—7, 11, 14, 19, 23, 26 and 27 the means and standard deviations are calculated from corrected values. The statistical significance of variation caused by breed, feeding and age are calculated from original values.

With 1-year olds (Table 3) the differences between live weight and carcass weight between breeds were significant in all experiments. Instead, with 1/2-year olds (Table 3) in experiments VIII and IX no statistically significant differences in live weight appeared between breeds, therefore the differences in carcass weights are due to the higher dressing-% of the crossbreds.

In quantities of meat and bone the differences between breeds (Table 4) were statistically significant with 1-year olds, whereas differences in percentages were significant only in a few cases. Also differences caused by age affected more clearly the quantities than the percentages.

In experiment IV (Tables 3 and 4) the difference caused by feeding was statistically significant with almost all the traits. During the indoor feeding season half of the animals were on abundant concentrate diet and the other half on silage fodder diet. During summertime one half were continuously

Table 3. Means and standard deviations of live and carcass weights by experiment.

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
1-year olds, n = 205										
	Live weight kg					Carcass weight kg				
I	328	23.1	*	—	—	168	12.3	***	—	—
II	298	17.8	***	—	—	140	10.5	***	—	—
III	336	23.9	**	***	**	158	12.2	**	***	**
IV	346	33.9 sires	*	***	—	158	18.0	***	—	—
V	289	28.0	*	—	**	134	14.2	*	—	*
VI	273	18.1	—	—	*	123	8.5	—	—	—
1/2-year olds, n = 174										
	Live weight kg					Carcass weight kg				
VII	203	16.4	**	—	—	100	8.1	***	—	—
VIII	200	16.2	—	—	*	98	9.1	*	—	*
IX	188	17.1	—	—	**	93	9.5	*	—	*
X	216	14.3	***	***	**	109	7.9	***	***	**

Statistical significance: *P < 0.05, **P < 0.01, ***P < 0.001.

indoor receiving abundantly concentrates and the other half were on mere pasture. During that period the growth rate of the two groups remained the same, difference in live weight had been gained already during indoor feeding period. However, animals in the first half carried about twice as much fats as in the second one.

With $\frac{1}{2}$ -year olds (Table 5) the differences were statistically significant in quantities and percentages of meat and bone. In these experiments there were in addition to the purebred Ay- and Fc-calves also Charolais crossbreds which in earlier experiments usually were thicker fleshed and less bony than the purebreds (RUOHOMÄKI and VARO 1967).

Table 4. Cutting results of whole carcass. Means and standard deviations by experiment. 1-year olds

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
n=205										
	Meat-kg					Meat-%				
I	129.7	10.3	***	—	—	76.3	1.2	—	—	—
II	104.8	8.5	***	—	—	73.8	0.9	—	—	—
III	118.2	10.1	**	***	**	73.7	1.5	—	—	—
IV	114.2	14.1		***	—	71.5	1.4		***	—
V	91.8	10.4	*	—	*	68.0	1.0	**	*	—
VI	93.2	6.9		—	—	75.4	1.0		—	—
	Fat-kg					Fat-%				
I	6.3	1.4	—	—	—	3.7	0.8	—	—	—
II	3.7	1.1	—	—	—	2.6	0.7	—	—	—
III	5.4	2.0	—	*	—	3.4	1.3	—	—	—
IV	9.6	2.8		***	—	5.6	1.2		***	—
V	13.6	2.1	—	*	*	10.1	1.3	*	*	—
VI	0.5	0.3		—	*	0.4	0.2		—	*
	Bone-kg					Bone-%				
I	30.9	2.3	*	—	—	18.2	0.8	**	—	—
II	32.6	3.3	***	—	—	21.1	0.8	—	—	—
III	32.5	2.9	—	—	**	20.3	1.4	—	—	—
IV	31.8	2.7 sires	**	—	—	20.2	1.6		***	—
V	27.9	2.9	—	—	*	20.7	0.9	—	**	—
VI	28.7	1.9		—	*	23.2	1.1		—	—
	Tendons %					Meat/Bone-ratio				
I	1.3	0.3	—	—	—	4.20	0.18	*	—	—
II	2.4	0.7	—	—	—	3.49	0.15	—	—	—
III	2.4	0.4	*	—	—	3.64	0.29	—	—	—
IV	2.5	0.3		—	—	3.57	0.31		***	—
V	1.0	0.1	—	*	—	3.25	0.22	*	**	—
VI	0.8	0.1		—	—	3.26	0.22		—	—

In all the tables the quantities are indicated: meat-kg, fat-kg, bone-kg and the percentages: meat-%, fat-% and bone-%.

Table 5. Cutting results of whole carcass. Means and standard deviations by experiment. $\frac{1}{2}$ -year olds

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
	Meat-kg					Meat-%				
IX	68.6	7.6	*		*	72.7	1.3	**		—
X	82.6	6.4	***	***	**	74.8	1.2	***	—	*
	Fat-kg					Fat-%				
IX	3.4	0.8	—		—	3.6	0.6	**		—
X	3.4	1.0	—	—	—	3.1	0.8	—	—	—
	Bone-kg					Bone-%				
IX	20.0	1.8	*		—	21.3	1.2	***		—
X	22.1	1.5	***	***	—	20.2	1.1	***	—	*
	Tendons %					Meat/Bone-ratio				
IX	2.3	0.4	—		—	3.43	0.30	**		—
X	1.7	0.4	—	—	—	3.74	0.23	***	—	—

Ist estimator: *live measurements*

Measuring technique


Before the measuring technique and results are presented some observations on the technique and errors by other researchers are explained.

According to WEBER (1957) the basic requirement for the measurements is that they are biologically reasonable. Regardless this were true, also the following sources of error were possible by Weber: 1) differences between animals, 2) differences between persons who perform the measuring, 3) differences between animals and persons and 4) errors in the measurements by one person. It was observed that no errors occur provided the head and feet remain in one position. The greatest difficulties and inaccuracies under the practical circumstances are caused by the movements of the animal.

In a study by MATHER et al. (1959) 99 heifers and 52 cows were measured twice and the differences in the measurements were compared. The difference was lowest in hip width, 0.5 % and the highest in the paunch girth, 5.8 %, except the body length of the heifers, 8.2 %. In a study by TOUCHBERRY and LUSH (1950) the greatest errors occurred in the measurements of length and the smallest in the depth of chest.

In the present study the measurements were taken of all the animals except the ones in experiments V and VI. Measurements were normally taken on

Fig. 2. Diagram of the measurements



the day of live weight determination, or, if the animal was weighed on several consequent days, on one of these occasions. All the measurements were performed by the same person who also during the growth of the animals had measured them several times. Thus it can be supposed that there appears a minimum of measurement errors in this material. Live animal measurements are presented in Figure 2.

Measurements 1–5 are taken with a measuring stick and 6, 7, 8, 10 and 13 with a measuring tape. Width of chest (3) has been measured on the back as the distance between the tips of the humeri and the width of back (4) as the distance between the hooks. Depth of chest (5) has been measured quite behind the front legs at the same point as the heart girth (6). Natural length (7) has been measured along the spine from withers till tail head, in the same way the oblique length (8) from the tip of humerus till tail head. The circumferences of front and hind cannon (10 and 13) are taken with a measuring tape at the narrowest part of the cannon (metacarpus and metatarsus). The widths of the cannons (11 and 14) are taken at the same point with a sliding gauge with which also the widths of knee (9) and hock (12) are measured.

Means of live measurements by experiment

In Tables 6 and 7 means and standard deviations of live measurements are presented by experiment.

The differencies in the widths of chest and back between breeds were statistically significant with both the 1-year olds (Table 6) and the $\frac{1}{2}$ -year olds (Table 7). On the other hand no equally distinct differencies in measurements of length and height appeared between breeds. Almost all differencies between breeds on measurements of legs were statistically significant, some of them even between sires.

Table 6. Means and standard deviations of live measurements by experiment.

1-year olds

n = 142

Epx. No.	Mean cm	SD cm	Statistical significance			Mean cm	SD cm	Statistical significance			
			Breed Feed. Age					Breed Feed. Age			
	1. Height at withers			2. Height of back							
I	111.1	3.4	—	—	—	116.0	3.8	—	—	—	
II	114.5	3.0	—	—	—	118.4	3.1	—	—	—	
III	118.0	2.9	—	*	**	123.8	2.9	—	*	*	
IV	115.0	3.2		*	—	119.2	3.5 sires	**	—	—	
	3. Width of chest			4. Width of back							
I	45.7	3.3	*		—	42.0	2.0	***		—	
II	38.3	2.8	***	—	—	38.0	2.4	***	—	—	
III	40.5	2.7	***	*	*	38.9	2.0	***	**	**	
IV	42.2	3.2		***	—	40.5	2.4		***	—	
	5. Depth of chest			6. Heart girth							
I	59.5	1.7	—		—	162.8	5.5	*		—	
II	59.3	1.5	—	—	—	155.7	4.1	*	—	—	
III	60.5	1.1	—	**	**	161.9	3.5	*	***	**	
IV	60.6	2.6 sires	*	**	—	162.5	6.6		***	—	
	7. Natural length			8. Oblique length							
I	114.6	5.0	—		—	129.7	4.7	—		—	
II	117.4	3.9	—	—	—	130.9	3.1	**	—	—	
III	117.9	4.0	—	*	**	131.4	4.1	—	—	**	
IV	116.6	3.1 sires	**	***	—	130.7	4.8		***	—	
	9. Width of knee			12. Width of hock							
I	9.4	0.4	*		—	8.7	0.3	**		—	
II	9.3	0.5	*	—	—	8.5	0.3	***	—	—	
III	9.7	0.4	*	***	***	8.9	0.3	***	—	*	
IV	9.6	0.4 sires	***	***	—	8.8	0.4 sires	***	—	—	
	10. Circumf. of front cannon			11. Width of front cannon							
I	17.2	0.9	*		—	6.0	0.2	***		—	
II	17.3	0.6	***	**	*	5.9	0.2	***	—	—	
III	18.4	0.8	***	**	*	6.1	0.2	**	*	**	
IV	18.1	0.8 sires	**	*	*	6.1	0.3 sires	***	**	—	
	13. Circumf. of hind cannon			14. Width of hind cannon							
I	19.3	0.8	*		—	7.3	0.3	**		—	
II	19.3	0.6	***	—	—	7.3	0.3	***	—	—	
III	20.3	0.9	***	**	*	7.5	0.3	***	**	**	
IV	19.9	0.8 sires	**	—	—	7.4	0.3 sires	***	—	—	

Table 7. Means and standard deviations of live measurements by experiment.

 $\frac{1}{2}$ -year olds

n = 174

Exp. No	Mean cm	SD cm	Statistical significance			Mean cm	SD cm	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
	1. Height at withers			2. Height of back						
VII	101.6	2.5	—	—	—	105.1	3.3	—	—	—
VIII	98.9	2.8	—	—	—	102.0	2.9	**	—	—
IX	98.2	3.2	—	—	—	103.0	3.5	—	—	—
X	100.9	2.8	*	*	*	106.0	3.2	*	**	—
	3. Width of chest			4. Width of back						
VII	33.0	1.8	**	—	—	30.8	0.9	***	—	—
VIII	31.7	1.9	***	—	—	31.0	1.6	***	—	—
IX	31.9	2.8	*	—	—	31.3	2.5	**	—	—
X	36.0	2.5	***	***	—	33.6	2.1	**	***	*
	5. Depth of chest			6. Heart girth						
VII	47.7	1.6	—	—	—	131.6	4.6	—	—	—
VIII	48.1	1.6	—	—	—	131.4	4.3	**	—	*
IX	49.0	1.9	—	—	—	131.5	4.4	—	—	*
X	50.1	2.1	—	***	*	137.5	4.7	*	***	*
	7. Natural length			8. Oblique length						
VII	101.5	3.0	—	—	—	115.1	2.6	—	—	—
VIII	100.6	4.1	***	—	—	112.2	4.3	***	—	—
IX	98.2	3.8	—	—	—	110.2	4.3	—	—	—
X	102.9	4.6	—	***	*	115.0	3.7	—	***	**
	9. Width of knee			12. Width of hock						
VIII	8.5	0.3	***	—	—	8.5	0.3	***	—	—
VIII	8.4	0.3	***	—	—	8.3	0.3	***	—	**
IX	8.4	0.4	**	—	—	8.2	0.3	***	—	—
X	8.8	0.4	***	*	—	8.3	0.4	***	**	—
	10. Circumf. of front cannon			11. Width of front cannon						
VII	15.5	0.7	**	—	—	5.2	0.2	**	—	—
VIII	15.4	0.7	***	—	—	5.1	0.2	***	—	—
IX	15.4	0.8	***	—	—	5.1	0.3	*	—	—
X	15.7	0.8	***	**	—	5.3	0.3	***	—	—
	13. Circumf. of hind cannon			14. Width of hind cannon						
VII	16.8	0.5	***	*	—	6.3	0.2	***	—	—
VIII	16.6	0.5	***	—	—	6.1	0.2	***	—	—
IX	17.1	0.9	***	—	—	6.2	0.3	—	—	—
X	17.6	0.8	***	***	—	6.5	0.3	***	**	—

According to a study by LINDSTRÖM and MAIJALA (1970) the following means of live measurements are presented of animals in a young bull rearing station:

Age 365 days	Ay 395 animals	Fc 179 animals
Heart girth	178 cm	173 cm
Natural length	133 »	128 »
Height at withers	119 »	118 »

As these figures are compared with the ones of the 1-year olds presented in Table 6 it can be seen that the mean of the heart girth is here about 15 cm and the length about 14 cm larger than in the present study. Difference in height was only 3 cm.

Stepwise regression analysis, results and discussion

Estimation of live weight, carcass weight and carcass composition with live measurements.

In Tables 8, 9 and 10 also live weight has been one of the estimating factors (part b.). Where no additional information is obtained with live measurements, the results are omitted from the tables.

Live weight. The R^2 ¹⁾ obtained for live weight with live measurements was 87 % for the 1-year olds and 82 % for the 1/2-year olds (Table 8, a). The most reliable estimators of live measurements for both age groups were the width of chest and the heart girth. With 1-year olds the second step was natural length but the value of its standardized regression coefficient was, however, lower than that of the heart girth and width of chest. With 1/2-year olds the natural length was replaced by the oblique length, and the height of back with height at withers.

The correlation of natural length on live weight was 0.659, and that of the oblique length 0.619 with the 1-year olds, with the 1/2-year olds the correlations were 0.510 and 0.546 respectively. The correlations of the height at withers and the height of back on live weight with 1-year olds were 0.542 and 0.598 and with 1/2-year olds 0.548 and 0.562 respectively. As the correlations of these measurements on live weight are almost equal, the reliability of the estimation would hardly decrease if the natural length was used instead of the oblique length and the height at withers instead of height of back. Application of fewer measurements would simplify the work and reduce errors.

Interrelationships between live weight and live measurements have been investigated in several studies. The most common measurements taken have been heart girth, width of chest, depth of chest, height at withers, height of back and natural length. In the more accurate definition of the measurement points, however, differences exist between different studies, which sometimes makes them incomparable. As materials consist of animals of different breeds and ages this naturally also reduces the possibilities of comparison.

¹⁾ R^2 = Coefficient of Determination, (%)

The R^2 obtained for live weight with heart girth and natural length was 79 % which is very close to the one obtained in the study by LINDSTRÖM and MAIJALA (1970) where the average R^2 -value obtained with the same traits was 81 % for both Ayrshire and Finncattle.

Table 8. Result of stepwise regression analysis.
 a) Live measurements — live weight and carcass weight.
 b) Live measurements and live weight — carcass weight.

Step	x	r	R	R^2	b	β
1-year olds, n = 142						
y = Live weight, $\bar{y} = 331$ kg						
a) 1.	Heart girth823	.823	67.7	1.928***	0.370
2.	Natural length659	.889	79.0	1.720***	0.250
3.	Width of chest790	.923	85.2	2.925***	0.325
4.	Width of hind cannon665	.929	86.3	12.066**	0.131
5.	Height of back598	.932	86.9	0.748*	0.091
y = Carcass weight, $\bar{y} = 156$ kg						
a) 1.	Heart girth806	.806	65.0	0.999***	0.360
2.	Width of chest797	.871	75.9	1.729***	0.361
3.	Natural length638	.910	82.8	0.933***	0.255
4.	Width of hind cannon660	.918	84.3	7.448***	0.152
b) 1.	Live weight938	.938	88.0	0.390***	0.734
2.	Width of back742	.943	88.9	0.823**	0.127
3.	Width of hock380	.947	89.7	3.661**	0.092
4.	Heart girth806	.948	89.9	0.280*	0.101
$1/2$ -year olds, n = 174						
y = Live weight, $\bar{y} = 203$ kg						
a) 1.	Heart girth778	.778	60.5	0.973***	0.275
2.	Width of chest752	.849	72.1	2.157***	0.317
3.	Height at withers548	.884	78.1	1.011***	0.181
4.	Width of hind cannon621	.895	80.1	9.402***	0.160
5.	Oblique length546	.903	81.5	0.585***	0.145
6.	Width of back681	.908	82.4	1.112**	0.135
y = Carcass weight, $\bar{y} = 100$ kg						
a) 1.	Heart girth779	.779	60.7	0.614***	0.318
2.	Width of chest764	.856	73.3	1.431***	0.385
3.	Natural length486	.878	77.1	0.305***	0.143
4.	Height at withers492	.885	78.3	0.349**	0.114
5.	Width of back687	.890	79.2	0.557*	0.123
6.	Width of knee514	.893	79.7	2.039*	0.086
b) 1.	Live weight922	.922	85.0	0.383***	0.700
2.	Width of chest764	.928	86.1	0.556***	0.149
3.	Heart girth779	.932	86.9	0.270**	0.140

Statistical significance: *P < 0.05, **P < 0.01, ***P < 0.001.

Table 9. Result of stepwise regression analysis.
 1-year olds a) Live measurements — carcass composition.

n = 142 b) Live measurements and live weight — carcass composition.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 116.6$ kg						
a) 1.	Width of chest796	.796	63.4	1.655***	0.431
2.	Heart girth783	.858	73.6	0.832***	0.374
3.	Natural length593	.886	78.5	0.732***	0.250
b) 1.	Live weight906	.906	82.1	0.345***	0.809
2.	Width of chest796	.915	83.7	0.687**	0.179
3.	Height of back459	.919	84.5	-0.394**	-0.113
4.	Width of hock357	.921	84.8	2.259*	0.070
y = Fat-kg, $\bar{y} = 6.6$ kg						
b) 1.	Depth of chest506	.506	25.6	0.288**	0.260
2.	Width of hock	-.182	.555	30.8	-1.999***	-0.338
3.	Live weight478	.625	39.1	0.032***	0.414
y = Bone-kg, $\bar{y} = 32.0$ kg						
a) 1.	Width of hind cannon643	.643	41.3	2.154**	0.220
2.	Natural length626	.735	54.0	0.183***	0.252
3.	Width of hock562	.768	59.0	2.044***	0.257
4.	Height of back574	.792	62.7	0.189***	0.219
5.	Heart girth518	.799	63.8	0.126*	0.132
b) 1.	Live weight671	.671	45.0	0.027**	0.262
2.	Width of hock562	.764	58.4	2.260***	0.284
3.	Height at withers560	.788	62.1	0.164**	0.179
4.	Width of hind cannon643	.797	63.5	1.665*	0.170
5.	Natural length626	.804	64.6	0.117*	0.161
y = Meat-%, $\bar{y} = 73.4$ %						
a) 1.	Width of chest337	.337	11.4	0.136**	0.302
2.	Height of back	-.117	.432	18.7	-0.139***	-0.340
3.	Heart girth319	.489	23.9	0.102***	0.394
4.	Width of hind cannon008	.515	26.5	-0.949*	-0.205
y = Fat-%, $\bar{y} = 4.0$ %						
a) 1.	Width of hock	-.294	.294	8.6	-0.965***	-0.319
2.	Depth of chest253	.406	16.5	0.159***	0.280
y = Bone-%, $\bar{y} = 20.1$ %						
a) 1.	Heart girth	-.455	.455	20.7	-0.108***	-0.430
2.	Width of hock273	.587	34.5	1.423***	0.394
3.	Width of chest	-.410	.631	39.8	-0.145***	-0.336
4.	Height of back014	.668	44.6	0.101***	0.258

Table 10. Result of stepwise regression analysis.
 $\frac{1}{2}$ -year olds a) Live measurements — carcass composition.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 76.6$ kg						
a) 1.	Width of chest795	.795	63.2	1.057***	0.406
2.	Heart girth740	.843	71.1	0.439***	0.288
3.	Natural length401	.871	75.9	0.288**	0.177
4.	Width of back748	.885	78.3	0.724**	0.236
b) 1.	Live weight882	.882	77.8	0.269***	0.602
2.	Width of chest795	.898	80.6	0.442*	0.170
3.	Width of hock333	.905	81.9	-2.736**	-0.140
4.	Heart girth740	.912	83.2	0.226*	0.148
5.	Width of back748	.917	84.1	0.461*	0.150
y = Fat-kg, $\bar{y} = 3.4$ kg						
a) 1.	Heart girth507	.507	25.7	0.064**	0.319
2.	Width of chest501	.556	30.9	0.108**	0.313
3.	Height at withers263	.572	32.7	0.102**	0.330
4.	Height of back128	.607	36.8	-0.080*	-0.289
y = Bone-kg, $\bar{y} = 21.2$ kg						
a) 1.	Width of knee730	.730	53.3	1.015**	0.249
2.	Depth of chest614	.801	64.2	0.169**	0.206
3.	Circumf. of hind cannon605	.832	69.2	0.389**	0.200
4.	Width of back581	.857	73.4	0.142**	0.196
5.	Width of hock692	.875	76.6	1.074**	0.233
6.	Oblique length508	.885	78.3	0.063**	0.153
b) 1.	Live weight796	.796	63.4	0.046***	0.440
2.	Width of knee730	.867	75.2	0.819**	0.201
3.	Width of hock692	.881	77.6	0.993**	0.216
4.	Height of back568	.892	79.6	0.068*	0.137
5.	Circumf. of hind cannon605	.898	80.6	0.263**	0.135
y = Meat-%, $\bar{y} = 73.9$ %						
a) 1.	Width of chest284	.284	8.1	0.188***	0.399
2.	Width of hock	-.232	.444	19.7	-1.270***	-0.360
y = Fat-%, $\bar{y} = 3.3$ %						
a) 1.	Heart girth305	.305	9.3	0.049**	0.305
y = Bone-%, $\bar{y} = 20.6$ %						
b) 1.	Width of chest	-.449	.449	20.2	-0.185***	-0.442
2.	Width of hock280	.635	40.3	1.482***	0.474
3.	Circumf. of hind cannon153	.668	44.6	0.397**	0.299
4.	Live weight	-.298	.694	48.2	-0.025*	-0.347

GRAVIR (1967) obtained an R^2 of 69 % for live weight with seven live measurements on 360-days old bulls, 57 % of this value was determined by heart girth alone.

In the study by SKJERVOLD (1958) several live measurements were taken. The correlations between six live measurements and live weight varied within 0.70–0.75.

In the last two studies the R^2 and the correlations remained somewhat lower than in the present study.

On the other hand, the results obtained in this study were on the same level as in the study by HUTH (1965), where the material consisted of 211 1-year old bulls. The following correlations were obtained:

Live weight — height of back and withers, mean	$r = 0.631$
— depth of chest	$r = 0.751$
— heart girth	$r = 0.859$
— width of back	$r = 0.836$

Correlations on the measurements of width are higher than on those of height.

In the study by JOHANSSON and HILDEMAN (1953) the heart girth, width of hips, height at withers, natural length and hind girth were measured on 10 000 animals. For 1175 $\frac{1}{2}$ -year olds and 171 young steers the correlations heart girth by live weight were 0.892–0.951. No additional information was obtained with other measurements.

The values varied even within one breed depending on the material, for the big animals the values were higher than for the small. In estimation of weight by heart girth the standard error was ± 6 % in live weight and ± 9 % in carcass weight. In weighing of live animals the standard error varied between ± 1 and ± 2 %.

For carcass weight (Table 8, a) the R^2 remained nearly 3 % lower for both age groups than for live weight. In both age groups the highest estimates were provided by width of chest, heart girth and natural length.

In the earlier mentioned study by GRAVIR an R^2 of 65 % for carcass weight was obtained with several live measurements. In the study by JOHANSSON and HILDEMAN the correlation between heart girth and carcass weight varied between 0.857–0.945.

In a study by BUSS (1968) the correlations of heart girth and carcass weight laid between 0.661 and 0.325.

In a study by LINDHÉ (1968) the material consisted of 218 animals, steers of six different breeds and crossbreeds. The regression between heart girth and carcass weight varied by breed between 2.67 and 3.36 (SE 9.3–16.1 kg), and between heart girth and live weight from 4.83 to 6.46 (SE 19.5–29.3 kg).

In several studies the correlations between live weight and carcass weight have been estimated. The correlations obtained by JOHANSSON and HILDEMAN were from 0.880 to 0.968 with $\frac{1}{2}$ -year olds and with young steers. BUSS obtained correlations of 0.783, 0.664 and 0.636 for the three breeds respectively. SEEBECK and TULLOH (1966) also observed in their study that carcass weight

was directly dependent on live weight, and that dressing-% increases with the live weight.

In the present study the correlations between live weight and carcass weight were with the measured 1-year olds 0.938 (with all the 1-year olds 0.940) and with $\frac{1}{2}$ -year olds 0.922 (with the dissected $\frac{1}{2}$ -year olds 0.924).

If it is possible to determine live weight through weighing, this alone provides a higher R^2 for carcass weight than live measurements. An R^2 of 88 % was obtained for 1-year olds and 85 % for $\frac{1}{2}$ -year olds (Table 8, b). In both age groups the live measurements increased the R^2 obtained with live weight only by about 2 %-units.

When carcass weight was estimated with mere live weight, the value of the regression coefficient (b) for both age groups was 0.50, thus if the live weight increases by one kilo, the carcass weight increases by 500 grams.

Like in the estimation of carcass weight the first three estimators for meat-kg were heart girth, width of chest and natural length for the 1-year olds (Table 9, a). The same value for R^2 for the $\frac{1}{2}$ -year olds, 78 %, was obtained with the above mentioned measurements and with the width of back (Table 10, a).


If the live weight is known in the estimation of meat-kg this alone provides an R^2 of 82 % for the 1-year olds (Table 9, b) and 78 % for the $\frac{1}{2}$ -year olds (Table 10, b). If the measurements are also included this gives additional information from 3 to 6 %-units.

In the study by VARO (1969) the material consisted of purebred Ayrshire and Finncattle of varying ages together with Charolais crosses. Among young animals the carcass meat quantity was best estimated by the ratio of carcass weight and size (= total of 10 measurements), the R was 0.949. When the older animals were concerned the best estimator of meat quantity was carcass weight, $r = 0.97$.

Foreign results are seldom comparable to the present ones because cutting practices in different countries differ from each other. Differences are apparent in comparison of the Finnish (Fig. 1) and the American (Fig. 3) carcass cutting schemes (LAINE 1971). Also other cutting schemes from other countries could be included as all of them differ from each other, and from the one used in Finland, to an extent which makes comparison almost impossible.

In several studies the quantities and percentages of the primal cuts in the whole carcass are investigated. In comparison of the results the differences in partitioning are of no significance when the quantities

Fig. 3.
WHOLESALE CUTS IN THE
U.S.A.



and percentages of meat, fat and bone of the whole carcass are compared.

Of the other results should also be mentioned those by SKJERVOLD (1958). With height over hips and width of rounds a correlation of $R = 0.68$ was obtained for the hind quarter, which was separated from the right hand side half between the 13th rib and the 1st lumbar vertebra. In addition to this when six measurements were included the correlation rose to $R = 0.71$. On the weight of the fore quarter the correlations were lower than the ones mentioned above.

The material by BUSCH et al. (1969) consisted of 181 steers. The mean yield of edible meat from one half of a carcass was 85.6 kg. The correlations of girth measurements on the quantity of edible meat remained between 0.41 and 0.81, and on the measurements of width 0.42—0.70, and on the measurements of length, 0.62 respectively. The correlation on carcass weight was 0.94. In the whole material (745 animals) carcass weight estimated the quantity of edible meat till 75—88 %, the measurements increased the coefficient by 2—4 %-units.

In the present study the R^2 for fat - kg of 39 % was obtained for the 1-year olds (Table 9, b) and of 37 % for the $\frac{1}{2}$ -year olds (Table 10, a).

In estimation of bone - kg the R^2 of 78 % was obtained for the $\frac{1}{2}$ -year olds (Table 10, a) and 64 % for the 1-year olds (Table 9, a).

For both the age groups several measurements were required to obtain the above mentioned values for R^2 . An accurate determination of the measuring sites cause difficulties in practice.

In both age groups rather low values were obtained for R^2 on meat - % and fat - % (Tables 9 and 10). For bone - % an R^2 of 45 % was obtained for both age groups.

In the study by NEIMAN—SØRENSEN et al. (1965) eight live measurements were taken. The results are presented together with those by NIELSEN (1962) obtained with 243 calves of the same breed.

Neiman—Sørensen:	Meat-%	Fat-%	Bone-%
R^2 (%)	6	2.5	13
Nielsen:			
Width of back, $r =$	-0.038	0.081	-0.041
Width of rounds $r =$	0.046	-0.011	-0.050

The correlations above are considerably lower than the ones obtained in the present study.

In the study by TALLIS et al. (1959) the obtained correlations for edible meat-% were: weight/height $r = 0.45$ and weight/width $r = -0.43$.

In the study by ORME et al. (1959 c) the correlations between carcass girths and the percentages of the primal cuts varied between -0.42 — -0.53 .

Ind estimator: initial weight and daily gain

Initial weight (weight at the beginning of the experiment)

The calves were weighed the day they were brought to the experiment station. Weights at birth are not known. Initial weights have been corrected by experiment for initial age unlike the other traits which were corrected for slaughter age.

Daily gain during experiment gr./day.

Daily gain was calculated according to the following model:

$$\frac{\text{Live weight at the end of exp.} - \text{Initial weight}}{\text{Slaughter age} - \text{Initial age}}$$

In calculations, no other figures for growth have been employed but the one for growth during experiment, because some calves were born in spring and the others in autumn. In the experiments the calves were of different ages at the first indoor feeding and pasture period. The calves born in spring started pasturing at about 3 months and the ones born in autumn at about 8 months.

Means of initial weights and daily gains by experiment

Table 11 presents the means and standard deviations of initial weights and daily gains by experiment.

Daily gain of the $\frac{1}{2}$ -year olds was 200 gr. higher than that of the 1-year olds. In the experiments for the $\frac{1}{2}$ -year olds all the calves were born in autumn

Table 11. Means and standard deviations of initial weight and daily gain by experiment.

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
1-year olds, n = 205										
Initial weight kg.										
I	42.1	4.7	—	—	—	873	69	—	—	—
II	38.1	2.8	**	—	—	749	47	***	—	—
III	39.7	3.9	—	—	***	840	62	*	***	*
IV	42.1	4.4 sires	***	—	**	877	96	***	—	—
V	43.3	8.7	—	—	*	779	83	*	—	—
VI	43.2	4.6	—	—	*	728	60	—	—	—
$\frac{1}{2}$ -year olds, n = 94										
Initial weight kg.										
Daily gain during experiment gr./day										
IX	40.1	4.0	***	—	***	940	96	—	—	—
X	41.7	4.5	***	—	—	1104	95	*	***	—

and received ample portions of fodder under almost uniform circumstances at the S.W. Finland Experiment Station. The experiments for the 1-year olds consisted of both steers and bulls. In several of them growth during the pasture period was low.

Stepwise regression analysis, results and discussion

Estimation of carcass weight and carcass composition with initial weight and daily gain.

For carcass weight the R^2 obtained was 86 % for the 1-year olds and 77 % for the $1/2$ -year olds. The respective values for meat - kg were 79 % and 71 % (Table 12). The R^2 obtained for fat - kg was 24 % for the 1-year olds and 29 % for the $1/2$ -year olds. The R^2 for bone - kg were 52 % and 56 % respectively. (Table 12).

Table. 12. Result of stepwise regression analysis. Initial weight and daily gain during experiment — carcass weight and carcass composition.

Step	x	r	R	R^2	b	β
1-year olds, n = 205						
y = Carcass weight ¹⁾ , $\bar{y} = 156$ kg.						
1.	Daily gain, gr./day907	.907	82.3	0.163***	0.857
2.	Initial weight, kg.410	.929	86.3	0.735***	0.208
y = Meat-kg, $\bar{y} = 109$ kg						
1.	Daily gain, gr./day884	.884	78.1	0.125***	0.864
2.	Initial weight, kg277	.889	79.0	0.206**	0.098
y = Fat-kg, $\bar{y} = 6.7$ kg						
1.	Daily gain, gr./day492	.492	24.1	0.012***	0.492
y = Bone-kg, $\bar{y} = 30.9$ kg						
1.	Daily gain, gr./day667	.667	44.5	0.022***	0.610
2.	Initial weight, kg401	.719	51.7	0.146***	0.275
$1/2$ -year olds, n = 94						
y = Carcass weight, $\bar{y} = 102$ kg.						
1.	Daily gain, gr./day840	.840	70.6	0.071***	0.786
2.	Initial weight, kg428	.880	77.4	0.537***	0.263
y = Meat-kg, $\bar{y} = 76.6$ kg						
1.	Daily gain, gr./day802	.802	64.3	0.055***	0.749
2.	Initial weight, kg.417	.843	71.1	0.429***	0.263
y = Fat-kg, $\bar{y} = 3.4$ kg						
1.	Daily gain, gr./day542	.542	29.4	0.005***	0.542
y = Bone-kg, $\bar{y} = 21.2$ kg						
1.	Daily gain, gr./day707	.707	50.0	0.012***	0.656
2.	Initial weight, kg383	.748	56.0	0.096***	0.249

¹⁾ n = 142

The correlations and regressions on the percentages of meat, fat and bone were quite low and therefore excluded from the calculations.

Daily gain was the first step in the estimation of all the traits. The correlations of initial age on all the traits was low and negative.

The different stages of growth have been abundantly studied in studies on both breeding and management. Determination of optimal growth is in key position considering the profitability of rearing beef cattle, as differences between breeds and management and the genetic variation have a decisive effect on the final result.

Only a few results of interrelations between the traits in question are presented here, as in the present study no other measurements for growth have been used but daily gain during experiment.

In the study by LINDSTRÖM and MAIJALA (1970) the following correlations were obtained between the weight at birth, at 180 days and at 365 days.

Weight at birth	Ay 335 animals		Fc 200 animals	
	r	r ²	r	r ²
— 180 days	0.608	37 %	0.588	34 %
— 365 days	0.537	29 %	0.413	17 %
	301 animals		142 animals	

The relation between birth weight and 180-day weight is considerably stronger than between birth weight and 365-day weight. The correlation between growth from 60-days to 180-days, and from 180-days to 365-days is nonsignificant.

In the present study the correlations between initial weights and live weights are considerably lower than the ones obtained by LINDSTRÖM and MAIJALA between weight at birth and live weight. With initial weight an R² of only 16 % on 165-day age, and of 11 % on 1-year were obtained. The mean initial age for the 1-year olds was 11 days and for the 1/2-year olds 9 days.

In Table 13 some results on the interrelation between growth and various carcass traits found in the literature are presented with the results of present study.

When the correlations between growth and the other traits from present study are compared to the ones presented in other studies it can be noticed that the correlations for growth found in this study especially on quantities are higher than the ones found in literature.

IIIrd estimator: carcass quality scores

Evaluation technique

The carcasses were evaluated after ca. 24 hours from slaughter. One person, who was uninformed of the breed or feeding group of the carcass, evaluated all the carcasses. The estimation was performed according to the practice in the slaughterhouse organisation. [The procedure has been applied since 1960, and the Meat Industrial Research Centre takes care of its further

Table 13. The correlations of daily gain on live and carcass weight and carcass composition found in the literature.

	BLACKWELL et al. 1962	du BOSE et al. 1967	BRACKELBERG & WILLHAM 1968	DIETERT 1969	HINKS & BECH ANDERSEN 1968	MARTIN & STARKENBURG 1965	NIELSEN 1962	SWIGER et al. 1965	WISMER-PEDERSEN 1969	Present study	1-yr. ½-yr.
Live weight76	.35	-	-	-	-	-	.75	-	.976	.925
Carcass weight72	.44	-	-	-	-	-	-	-	.907	.840
Boneless steak and roast kg	-	.45	-	-	-	-	-	-	-	-	-
Boneless steak and roast %	-	.11	-	-	-	-	-	-	-	-	-
Primal yield %	-	-	-	.15	-	-	-	-	-	-	-
Boneless retail cut kg	-	-	-	-	-	-	-	.65	-	-	-
Boneless retail cut %	-	-	-	-	-	-	-	-.35	-	-	-
Meat-kg	-	-	-	-	-	-	-	-	-	.884	.802
Meat-%	-	-	-.21	-	.0	-.04	.136	-	.0-	.287	.182
									.23		
Fat-kg	-	-	-	-	-	-	-	.36	-	.492	.542
Fat-%	-	-	.23	-	.06	-	.058	-	-.37	.127	.321
									-.10		
Bone-kg	-	-	-	-	-	-	-	-	-	.667	.707
Bone-%	-	-	-.21	-	-.05	-	-.252	-	-	-.395	-.305

development. The last slight changes were made in 1969. Since 1972 a new evaluation technique has been employed which partly differs from the one described here.

In evaluation of carcass fleshiness attention is paid to the following parts: outer, inner and overall steak lines, rump, sirloin, fore back, neck, shoulders, brisket + flanks and hind shanks. The mean of the points given to these ten parts is presented as fleshiness scores graded from 1 to 15, like each individual score of fleshiness given to an evaluated part in a carcass.

The scale for fat content and distribution was from - 7 till + 4, where points below one indicated a too high fat content.

The colours of meat and fat are estimated both separately from - 3 to + 5. Colour points are means of colour points for fat and meat.

Fleshiness scores + fat content + colour points build up the total score according to which the carcass quality class is determined.

The total scores correspond to the carcass quality classes as follows:

22 points	class	E (10)
20 »	»	I + (9)
18 »	»	I (8)
16 »	»	I - (7)
14 »	»	II + (6)
12 »	»	II (5)
10 »	»	II - (4)

In brackets is given the numerical value used in punch cards, which are also used in tables to indicate carcass quality class.

Steers, young bulls and ½-year olds are evaluated according to the same scale. Some carcasses, however, were so scarce in fat that fat colour could not be evaluated, thus it was excluded from all results. Still, it is included in colour points as the mean of fat and meat colour.

Means of carcass quality scores by experiment

Table 14 presents the means and standard deviations of carcass quality scores by experiment.

The carcass evaluation system has been especially criticized as it follows the shape of the carcass but provides no information of the quantities and percentages of different types of tissue. Also in the present material, differences between breeds are largely caused by differences in shape as carcasses of crossbred animals differ from the purebreds to an extent which affects especially the evaluation of the hind quarters. On the other hand crossbred carcasses contained more meat than purebreds, but the Finncattle carcasses generally obtained the lowest scores because of the narrow shape of their hind quarters, although their meat content usually was as high as that of the Ayrshires (RUOHOMÄKI 1967, RUOHOMÄKI and VARO 1967).

Stepwise regression analysis, results and discussion

Estimation of carcass composition with carcass quality scores.

In addition to quality scores also carcass weight has acted as one estimator of other carcass traits (Tables 15 and 16).

With carcass quality scores for meat-kg the R^2 obtained was 47 % for the 1-year olds (Table 15, a) and 59 % for the ½-year olds (Table 16, a).

When carcass weight was taken as an estimator of meat-kg in addition to the carcass quality scores this alone improved the value of R^2 up to 96 % in both age groups. Quality scores increased the R^2 by only 0.6 %-units for the 1-year olds (Table 15, b), and 1% -unit for the ½-year olds (Table 16, b). The standardized regression coefficient was for the 1-year olds about 20 times and for the ½-year olds about 10 times as high as that of the scores.

When carcass weight alone estimated the meat-kg, the value of the regression coefficient (b) was 0.7757*** for the 1-year olds, and 0.7960*** for the ½-year olds. It can be seen that if carcass weight increases by one kilo the meat-kg increases by about 800 grams. Carcass quality scores have no significance in estimation of meat-kg if carcass weight is known.

Table 17 presents the results of the stepwise regression analysis for the 1-year olds by steps when meat-kg has been estimated with carcass weight and carcass quality scores.

The R^2 increases only by 0.6 % with the scores of shoulders and not at all with the fleshiness scores. The standard error of the estimates (SE \pm 2.053)

Table 14. Means and standard deviations of carcass quality scores by experiment.

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
1-year olds, n = 205										
Fleshiness scores						Fat content and distribution				
I	13.4	0.5	***	—	—	3.8	0.4	—	—	—
II	9.9	1.2	**	—	—	2.8	0.6	*	—	—
III	10.6	1.0	***	**	—	3.4	0.5	—	—	—
IV	11.6	1.0	—	***	—	3.0	0.7	—	—	—
V	10.0	1.1	**	*	—	3.5	0.4	—	—	—
VI	10.2	1.1	—	—	—	2.9	0.3	—	—	—
Colour of meat						Colour points total				
I	4.0	0.8	*	—	—	4.4	0.7	—	—	—
II	3.3	0.7	*	—	—	3.3	0.7	*	—	—
III	4.0	0.8	—	—	—	4.1	0.8	—	—	—
IV	3.7	0.6	—	***	—	3.8	0.5	—	***	—
V	3.9	0.6	—	—	—	4.5	0.3	—	—	—
VI	3.0	0.3	—	—	—	3.1	0.3	—	—	—
Points total						1) Carcass class				
I	21.6	1.0	***	—	—	9.5	0.4	***	—	—
II	16.0	2.2	**	—	—	7.0	1.1	**	—	—
III	18.1	1.8	***	*	—	8.1	0.9	***	—	—
IV	18.3	1.7	—	***	—	8.3	0.8	—	***	—
V	18.0	1.2	***	**	—	7.9	0.7	**	*	—
VI	16.1	1.4	—	—	—	7.4	0.8	—	—	—
1/2-year olds, n = 94										
Fleshiness scores						Fat content and distribution				
IX	10.9	1.2	***	—	—	3.7	0.5	—	—	—
X	12.2	1.1	***	—	—	3.7	0.5	—	—	—
Colour of meat						Colour points total				
IX	4.7	0.5	—	—	—	4.8	0.5	—	—	—
X	4.7	0.7	—	—	—	4.8	0.4	—	—	*
Points total						1) Carcass class				
IX	19.3	1.7	***	—	—	8.5	0.9	***	—	—
X	20.8	1.5	***	—	*	9.1	0.8	***	—	*

1) Value 7 corresponds to carcass class I—, value 8 to carcass class I and value 9 to carcass class I+.

Table 15. Result of stepwise regression analysis.

1-year olds a) Carcass quality scores — carcass composition.

n = 205 b) Carcass quality scores and carcass weight — carcass composition.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 109$ kg						
a) 1.	Fleshiness scores686	.686	47.1	7.158***	0.686
b) 1.	Carcass weight980	.980	96.0	0.738***	0.933
2.	Shoulders430	.983	96.6	0.312*	0.045
3.	Fleshiness scores686	.983	96.6	0.474*	0.045
y = Fat-kg, $\bar{y} = 6.7$ kg						
a) 1.	Colour of meat290	.290	8.4	0.742***	0.259
2.	Fleshiness scores266	.370	13.7	1.029***	0.542
3.	Hind shanks116	.391	15.3	-0.307*	-0.234
4.	Shoulders103	.415	17.2	-0.240*	-0.193
b) 1.	Carcass weight484	.484	23.4	0.084***	0.586
2.	Colour of meat290	.556	30.9	0.831***	0.290
3.	Hind shanks116	.585	34.2	-0.283**	-0.215
y = Bone-kg, $\bar{y} = 30.9$ kg						
a) 1.	Hind shanks342	.342	11.7	0.490***	0.267
2.	Outer steak line341	.386	14.9	0.553***	0.298
3.	Colour of meat	-.141	.430	18.5	-0.735**	-0.183
4.	Shoulders083	.432	18.7	-0.294*	-0.168
b) 1.	Carcass weight731	.731	53.4	0.177***	0.884
2.	Fleshiness scores314	.762	58.1	-1.453***	-0.548
3.	Colour of meat	-.141	.774	59.9	-0.562**	-0.140
4.	Hind shanks342	.784	61.5	0.373**	0.203
5.	Outer steak line341	.793	62.9	0.360**	0.193
y = Meat-%, $\bar{y} = 72.9$ %						
a) 1.	Fleshiness scores499	.499	24.9	0.455***	0.374
2.	Shoulders451	.520	27.0	0.167*	0.210
3.	Colour points total	-.075	.541	29.3	-0.313*	-0.150
y = Fat-%, $\bar{y} = 4.4$ %						
b) 1.	Colour points total332	.332	11.0	0.604***	0.350
2.	Hind shanks	-.085	.352	12.4	-0.154**	-0.222
3.	Carcass weight088	.390	15.2	0.014*	0.196
y = Bone-%, $\bar{y} = 20.7$ %						
b) 1.	Fleshiness scores	-.555	.555	30.8	-1.084***	-0.916
2.	Colour of meat	-.300	.599	35.9	-0.414***	-0.232
3.	Hind shanks	-.319	.619	38.3	0.232***	0.283
4.	Outer steak line	-.342	.637	40.6	0.245***	0.296
5.	Carcass weight	-.446	.654	42.8	-0.018**	-0.204
6.	Brisket and flank	-.261	.663	44.0	0.122*	0.140

Table 16. Result of stepwise regression analysis.

 $\frac{1}{2}$ -year olds a) Carcass quality scores — carcass composition.

n = 94 b) Carcass quality scores and carcass weight — carcass composition.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 76.6$ kg						
a) 1.	Fleshiness scores718	.718	51.6	4.714***	0.763
2.	Overall steak line635	.738	54.5	0.989*	0.208
3.	Sirloin222	.753	56.7	-1.055*	-0.178
4.	Rump314	.765	58.5	-0.862*	-0.177
b) 1.	Carcass weight982	.982	96.4	0.734***	0.917
2.	Fleshiness scores718	.985	97.0	0.649**	0.105
3.	Neck528	.986	97.2	0.239*	0.055
4.	Carcass quality class585	.987	97.4	-0.466*	-0.055
y = Fat-kg, $\bar{y} = 3.4$ kg						
a) 1.	Points total435	.435	18.9	0.559***	0.945
2.	Carcass quality class308	.490	24.0	-0.621*	-0.556
y = Bone-kg, $\bar{y} = 21.2$ kg						
a) 1.	Hind shanks337	.337	11.4	0.415**	0.337
b) 1.	Carcass weight773	.773	59.8	0.199***	1.041
2.	Fore back089	.844	71.2	-0.326***	-0.283
3.	Shoulders199	.858	73.6	-0.197**	-0.218
y = Meat-%, $\bar{y} = 73.9$ %						
a) 1.	Fleshiness scores560	.560	31.4	0.431**	0.385
2.	Overall steak line519	.585	34.2	0.210*	0.243
y = Fat-%, $\bar{y} = 3.3$ %						
a) 1.	Points total274	.274	7.5	0.352**	0.748
2.	Carcass quality class167	.344	11.8	-0.460*	-0.518
y = Bone-%, $\bar{y} = 20.6$ %						
a) 1.	Fleshiness scores	-.676	.676	45.7	-0.277*	-0.279
2.	Fore back	-.675	.720	51.8	-0.243**	-0.310
3.	Neck	-.636	.735	54.0	-0.154*	-0.222

is in the first step ($3 \times \text{SE}$) 12.318 kg and in the third step correspondingly 11.562 kg. The remaining difference is only about 0.750 kg, therefore no significant improvement appears in carcass evaluation when carcass quality scores are included (Table 17).

In estimation of fat-kg the significance of the scores was quite low (Tables 15 and 16). The correlation of fat content scores on fat-kg was 0.065 for the 1-year olds and 0.308 for the $\frac{1}{2}$ -year olds.

With carcass quality scores an R² of 11–19 % was obtained on bone-kg (Tables 15, a and 16, a). When the carcass weight was included the value

Table 17. Result of stepwise regression analysis.
 1-year olds Carcass quality scores and carcass weight — meat-kg.
 n = 205

Step	x	r	R	R ²	b	β
y = Meat-kg, y = 109 kg						
1. Carcass weight980	.980	96.0	0.7757***	0.9809
standard error of estimation ± 2.053 kg						
1. Carcass weight980	.980	96.0	0.7750***	0.9547
2. Shoulders430	.983	96.6	0.4741***	0.0961
standard error of estimation ± 1.943 kg						
1. Carcass weight980	.980	96.0	0.7385***	0.9339
2. Shoulders430	.983	96.6	0.3125**	0.0455
3. Fleshiness scores686	.983	96.6	0.4740*	0.0455
standard error of estimation ± 1.927 kg						

increased to 63 % with the 1-year olds (Table 15, b) and to 74 % with the ½-year olds (Table 16, b).

In estimation of meat - % an R² of 19 % was obtained with 1-year olds and 34 % with ½-year olds (Tables 15 and 16). In estimation of fat - % R²-values of 12 % and 15 % were obtained respectively (Tables 15 and 16). For bone % an R² of 44 % for the 1-year olds and 54 % for the ½-year olds were obtained (Tables 15 and 16).

Carcass quality scores are particularly developed to serve commercial purposes. The price for meat is in each country determined by consumption habits which are different from country to country and in different times. For this reason the evaluation methods differ to such a great extent from each other. The common factor among them is, however, that they take into consideration carcass weight, the age and sex of the animal, meat and fat content and often also the colour of meat and fat.

As several weaknesses have been found in the evaluation methods they are being developed to meet their purpose better. Evaluation is sense-dependent and therefore also the possibility of misevaluation is greater than in methods based on measurements and weights. During the last few years in the newly applied methods also some measurements are included together with the scores. For example, measurement of the thickness of fat cover area of the MLD and weight of the intestinal fats.

It is found in several investigations that carcass evaluation based on mere sense-dependent evaluation meets the requirements of scientific study no better than live animal evaluation.

Correlations collected from the literature between carcass quality scores and different carcass traits are presented in Table 18. Also correlations obtained in present study are included.

Table 18. Correlations between carcass quality scores and carcass weight and composition found in the literature.

	Fleshiness scores		Fat content and distribution	Points total
	1)	2)		
Carcass weight				
BRÄNNÄNG 196642	— .45		.42
CUNDIFF et al. 196416
WILSON et al. 196415
Present study	1-yr.	.652	.141	.478
	1/2-yr.	.628	.289	.593
Fat-kg				
GOTTSCH et al. 196175	
SWIGER et al. 196531
VARO 1969	R =		.518	
Present study	1-yr.	.266	.065	.284
	1/2-yr.	.360	.308	.435
Boneless steak and roast %				
du BOSE et al. 1967				— .16
Primal yield %				
DIETERT 196915
Retail cut %				
CUNDIFF et al. 1964				— .34
SWIGER at al. 1965				— .44
Meat-%				
HINKS & BECH ANDERSEN 1968	.28	.44	— .24	.35
MARTIN & STARKENBURG 1965 .				— .26
NEIMAN-SØRENSEN et al. 196536	.34	— .13	.29
NIELSEN 1962359	.341		.291
Present study	1-yr.	.499	.062	.318
	1/2-yr.	.560	.123	.467
Fat-%				
MARTIN & STARKENBURG 1965 ...				— .16
NEIMAN-SØRENSEN et al. 196507	— .09	.35	.002
NIELSEN 1962	— .069	— .094		.014
Present study	1-yr.	.036	.070	.158
	1/2-yr.	.176	.262	.274
Bone-%				
NEIMAN-SØRENSEN et al. 1965 ...	— .41	— .36	— .23	— .37
NIELSEN 1962	— .408	— .360		— .414
Present study	1-yr.	— .555	— .145	— .499
	1/2-yr.	— .676	— .225	— .622

1) for the fore quarter, 2) for the hind quarter except in present study for both quarters.

When the correlations in Table 18 are compared with each other it can be found that they largely vary between studies and even within present study. Especially correlations on fat content with 1-year olds are dissimilar to the ones with $\frac{1}{2}$ -year olds.

Generally taking interrelations are weak, according to some studies even nonsignificant, as the most accurate information on meat-kg is provided by carcass weight and in estimation of other traits the scores provide only low accuracies. As already stated one of the weakest points in evaluation is that it follows the shape of the carcass. The shape is also affected by the dimensions of skeleton as SKJERVOLD (1958) mentions in his study.

According to a study by HAMMOND (1958) the short and plump animals have thicker muscles than the long and narrow ones. Thickly fleshed carcasses endure keeping better and processed meat is also tastier than meat obtained from thinly fleshed carcasses.

Also WITT (1965) in his study presents that the smaller surface area compared to the volume of the muscle the more economical and tastier the meat is after processing. Also de BOER et al. (1969) and PIRCHNER (1965) have expressed similar opinions. Meat-kg is not only a quantitative but also a qualitative trait as with the increase in meat content the bone content decreases and larger muscles are more valuable than the small ones.

Also results of the present study and earlier ones with Charolais crossbreds have shown that they comprise more meat and less bone than purebreds. They obtain higher quality scores, too, which proves that at least in these breed comparisons the plump shape of the carcass has indicated higher meat content (RUOHOMÄKI 1967, RUOHOMÄKI and VARO 1967).

As a conclusion it can be stated that information of carcass traits obtained with carcass quality scores is inadequate and that beside sense-dependent evaluation or instead of that an evaluation method which is entirely based on measurements and weights should be developed.

The evaluation method applied since the beginning of 1972 can not be compared to the present one as no carcass dissection was performed in the last experiments.

IVth estimator: *live weight, carcass weight, dressing-%, by-products and, only for the 1½-year olds, cutting results of flanks*

Determination of live weight and carcass weight has already been explained with the estimated traits. In the present chapter they act as estimators with dressing-%, by-products and flank cutting results.

Dressing-%

Dressing-% (carcass yield) is calculated as the ratio of carcass weight to live weight. In several studies the dressing-% has been calculated of the warm carcass weight which often makes it higher in foreign studies. Kidney fats which remain in the carcass for their part also increase the dressing-%.

By-products

In connection with slaughter the head, tongue, lungs, liver, heart, diaphragm, bowel and kidney fats, kidneys, spleen, skin and thyroid gland were weighed in most experiments. In calculations, however, only the head, liver, heart and kidney fats were included. In experiments I—IV, IX and X also bowel fats were weighed. The percentage of by-products was calculated from carcass weight.

Dissection of flanks

The flanks were dissected from the carcasses of the $\frac{1}{2}$ -year olds and partitioned into meat and fat — flanks comprise no bones. The flanks were dissected according to the scheme in Figure 1.

Means of dressing-%, by-products and cutting results of flanks by experiment

Table 19 presents the means and standard deviations of dressing-%, by-products and cutting results of flanks.

Stepwise regression analysis, results and discussion

Estimation of carcass composition with live weight, dressing-%, by-products and cutting results of flanks.

The R^2 for meat-kg increased only by 0.4 %-units in both age groups from the value obtained with mere carcass weight (Tables 20 and 21).

The R^2 for fat-kg for 1-year olds ($n = 142$) was 62 % (Table 20) and for $\frac{1}{2}$ -year olds 48 % (Table 21), when the traits in a) acted as estimators. When in the estimation of fat-kg the traits mentioned in b) were estimators an R^2 of 60 % was obtained for the 1-year olds (Table 20). The standardized regression coefficient for kidney fat (kg) was when calculated in both ways about twice as high as that of the other estimators.

When fat-kg was estimated for the $\frac{1}{2}$ -year olds with carcass weight, kidney and bowel fats plus cutting results of flanks, an R^2 of 70 % was obtained (Table 21, b). With mere fat-kg in flanks a 10 % higher value for R^2 was obtained than the one obtained with all the traits in a). Bowel fats would have been the fourth step in the stepwise regression analysis but the value of the regression coefficient was nonsignificant. The correlation of bowel fat (kg) on carcass fat-kg was 0.630 for the 1-year olds and only 0.181 for the $\frac{1}{2}$ -year olds.

Inclusion of bowel fats increases the R^2 of fat-kg for 1-year olds approximately by 3 %-units and, as the regression coefficient for the $\frac{1}{2}$ -year olds was nonsignificant, it seems useless to weigh carcass bowel fats for this purpose. Removal and weighing of bowel fats requires quite much extra work as the intestines are removed immediately after slaughter and taken to another room to be cleaned and washed.

Instead, the R^2 for fat-kg obtained with flank fats is considerably higher than those obtained with any other trait, and removal of one flank and partition-

Table 19. Means and standard deviations of dressing-%, percentages of carcass by-products and cutting results of flanks by experiment.

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
1-year olds, n = 205										
Dressing-%										
I	51.0	2.2	*		—	6.0	0.3	**		—
II	47.0	1.2	—	—	—	6.6	0.5	—	—	—
III	46.8	1.6	—	—	—	6.3	0.5	*	—	—
IV	45.4	1.3		***	—	6.2	0.5		***	—
V	46.2	1.7	—	**	—	7.3	0.6	—	—	—
VI	44.9	1.5		—	—	7.0	0.3		—	—
Head, %										
I	6.0	0.3				6.0	0.3	**		—
II	6.6	0.5				6.6	0.5	—	—	—
III	6.3	0.5				6.3	0.5	*	—	—
IV	6.2	0.5		***	—	6.2	0.5		***	—
V	7.3	0.6		**	—	7.3	0.6	—	—	—
VI	7.0	0.3		—	—	7.0	0.3		—	—
Liver, %										
I	2.5	0.2	—		—	0.8	0.1	—		—
II	3.3	0.2	**	—	—	1.1	0.2	—	—	—
III	3.1	0.3	*	***	—	1.0	0.1	*	—	—
IV	2.9	0.2		**	—	0.9	0.1		—	—
V	3.1	0.2	—	**	—	0.8	0.1	—	—	—
VI	3.1	0.2		—	—	0.8	0.1		—	—
Heart, %										
I	0.8	0.1				0.8	0.1	—		—
II	1.1	0.2				1.1	0.2	—	—	—
III	1.0	0.1	*	***	—	1.0	0.1	*	—	—
IV	0.9	0.1		**	—	0.9	0.1		—	—
V	0.8	0.1	—	**	—	0.8	0.1	—	—	—
VI	0.8	0.1		—	—	0.8	0.1		—	—
Kidney fat, %										
I	2.4	0.8	—		—	2.2	0.6	—		—
II	2.2	0.8	—	—	—	1.9	0.6	—	—	—
III	2.1	0.7	—	—	—	2.2	0.6	—	—	—
IV	2.3	0.6		***	—	3.4	0.9		***	—
V	1.8	0.3	*	*	—					
VI	0.3	0.1		—	—					
Bowel fat, % (n = 142)										
I	2.2	0.6	—		—	2.2	0.6	—		—
II	1.9	0.6	—	—	—	1.9	0.6	—	—	—
III	2.2	0.6	—	—	—	2.2	0.6	—	—	—
IV	3.4	0.9		***	—	3.4	0.9		***	—
V	1.8	0.3	*	*	—					
VI	0.3	0.1		—	—					
1/2-year olds, n = 94										
Dressing-%										
IX	49.6	1.4	*		—	6.9	0.5	—		—
X	50.2	1.7	***	—	—	6.6	0.5	***	*	—
Head, %										
IX	6.9	0.5				6.9	0.5	—		—
X	6.6	0.5	***	—	—	6.6	0.5	***	*	—
Liver, %										
IX	3.5	0.4	**		—	0.9	0.1	*		—
X	3.4	0.3	***	—	—	0.9	0.2	—	—	—
Heart, %										
IX	0.9	0.1				0.9	0.1	*		—
X	0.9	0.2				0.9	0.2	—	—	—
Kidney fat, %										
IX	2.0	0.4	***		—	2.0	0.4	***		—
X	2.4	0.5	***	—	—	2.4	0.7	*	—	—
Bowel fat, %										
IX	2.0	0.4	***		—	2.0	0.4	***		—
X	2.4	0.5	***	—	—	2.4	0.7	*	—	—
Flanks, kg										
IX	4.9	0.8	*		—	5.2	0.5	**		—
X	6.4	0.7	***	***	—	5.7	0.5	*	—	—
Flanks, %										
IX	5.2	0.5				5.2	0.5	**		—
X	5.7	0.5		***	—	5.7	0.5	*	—	—
Meat-% in flanks										
IX	82.2	2.3	—		—	18.1	2.2	—		—
X	84.2	4.1	—	—	—	15.6	4.2	—	—	—
Fat-% in flanks										
IX	18.1	2.2				18.1	2.2	—		—
X	15.6	4.2				15.6	4.2	—	—	—

Table 20. Result of stepwise regression analysis.

1-year olds a) Live and carcass weight, dressing-% and carcass by-products — carcass composition.
 b) Carcass weight, bowel and kidney fat — carcass fat-kg and -%.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 109$ kg						
a) 1.	Carcass weight980	.980	96.0	0.762***	0.963
2.	Kidney fat, %	-.084	.981	96.2	-0.569*	-0.031
3.	Dressing-%476	.982	96.4	0.247**	0.034
y = Fat-kg, $\bar{y} = 6.6$ kg						
a) 1.	Kidney fat, kg700	.700	49.0	2.152***	1.171
2.	Kidney fat, %535	.752	56.6	-1.724***	-0.556
3.	Liver, %	-.327	.763	58.2	-1.631**	-0.185
4.	Heart, %	-.102	.777	60.4	5.970*	0.363
5.	Head, %	-.430	.786	61.8	-0.808*	-0.172
b) 1.	Kidney fat, kg700	.700	49.0	1.931***	1.051
2.	Kidney fat, %535	.752	56.6	-1.661***	-0.536
3.	Bowel fat, kg630	.772	59.6	0.373**	0.232
y = Bone-kg, $\bar{y} = 30.9$ kg						
a) 1.	Carcass weight731	.731	53.4	0.097***	0.485
2.	Kidney fat, kg011	.759	57.6	-0.655***	-0.234
3.	Live weight713	.767	58.8	0.035*	0.332
y = Meat-%, $\bar{y} = 72.9$ %						
a) 1.	Carcass weight367	.367	13.5	0.078***	0.852
2.	Kidney fat, %	-.248	.434	18.8	-0.463***	-0.217
3.	Live weight279	.463	21.4	-0.022*	-0.459
4.	Heart, kg047	.482	23.2	-1.034*	-0.147
y = Fat-%, $\bar{y} = 4.0$ %						
a) 1.	Kidney fat, kg645	.645	41.6	0.610***	0.648
2.	Liver, kg	-.114	.670	44.9	-0.688***	-0.277
3.	Heart, %009	.688	47.3	1.652**	0.209
4.	Head, %	-.253	.710	50.4	-0.503**	-0.209
b) 1.	Kidney fat, kg645	.645	41.6	0.468***	0.497
2.	Bowel fat, %543	.689	47.5	0.421***	0.283
y = Bone-%, $\bar{y} = 20.7$ %						
a) 1.	Carcass weight	-.446	.446	19.9	-0.077***	-0.865
2.	Kidney fat, %	-.279	.539	29.1	-0.648***	-0.313
3.	Live weight	-.393	.555	30.8	0.017*	0.369
4.	Heart, kg	-.055	.567	32.1	0.885*	0.129

In estimation of meat-kg and -%, and bone-kg and -% the no. of animals is 205, and in that of fat-kg and -% 142.

Table 21. Result of stepwise regression analysis.
 $\frac{1}{2}$ -year olds a) Live and carcass weight, dressing-% and carcass by-products — carcass composition.
 n = 94 b) Carcass weight, bowel and kidney fat, and cutting results of flanks — carcass fat-kg and -%.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 76.6$ kg						
a) 1.	Carcass weight982	.982	96.4	0.928***	1.146
2.	Live weight882	.984	96.8	-0.079***	-0.176
y = Fat-kg, $\bar{y} = 3.4$ kg						
a) 1.	Live weight553	.553	30.6	0.143*	0.241
2.	Kidney fat, kg548	.654	42.8	0.709***	0.375
3.	Liver, kg508	.691	47.7	0.609**	0.271
b) 1.	Fat-kg in flanks759	.759	57.6	3.605***	1.056
2.	Kidney fat, kg548	.808	65.3	0.488***	0.258
3.	Fat-% in flanks529	.838	70.2	-0.120***	-0.455
y = Bone-kg, $\bar{y} = 21.2$ kg						
a) 1.	Live weight796	.796	63.4	0.069***	0.658
2.	Head, kg620	.819	67.1	0.753**	0.237
y = Meat-%, $\bar{y} = 73.9$ %						
a) 1.	Dressing-%421	.421	17.7	0.242**	0.306
2.	Head, %	-.359	.482	23.2	-0.725**	-0.276
3.	Kidney fat, %	-.210	.519	26.9	-0.551*	-0.195
y = Fat-%, $\bar{y} = 3.3$ %						
a) 1.	Kidney fat, kg518	.518	26.8	0.783***	0.520
2.	Liver, %282	.592	35.0	0.613***	0.286
b) 1.	Fat-kg in flanks689	.689	47.5	1.431***	0.526
2.	Kidney fat, %452	.748	56.0	0.465***	0.283
3.	Flanks, %508	.765	58.5	0.297**	0.187
y = Bone-%, $\bar{y} = 20.6$ %						
a) 1.	Carcass weight	-.435	.435	18.9	-0.154***	-1.187
2.	Live weight	-.298	.514	26.4	0.046**	0.650
3.	Head, kg	-.067	.553	30.6	0.552*	0.255

ing it into meat and fat is a simple and fast procedure. If only one flank is removed per carcass this should not decrease its commercial value too much.

With 1-year olds the R² for fat-% obtained with traits in a) was 50 % (Table 20) and with traits in b) 47.5 % (Table 20), the best estimating trait in both a) and b) being kidney fat-kg. Correspondingly with $\frac{1}{2}$ -year olds the R² obtained with traits in a) on fat-% was 35 % and with those in b) 58.5 % (Table 21).

The R^2 -values obtained, 50 % and 58.5 %, can be considered quite satisfactory as estimation of the percentages has been fairly unreliable in general.

Estimation of bone - kg was less certain with the 1-year olds than with the $\frac{1}{2}$ -year olds, R^2 -values were 59 % and 67 % (Tables 20 and 21). The correlation of head (kg) on bone-kg was 0.553 for 1-year olds and 0.630 for $\frac{1}{2}$ -year olds.

Estimation of meat and bone percentages was unreliable in both age groups. Carcass weight and live weight were the best estimators of bone-%, their correlations on bone-% were negative (Tables 20 and 21).

In quite a few studies carcass characteristics have been estimated with the same traits as here. In several studies, however, in the estimation of a trait thickness of fat cover, the area of the MLD and other carcass measurements have been used in addition (EPLEY et al. 1970, HENDERSON et al. 1966 a, b, LEVY et al. 1968, du BOSE et al. 1967, DUMONT et al. 1961, ÇUNDIFF et al. 1967 and BRÄNNÄNG and NILSSON 1969).

Through combination of several traits high multiple correlations and R^2 -values have been obtained for percentages, too. Especially estimation of fat has been more reliable in several other studies than in the present one, although the R^2 -values for fat-kg and fat-% in the present study can be regarded fairly satisfactory.

Table 22 presents the interrelations literature provides between live weight, carcass weight, dressing-%, and by-products on carcass traits. Also some correlations obtained in the present study are included.

In several respects the results differ largely from each other. For example the correlations between carcass weight and carcass fat obtained by ALLEN et al. (1968), FIELD et al. (1966) and BUTTERFIELD (1965) were much higher than the ones obtained in the present study.

Vth estimator: cutting results of shanks

Dissection of shanks

Dissection of the shanks, like that of the flanks described in Chapter IV, require a partial destruction of a carcass. Removal of the shanks from the carcass and cutting them into selections is quite a simple and fast procedure. It is, however, difficult to estimate how much the removal of the shanks reduces the commercial value of a carcass.

Only the fore shanks were removed from the carcasses in experiments IX and X. They were cut into meat, bone and tendon selections, no fat was found. From the other carcasses of the 1-year olds, except in experiment I, the fore and hind shank on the right hand side of the carcasses were removed and dissected boneless, tendons were not removed from the muscular tissue.

In experiments V and VI there were differences in dissection and cutting techniques of the shanks if compared to those in II, III and IV. In all of them, however, tibia fibula has been separated in one piece. In experiments V and VI part of the meat in the fore shank remained with the

Table 22. Correlations between carcass composition and live weight, carcass weight, dressing-% and by-products found in the literature.

	Meat kg	Fat kg	Bone kg	Meat %	Fat %	Bone %
Live weight						
du BOSE et al. 196785 ¹⁾	—	—	.02 ²⁾	—	—
BRACKELSBURG & WILLHAM 1968 .	—	—	—	-.28	.18	-.33
SCHMITTEN 1968	—	—	—	.27	—	—
VARO 196990	.36	—	—	—	—
Present study	1-yr. .911	.496	.713	.279	.120	-.393
	1/2-yr. .882	.553	.796	.186	.305	-.298
Carcass weight						
ALLEN et al. 196886	.80	.77	-.03	.04	.0
du BOSE et al. 196794 ¹⁾	—	—	-.03 ²⁾	—	—
BRACKELSBURG et al. 196841	.08	—	.10	-.12	—
BRACKELSBURG & WILLHAM 1968 .	—	—	—	-.30	.35	-.40
BUTTERFIELD 196597	.84	.80	—	—	—
DINKEL et al. 1965	—	—	—	-.49	—	—
FIELD et al. 1966	R ² = 97 %	82 %	92 %	31 %	48 %	54 %
Present study	1-yr. .980	.484	.731	.367	.088	-.446
	1/2-yr. .982	.504	.773	.324	.264	-.435
Dressing-%						
NIELSEN 1962	—	—	—	.25	-.07	-.26
WISMER-PEDERSEN 1969	bulls	—	—	.32	-.06	—
	calves	—	—	.42	-.09	—
				.39	-.15	
Present study	1-yr. .476	.103	.289	.340	-.073	-.254
	1/2-yr. .491	.101	.167	.421	-.043	-.431
Kidney and bowel fat, kg						
BUTTERFIELD 1965	—	.91	—	—	—	—
VARO 1969	—	.46	—	—	—	—
Kidney fat kg						
du BOSE et al. 196730 ¹⁾	—	—	-.53 ²⁾	—	—
HINKS & BECH ANDERSEN 1968	—	—	—	-.40	—	—
Present study	1-yr. —	.700	—	—	.645	—
	1/2-yr. —	.548	—	—	.518	—

1) Boneless roast and steak, kg.

2) » » » » » %

shoulders, and part of the hind shanks with the rounds. In comparison of the mean results (Table 23) the difference in dissection can be observed.

Means of cutting results of shanks by experiment

Table 23 presents the means and standard deviations of the cutting results of shanks by experiment.

Differences in dissection techniques are most apparent in bone-% and meat-% where differences between experiments II, III and IV on one hand and V and VI on the other hand are distinct (Table 23).

Table 23. Means and standard deviations of cutting results of fore and hind shanks by experiment.

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
1-year olds, n = 181										
Hind shank, kg						Hind shank, %				
II	3.8	0.3	***	—	—	2.6	0.2	—	—	—
III	4.4	0.5	**	*	*	2.7	0.3	—	—	—
IV	4.0	0.3 sires	***	**	—	2.5	0.2	—	***	—
V	3.5	0.4	*	—	—	2.6	0.2	—	—	—
VI	3.6	0.3	—	—	—	2.9	0.2	—	—	—
Meat-% in hind shank						Bone-% in hind shank				
II	55.2	1.8	—	—	—	44.8	1.8	—	—	—
III	59.1	2.4	—	—	—	40.9	2.3	—	—	—
IV	57.9	2.8	—	**	—	42.4	2.8	—	**	—
V	51.6	1.7	—	—	*	48.4	1.7	—	—	*
VI	53.5	1.9	—	*	—	46.7	1.9	—	*	—
Fore shank, kg						Fore shank %				
II	3.2	0.5	**	—	**	2.3	0.3	—	—	**
II	3.9	0.4	**	*	**	2.4	0.1	—	—	—
IV	3.7	0.4 sires	*	**	—	2.6	0.2	—	**	—
V	2.5	0.2	**	—	**	1.9	0.1	—	—	—
VI	2.3	0.2	—	—	—	1.9	0.1 sires	**	—	—
Meat-% in fore shank						Bone-% in fore shank				
II	56.4	2.7	—	—	—	43.6	2.7	—	—	—
III	55.3	2.1	—	—	—	44.8	2.0	—	—	—
IV	57.4	2.6	—	**	—	43.0	2.3	—	**	—
V	52.0	1.1	—	***	—	48.0	1.1	—	***	—
VI	52.1	2.5	—	—	—	47.7	2.2	—	—	—
1/2-year olds, n = 94										
Fore shanks, kg						Fore shanks, %				
IX	4.9	0.9	**	—	—	5.1	0.2	**	—	**
X	5.4	0.4	***	***	—	4.9	0.3	*	—	—
Meat-% in fore shanks						Bone-% in fore shanks				
IX	47.2	2.8	—	—	—	45.8	1.9	*	—	—
X	51.9	3.6	—	—	—	44.4	2.6	**	—	—

With 1/2-year olds bone-% and meat-% did not total 100 % because tendons are dissected into another selection. The share of tendons is 4–6 %, which is not, however, included in the calculations.

Stepwise regression analysis, results and discussion

Estimation of carcass composition with cutting results of shanks.

In addition to the cutting results of the shanks, carcass weight has acted as one estimator (Tables 24 and 25). Where carcass weight has produced no additional information the results have been omitted from the tables.

Table 24. Result of stepwise regression analysis.
 1-year olds a) Cutting results of fore and hind shank — carcass meat-kg, bone-kg and bone-%.
 n = 181 b) Carcass weight and cutting results of fore and hind shank — carcass meat-kg,
 bone-kg and bone-%.
 c) Carcass weight and cutting results of fore shank — carcass bone-kg and
 bone-%.
 d) Carcass weight and cutting results of hind shank — carcass bone-kg and
 bone-%.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 106$ kg						
a) 1.	Fore shank, kg690	.690	47.6	7.657***	0.257
2.	Fore shank, %	-.224	.916	83.9	-11.582***	-0.202
3.	Hind shank, %	-.422	.929	86.3	-34.950***	-0.623
4.	Hind shank, kg642	.975	95.1	21.312***	0.724
b) 1.	Carcass weight984	.984	96.8	0.827***	1.052
2.	Hind shank, kg642	.985	97.0	-2.956***	-0.100
3.	Bone-% in hind shank	-.040	.986	97.2	-0.139*	-0.029
y = Bone-kg, $\bar{y} = 30.9$ kg						
a) 1.	Bone-kg in hind shank768	.768	59.0	7.461***	0.420
2.	Fore shank, kg664	.798	63.7	6.547***	0.853
3.	Fore shank, %008	.818	66.9	-3.760***	-0.255
4.	Meat-kg in fore shank562	.825	68.1	-4.508*	-0.368
c) 1.	Carcass weight716	.716	51.3	0.071***	0.352
2.	Bone-kg in fore shank685	.780	60.8	9.071***	0.538
3.	Fore shank, %008	.787	61.9	-2.298*	-0.156
d) 1.	Bone-kg in hind shank768	.768	59.0	9.275***	0.523
2.	Carcass weight716	.810	65.6	0.072***	0.355
y = Bone-%, $\bar{y} = 21.0$ %						
a) 1.	Hind shank, %564	.564	31.8	5.454***	0.828
2.	Bone-% in hind shank166	.687	47.2	0.298***	0.534
3.	Bone-kg in hind shank051	.715	51.1	-3.520***	-0.435
4.	Bone-kg in fore shank061	.743	55.2	2.245***	0.292
c) 1.	Carcass weight	-.464	.464	21.5	-0.066***	-0.723
2.	Bone-kg in fore shank061	.641	41.1	3.578***	0.465
3.	Bone-% in fore shank412	.654	42.8	0.092***	0.153
d) 1.	Hind shank, %564	.564	31.8	4.058***	0.616
2.	Bone-% in hind shank166	.687	47.2	0.210***	0.376
3.	Carcass weight	-.464	.716	51.3	-0.020***	-0.218

With mere weight and percentage of the shanks the meat-kg could be estimated with R² of 95 % to 96 %. When carcass weight was included as one estimator of meat-kg this alone gave a higher R²-value than the cutting results of the shanks. The shanks increased the R² with 0.4–1.8 %-units (Tables 24 and 25).

Table 25. Result of stepwise regression analysis.
 $\frac{1}{2}$ -year olds a) Cutting results of fore shanks — carcass composition.
 n = 94 b) Carcass weight and cutting results of fore shanks — carcass composition.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 76.6$ kg						
a) 1.	Meat-kg in fore shanks656	.656	43.0	30.182***	1.242
2.	Fore shanks, %	-.340	.867	75.2	-17.459***	-0.692
3.	Meat-% in fore shanks154	.979	95.8	-1.259***	-0.598
b) 1.	Carcass weight982	.982	96.4	0.772***	0.953
2.	Meat-% in fore shanks154	.987	97.4	0.146***	0.069
3.	Fore shanks, %	-.340	.990	98.0	-2.075***	-0.082
4.	Bone-% in fore shanks	-.158	.991	98.2	-0.142*	-0.048
y = Fat-kg, $\bar{y} = 3.4$ kg						
b) 1.	Carcass weight544	.544	29.6	0.058***	0.544
y = Bone-kg, $\bar{y} = 21.2$ kg						
a) 1.	Bone-kg in fore shanks742	.742	55.1	20.945***	3.511
2.	Fore shanks, kg544	.780	60.8	-6.257***	-2.538
3.	Bone-% in fore shanks337	.864	74.6	-0.794***	-1.136
4.	Fore shanks, %062	.907	82.1	-1.783***	-0.299
y = Meat-%, $\bar{y} = 73.9$ %						
a) 1.	Meat-% in fore shanks543	.543	29.5	0.099*	0.227
2.	Fore shanks, %	-.437	.713	50.8	-2.372***	-0.518
3.	Meat-kg in fore shanks339	.735	53.7	1.003*	0.227
4.	Bone-% in fore shanks	-.518	.749	56.1	-0.121*	-0.225
y = Fat-%, $\bar{y} = 3.3$ %						
b) 1.	Carcass weight264	.264	7.0	0.022*	0.264
y = Bone-%, $\bar{y} = 20.6$ %						
a) 1.	Bone-% in fore shanks603	.603	36.4	0.213***	0.449
2.	Fore shanks, %507	.788	62.1	2.468***	0.608
3.	Meat-kg in fore shanks	-.357	.836	69.9	-1.298***	-0.332

In the estimation of fat-kg and fat-% the results obtained with shank cutting results were unreliable in both age groups.

For the 1-year olds an R² of 68 % was obtained for bone-kg with the cutting results of shanks (Table 24, a), carcass weight gave no additional information to this value. The value of R² did not increase in the experiments (II—IV) where the shanks had been dissected uniformly.

When the estimation was performed separately with the fore shanks and again with the hind shanks and carcass weight, the result remained less reliable than the one obtained with mere cutting results of the fore and hind shanks (Table 24, c, d).

With cutting results of the fore shanks an R² of 82 % was obtained with the $\frac{1}{2}$ -year olds (Table 25, a). With carcass weight and cutting results of fore shanks the R² was 83 %.

Estimation of meat-% was rather unreliable with the 1-year olds, but for the 1/2-year olds an R^2 of 56 % was obtained with the mere cutting of the fore shanks (Table 25, a).

The R^2 for bone % was 55 % for the 1-year olds (Table 24, a). In the experiments where the shanks were dissected in the same way (experiments II–IV), the R^2 obtained was about 10 %-units higher than the one for the whole material. For the 1/2-year olds the R^2 for bone-% was 70 % (Table 25, a).

According to personal impression it seems probable that with the cutting results of mere fore or hind shank the same degree of reliability can be achieved as with dissection of both the fore and the hind shanks. In earlier studies at the S.W. Finland Experiment Station in experiments on Charolais cross-breeds similar values for R^2 for bone-kg and bone-% were obtained with cutting results of the fore shanks as in the present study with the 1/2-year olds. In all the experiments the fore shanks were removed and cut into selections in the same manner as in the present study in the experiments for the 1/2-year olds.

In several studies attention has been paid to the information of carcass composition obtained with dissection and cutting of the shanks. This is quite natural considering that removal of the shanks requires no high expenses and no entire destruction of the carcass.

In the study by CALLOW (1962) the following correlations were obtained with shank cutting results:

Fore shank meat-kg	—	carcass meat-kg	$r = 0.96$
Hind »	»	»	$r = 0.62$
Fore » fat-kg	—	» fat-kg	$r = 0.53$
Hind »	»	»	$r = 0.58$
Fore » bone-kg	—	» bone-kg	$r = 0.98$
(radius + ulna)			
Hind »	»	»	$r = 0.97$
(tibia fibula)			

In a study by BUTTERFIELD (1965) one half of 29 carcasses were dissected into muscles and bones and these were compared to the meat and bone quantities in the other half. Between separate bones and the bone quantity correlations of 0.92–0.99 were obtained. Radius and ulna gave the highest estimations. The correlations between separate muscles and meat quantity varied between 0.87 and 0.99. The correlation between fore shank meat and the total meat quantity was 0.95. Correlations for the percentages were low.

The material in a study by ORME et al. (1959 b) comprised 8 A. Angus and 23 Hereford steers, of which the left half was dissected. The results follow:

Fore shank bone-kg	—	carcass meat-kg	$r = 0.66$
» » » %	—	» meat-%	$r = 0$
Hind » bone-kg	—	» meat-%	$r = 0.69$
» » » %	—	» meat-%	$r = -0.01$

In a study by BRACKELSBURG et al. (1968) the correlation obtained between fore shank meat-% and carcass meat-% was 0.61, the correlation between the corresponding bone-% was 0.43, and between fat-% 0.23.

COLE et al. (1960) obtained a correlation of 0.81 between fore shank meat-kg and carcass meat-kg and through addition of carcass weight the R-value rose up to 0.89.

Materials in these studies have been quite different from the one in the present study, therefore there remains only little ground for comparison. In estimation of fat the present results with shank dissection provided virtually no information. Instead in some foreign studies relatively high correlations have been obtained even in estimation of fat, which evidently is due to the abundancy of fat even in the shanks of the breeds studied. In the present study no fat is cut off the shanks as it is very scarce in young animals of our breeds.

In several studies the cutting results of the shanks have been estimators of other carcass traits in addition to other estimating characteristics. But because the thickness of fat cover and the area of the MLD have often been included, there is no reason to do any comparisons here. (Some results are presented in Chapter IV).

Vith estimator: *cutting results of the fore and hind quarters*

As the fore quarters of a carcass, excluding the shoulders, are regarded as the cheaper part, and the shoulders with the hind quarters — the rounds and the back — as the more valuable parts of a carcass, a principle was established that if the carcass traits can not be properly estimated with any other method, the cheaper part i.e. the fore quarters could be dissected.

In all experiments the rounds were dissected of the hind quarters behind the last lumbar vertebra. The bones were removed of the steaks in the rounds, different muscles were not, however, separated but the flesh was weighed in one piece. The MLD was removed of the back. The other muscles of the back were not weighed separately. The dissection of the inner fillets differed between experiments, therefore they are excluded from the calculations.

In all the carcasses the fore and hind quarters were separately partitioned into meat, fat, bone and tendons.

Means of cutting results of the fore and hind quarters by experiment

Tables 26 and 27 present the means and standard deviations of the cutting results of the fore and hind quarters by experiment.

The proportions of the fore and hind quarters to the total carcass weight remained rather constant from one experiment to another. The differences in percentages between breeds were, however, statistically significant in all experiments. Differences in percentages of meat, fat and bone were only in a few cases statistically significant for the 1-year olds. For the ½-year olds the differences of most traits were statistically significant between breeds.

Table 26. Means and standard deviations of cutting results of the fore quarters by experiment.

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
1-year olds, n = 205										
Fore quarters, kg						Fore quarters, %				
I	73.0	5.8	*		—	43.0	0.9	***		—
II	60.9	4.3	***		—	43.0	1.0	***	—	***
III	68.4	5.3	*	***	**	42.7	0.7	***	*	—
IV	68.2	8.2		***	—	42.6	0.8		—	—
V	58.7	6.4	—	—	**	43.6	1.0	**	—	—
VI	55.2	4.2		—	—	44.7	1.4		—	—
Meat-kg						Meat-%				
I	54.3	4.5	*		—	74.4	1.3	—		—
II	43.4	3.4	**	—	—	71.3	1.5	—	—	—
III	48.7	4.2	*	***	**	71.2	1.7	—	—	—
IV	47.2	6.3		***	—	69.1	1.9		*	—
V	38.5	4.6	—	—	*	65.4	1.3	—	—	—
VI	40.2	3.4		—	—	72.7	1.3		—	—
Fat-%						Bone-%				
I	3.3	0.8	—		—	20.1	0.8	*		—
II	3.0	0.9	—	—	—	23.4	1.0	—	—	—
III	3.4	1.3	—	—	—	22.9	1.4	—	*	—
IV	5.3	1.1		***	—	22.7	1.8		***	—
V	10.0	1.2	*	—	—	22.7	1.2	—	*	—
VI	0.4	0.2		—	*	25.3	0.4		—	—
$\frac{1}{2}$ -year olds, n = 94										
Fore quarters, kg						Fore quarters, %				
IX	39.1	3.9	—		*	41.5	0.8	**		—
X	45.5	3.2	***	***	**	41.3	0.6	***	—	—
Meat-kg						Meat-%				
IX	27.3	3.0	—		*	69.9	1.4	*		—
X	33.1	2.6	***	***	**	72.5	1.9	**	—	—
Fat-%						Bone-%				
IX	3.1	0.6	**		—	23.7	1.3	***		—
X	2.5	1.0	—	—	—	22.8	1.6	***	—	—

Stepwise regression analysis, results and discussion

Estimation of carcass composition with carcass weight and cutting results of the fore and hind quarters.

Carcass weight estimated the meat-kg with an R^2 of 96% as already mentioned. With the addition of the cutting results of the fore and hind

Table 27. Means and standard deviations of cutting results of hind quarters by experiment

Exp. No.	Mean	SD	Statistical significance			Mean	SD	Statistical significance		
			Breed	Feed.	Age			Breed	Feed.	Age
1-year olds, n = 205										
Hind quarters, kg						Hind quarters, %				
I	97.0	7.3	***		—	57.0	0.9	***		—
II	81.1	6.6	***		—	57.0	1.0	***		***
III	92.3	7.2	**	***	**	57.3	0.7	***	*	—
IV	91.6	10.5		***	—	57.4	0.8			—
V	76.2	8.2	*		*	56.4	1.0	**		—
VI	68.4	4.8			—	55.3	1.4			—
Meat-kg						Meat-%				
I	75.4	5.9	***		—	77.7	1.2			—
II	61.4	5.4	***		—	75.5	1.0			—
III	69.5	6.1	**	***	**	75.5	1.6			—
IV	67.0	8.1		***	—	73.2	1.3		***	—
V	53.3	6.0	**		*	69.9	1.1	**	*	—
VI	53.1	4.0			—	77.6	1.0			—
Fat-%						Bone-%				
I	3.9	0.9			—	16.9	0.5	***		—
II	2.3	0.7			—	19.5	0.8			—
III	3.3	1.3			—	18.4	1.5			—
IV	6.0	1.4		***	—	18.6	1.5		***	—
V	10.1	1.5	*	**	—	19.2	0.8		**	—
VI	0.4	0.2			*	21.6	1.1			—
1/2-year olds, n = 94										
Hind quarters, kg						Hind quarters, %				
IX	55.2	5.9	*		*	58.7	0.8	**		—
X	64.8	4.9	***	***	*	58.7	0.6	***		—
Meat-kg						Meat-%				
IX	41.3	4.7	**		*	74.7	1.4	**		—
X	49.5	3.9	***	***	*	76.4	1.1	***	**	—
Fat-%						Bone-%				
IX	4.0	0.7	*		—	19.6	1.2	**		—
X	3.5	0.9			—	18.3	1.0	***		*

quarters the coefficient increased by 2—3 %-units in both age groups (Tables 28 and 30). It is, however, questionable whether for such a modest, less than 3 %-units, increase in the coefficient it is reasonable to perform a costly dissection.

In the estimation of fat-kg considerably higher values of R^2 , 86 %-87 %, were obtained with cutting results of the fore and hind quarters with the 1-year olds than with any earlier estimator (Table 28, a, b).

Table 28. Result of stepwise regression analysis.
 1-year olds a) Carcass weight and cutting results of fore quarters — quantities of carcass
 n = 205 meat, fat and bone.
 b) Carcass weight and cutting results of hind quarters — quantities of carcass
 meat, fat and bone.

Step	x	r	R	R ²	b	β
y = Meat-kg, $\bar{y} = 109$ kg						
a) 1.	Carcass weight980	.980	96.0	0.793***	1.002
2.	Meat-% in fore quarters433	.989	97.8	0.939***	0.138
3.	Fat-kg » » »397	.990	98.0	-2.084***	-0.158
4.	Fat-% » » »009	.991	98.2	1.239***	0.119
b) 1.	Carcass weight980	.980	96.0	0.737***	0.932
2.	Meat-% in hind quarters440	.992	98.4	1.227***	0.147
3.	Boneless steaks of round, %246	.992	98.4	0.120*	0.021
y = Fat-kg, $\bar{y} = 6.7$ kg						
a) 1.	Fat-kg in fore quarters918	.918	84.3	2.088***	0.871
2.	Bone-% » » »	-.474	.922	85.1	-0.150***	-0.108
3.	Fore quarters, %	-.044	.925	85.6	-0.135*	-0.060
b) 1.	Fat-% in hind quarters830	.830	68.9	1.197***	0.714
2.	Carcass weight484	.928	86.1	0.048*	0.336
3.	Fat-kg in hind quarters.....	.579	.930	86.5	0.059*	0.062
4.	Meat-kg » » »396	.931	86.7	-0.142**	-0.459
5.	Hind quarters, kg496	.933	87.0	0.128**	0.525
y = Bone-kg, $\bar{y} = 30.9$ kg						
a) 1.	Bone-kg in fore quarters902	.902	81.4	1.983***	0.924
2.	Fore quarters, %039	.910	82.8	-0.341***	-0.122
b) 1.	Bone-kg in hind quarters904	.904	81.7	1.473***	0.765
2.	Hind quarters, %	-.030	.913	83.4	-0.360***	-0.128
3.	Rounds with hind shanks, kg761	.919	84.5	0.106***	0.188
4.	Boneless steaks of round, %	-.131	.920	84.6	-0.089*	-0.062

With the 1/2-year olds the R² was 73 % with the cutting results of the fore quarters and 86 % with those of the hind quarters. (Table 30, a, b). With kidney fats and cutting results of the flanks the R² obtained was 70% (Chapter IV). For this reason the purpose of cutting the fore quarters becomes questionable considering cutting of large materials.

Almost uniform R²-values were obtained for bone-kg with cutting results of both the fore and hind quarters, R²-values were from 83 % to 85 % and 95 % (Tables 28 and 30).

The R² for meat-% was 76 % with cutting results of fore quarters and 82 % with cutting results of hind quarters for both age groups. (Tables 29 and 30).

The R² obtained for fat-% was 12—14 %-units higher with the cutting results of hind quarters than with those of the fore quarters (Tables 29 and 30).

In estimation of bone-% a higher result was obtained here than with any other earlier presented estimator: the R^2 for the 1-year olds was 88 % (Table 29, a, b) with cutting results of both the fore and hind quarters and for the 1/2-year olds 81 % and 91 % respectively (Table 30, a, b).

In the literature no studies were found in which the carcass traits were estimated with the cutting results of the fore or hind quarters alone. However, results obtained in other countries are seldom comparable to the ones presented here because even the dissection of a carcass into fore and hind quarters may be done between the 6th and the 7th rib till the 13th rib and the 1st lumbar vertebra. The weight of the fore and hind quarters separately and their proportion of carcass weight thus varies according to the separation of the quarters.

Table 29. Result of stepwise regression analysis.
 1-year olds a) Carcass weight and cutting results of fore quarters — percentages of carcass
 n = 205 meat, fat and bone.
 b) Carcass weight and cutting results of hind quarters — percentages of carcass
 meat, fat and bone.

Step	x	r	R	R^2	b	β
$y = \text{Meat-\%}, \bar{y} = 72.9 \%$						
a) 1.	Meat-% in fore quarters840	.840	70.6	0.440***	0.558
2.	Carcass weight367	.847	71.7	0.057***	0.625
3.	Fat-kg in fore quarters	-.183	.855	73.3	-0.461***	-0.300
4.	Bone-kg » » »	-.011	.871	75.9	-0.371***	-0.377
b) 1.	Meat-% in hind quarters892	.892	79.6	0.791***	0.812
2.	Hind quarters, %	-.115	.899	80.8	-0.160***	-0.124
3.	Bone-% in hind quarters	-.516	.905	81.9	-0.094*	-0.090
4.	Boneless steaks, kg461	.908	82.4	0.035*	0.084
$y = \text{Fat-\%}, \bar{y} = 4.4 \%$						
a) 1.	Fat-% in fore quarters870	.870	75.7	0.840***	0.838
2.	Bone-% » » »	-.349	.876	76.7	-0.082**	-0.112
b) 1.	Fat-% in hind quarters944	.944	89.1	0.838***	0.944
$y = \text{Bone-\%}, \bar{y} = 20.7 \%$						
a) 1.	Bone-% in fore quarters918	.918	84.3	0.757***	0.879
2.	Fore quarters, %	-.017	.931	86.7	0.194***	0.156
3.	Fat-% in fore quarters	-.355	.934	87.2	-0.110***	-0.093
4.	Meat-kg » » »	-.512	.936	87.6	-0.001*	-0.076
b) 1.	Bone-% in hind quarters913	.913	83.4	0.513***	0.506
2.	Boneless steaks of round, %	-.494	.921	84.8	-0.070***	-0.110
3.	Rounds with hind shanks, %352	.926	85.7	0.093*	0.070
4.	Carcass weight	-.446	.930	86.5	-0.036***	-0.401
5.	Bone-kg in hind quarters202	.935	87.4	0.267***	0.311
6.	Fat-% » » »	-.388	.937	87.8	-0.149*	-0.143
7.	Meat-% » » »	-.463	.938	88.0	-0.088*	-0.093

Table 30. Result of stepwise regression analysis.
¹/₂-year olds a) Carcass weight and cutting results of fore quarters — carcass composition.
n = 94 b) Carcass weight and cutting results of hind quarters — carcass composition.

Step	x	r	R	R ²	b	β
y = Meat-kg, \bar{y} = 76.6 kg						
a) 1.	Carcass weight982	.982	96.4	0.813***	1.004
2.	Meat-% in fore quarters468	.994	98.8	0.488***	0.117
3.	Bone-kg » » »617	.994	98.8	-0.722***	-0.084
b) 1.	Meat-kg in hind quarters987	.987	97.4	1.155***	0.708
2.	Hind quarters, %159	.995	99.0	-0.786***	-0.081
3.	Carcass weight982	.996	99.2	0.231***	0.285
4.	Bone-% in hind quarters	-.543	.996	99.2	-0.312***	-0.047
y = Fat-kg, \bar{y} = 3.4 kg						
a) 1.	Fat-kg in fore quarters823	.823	67.7	1.490***	0.721
2.	Carcass weight544	.852	72.6	0.026***	0.243
b) 1.	Fat-kg in hind quarters925	.925	85.6	1.426***	0.925
y = Bone-kg, \bar{y} = 21.2 kg						
a) 1.	Bone-kg in fore quarters965	.965	93.1	1.872***	0.921
2.	Fore quarters, %	-.024	.974	94.9	-0.297***	-0.131
3.	Meat-kg in fore quarters653	.977	95.5	0.056**	0.096
b) 1.	Bone-kg in hind quarters970	.970	94.1	1.825***	0.988
2.	Hind quarters, %024	.977	95.5	-0.275***	-0.121
y = Meat-%, \bar{y} = 73.9 %						
a) 1.	Meat-% in fore quarters865	.865	74.8	0.693***	0.920
2.	Fat-% » » »	-.247	.973	76.2	0.194*	0.134
b) 1.	Meat-% in hind quarters883	.883	78.0	0.706***	0.709
2.	Bone-% » » »	-.718	.904	81.7	-0.242**	-0.201
3.	Boneless steaks, kg513	.908	82.4	0.060(*)	0.105
y = Fat-%, \bar{y} = 3.3 %						
a) 1.	Fat-% in fore quarters809	.809	65.4	0.422**	0.499
2.	Fat-kg » » »795	.821	67.4	0.558*	0.339
b) 1.	Fat-% in hind quarters902	.902	81.4	0.821***	0.902
y = Bone-%, \bar{y} = 20.6 %						
a) 1.	Bone-% in fore quarters856	.856	73.3	0.365***	0.487
2.	Meat-% » » »	-.699	.874	76.4	-0.335***	-0.502
3.	Fat-% » » »	-.151	.902	81.4	-0.383***	-0.298
b) 1.	Bone-% in hind quarters956	.956	91.4	1.020***	0.956

VIIth estimator: detailed cutting results of the carcasses

In all experiments the outer fillets and rounds including and excluding bones were weighed separately. The detailed cutting method applied to the carcasses in experiments IX and X is presented on page 395. The cutting results of flanks and shank have already been explained in Chapters IV and V. The correlations between other carcass parts and carcass composition will be discussed below.

Results and discussion

Estimation of carcass composition with detailed cutting results.

With the $\frac{1}{2}$ -year olds the R^2 obtained for meat-kg with the weights of the hind quarter cuts and carcass weight was 97 %. In addition to the carcass weight the only statistically significant factor was the weight of the back. With carcass weight the cuts of the fore quarters were nonsignificant. The carcass weight was the first estimator for bone-kg, 60 %. Of the other cuts the weight of the back increased the value by 3 %-units. For fat-kg the R^2 obtained with mere weight of flanks was 45 %. The results of the estimation of meat-%, fat-% and bone-% remained low.

Only in few studies the quantities of meat, fat and bone in some cuts of a carcass are compared to the respective traits in the whole carcass. Instead, in most of the studies the percentages of meat, fat and bone in different cuts of a carcass are compared to the respective percentages in the carcass.

However, in a study by COLE et al. (1960) the correlations of the meat-kg in the rounds, shoulders and back on the meat-kg of the whole carcass have been calculated. The following correlations were obtained: 0.95, 0.93 and 0.80 respectively. In the present study the corresponding correlations were 0.969, 0.881 and 0.894 which are very close to those obtained by Cole et al. In the same study the following values for R were obtained with the stepwise regression analysis on the meat-kg in the cuts:

Carcass weight + round	meat, kg - carcass	meat-kg	R = 0.97
» + shoulder	» » - »	»	R = 0.94
» + back	» » - »	»	R = 0.90-0.85

Below are presented the correlations by THORTON and HINER (1965) and the correlations obtained in the present study between the weight of rounds and the weights of carcass meat, fat and bone. In the study in question the material comprised 43 A. Angus and 80 Hereford calves, age 6 months, thus the results obtained are well comparable to the ones obtained in this study for the $\frac{1}{2}$ -year olds.

	Meat kg		Fat-kg		Bone-kg	
	AA	H	AA	H	AA	H
Thorton & Hiner: weight of the rounds	0.93	0.96	0.24	0.84	0.80	0.95
Present study $\frac{1}{2}$ -yr.	0.963		0.476		0.786	
1-yr.	0.943		0.394		0.761	

In the study by VARO (1969) the weight of the rounds was the first estimator of carcass meat-kg, fat-kg and bone -kg in the younger part of the material.

When the rounds were omitted, the R-value obtained with the ratio of carcass weight and size for the meat-kg was 0.949, the percentage of the rounds increased the R-value till 0.962. In the estimation of bone-kg the first estimator was the weight of head, the R-value obtained was 0.691, and the second one the percentage of the rounds, R-value obtained was 0.779. The highest estimate for the percentage of meat and bone was also obtained with the rounds.

It can be observed that the correlations obtained for the meat-kg and bone-kg in the present study are uniform with those obtained in the other studies.

Table 31 presents correlations found in the literature of the percentages of meat, fat and bone of some cuts of a carcass on the respective percentages in the whole carcass. Beside them the corresponding correlations found in the present study are presented.

The correlations for the meat-% and fat-% found in the other studies (Table 31) are in several of them higher than the ones obtained here. The correlation of the bone-% in the rounds on the carcass bone-% was considerably lower in the other studies than in the present study.

Table 31. Correlations between percentages of meat, fat and bone in carcass cuts and respective percentages of the whole carcass found in the literature.

	Round	Back	Flanks	Short plate	Brisket	Fore back	Shoulder	Neck
Meat-%								
BRACKELSBURG et al. 1968	.81— .89	.89	.87	.90	.91	.96	.97 ¹⁾	—
HERTKAMPF ref. WENIGER 1965	.813	—	.710	.712	.753	.741	.872	.649
HINKS & BECH ANDERSEN 1968	—	—	—	—	.28	—	—	.52
Present study ^{1/2} -yr.	.738	.668	.186	.454	—	—	.539	.617
Fat-%								
BRACKELSBURG et al. 1968	.82— .87	.92	.89	.92	.88	.96	.96 ¹⁾	—
BUSS 1968	—	.751— .852	—	—	—	—	—	—
HERKAMPF ref. WENIGER 1965	.923	—	.717	.771	.764	.780	.832	.604
Present study ^{1/2} -yr.	.761	—	.566	—	—	—	—	—
Bone-%								
BRACKELSBURG et al. 1968	.28— .61	.64	.38 ²⁾	.66	.58	.60	.70 ¹⁾	—
HERTKAMPF ref. WENIGER 1965	.696	—	—	.445	.554	.587	.629	.428
Present study ^{1/2} -yr.	.908	.785	—	—	—	—	—	—

¹⁾ shoulder + neck, ²⁾ flanks comprise ends of ribs.

The carcass meat quantity has been estimated by some separate muscles. In the study by LILJEDAHL (1965) the weights of *Musculus pectoralis profundus* and *M.p. superficialis* were compared to the meat quantity in the carcass. The correlations obtained varied between 0.73–0.98. In the present study the correlations obtained in the experiments III and V (material in all 78 1-year olds) between the muscles in question and the carcass meat-kg was 0.774.

In the study by JENSEN (1967) the relations between separate muscles and the meat quantities of a carcass of three different weight categories were compared. The highest correlation, $r = 0.94$, was obtained with the two muscles of the rounds. Of the 20 muscles studied the muscles in the rounds and the shanks were the best estimators of meat-kg.

ORME et al. (1959 a) obtained correlations of 0.795 to 0.957 with separate muscles and the meat quantity with a material comprising 43 animals of 9½–11 years of age. The correlation between the weight of the MLD and the meat quantity was 0.843. In the present study the corresponding correlations were 0.817 for the 1-year olds and 0.835 for the ½-year olds.

There are several studies on the relation between the weight and area of the MLD and the different traits of a carcass. In earlier experiments the interrelations between area of the MLD on the other traits in a carcass have been studied (RUOHOMÄKI and VARO 1967) but, like in the present one, the correlations obtained have been low.

As in several studies the correlations between the different carcass components — meat, fat and bone — are calculated. Table 32 presents the correlations found in the literature with the corresponding correlations obtained in the present study.

In several studies it has been observed that no adequately reliable and accurate information of a carcass for scientific purpose can be obtained with dissection of only some parts or with mere partial dissection of a carcass. For example ALLEN et al. (1969) dissected only one half of the carcasses into salable boneless retail cuts but the R^2 obtained corresponded only 71 % of the result obtained by the complete cutting of the other half.

Estimation of all the traits in Chapters VI and VII requires destruction of the whole carcass. With the bigger parts, like hind quarters, the fore quarters or the rounds, higher R^2 values are obtained than with the small parts. Large parts represent quite a large portion of the quantities of meat, fat and bone which makes it self-evident that these parts are more accurate estimators of the named traits than the smaller parts. According to a personal opinion there is no great difference whether a larger or a smaller part of a carcass is separated for research if the effect of both of them is destructive for the carcass. However, if the quantities and percentages of the whole carcass could be estimated with cutting only a small part of the carcass it would save labour. When separation of even a small part, for instance separate muscles or bones, requires dissection of the carcass into two parts and destruction of one of them, this also reduces the value of the carcass.

The same applies to the partition and study of the three rib joint, which is quite abundantly used in foreign studies. Dissection of one part and destruction

Table 32. Correlations between quantities and percentages of carcass meat, fat and bone found in the literature.

		Fat-kg	Bone-kg	Meat-%	Fat-%	Bone-%
Meat-kg						
ALLEN et al. 196840	.90	—	—	—
THORTON & HINER 1965		-.06	.45	—	—	—
		-.20	.63			
COLE et al. 1960		—	.75	—	—	—
Present study	1-yr.	.399	.666	.513	.006	.502
	1/2-yr.	.478	.676	.480	.195	-.548
Fat-kg						
ALLEN et al. 1968		—	—	—	—	.28
THORTON & HINER 1965		—	.10	—	—	—
			-.31			
Present study	1-yr.	—	.166	-.197	.867	-.515
	1/2-yr.	—	.345	-.169	.942	-.324
Bone-kg						
Present study	1-yr.	—	—	-.018	-.154	.186
	1/2-yr.	—	—	-.166	.130	.219
Meat-%						
ALLEN et al. 1968		—	—	—	-.98	.74
GOTTSCH et al. 1961		—	—	—	-.91	—
HENDERSON et al. 1966 b.		—	—	—	-.95	-.77
HINKS & BECH ANDERSEN 1968		—	—	—	-.72	-.64
SCHÖN 1969		—	—	—	-.936	-.445
WISMER-PEDERSEN 1969		—	—	—	-.84	-.72
					-.92	-.83
Present study	1-yr.	—	—	—	-.406	-.583
	1/2-yr.	—	—	—	-.303	-.762
Fat-%						
ALLEN et al. 1968		—	—	—	—	-.84
SCHÖN 1969		—	—	—	—	-.686
Present study	1-yr.	—	—	—	—	-.400
	1/2-yr.	—	—	—	—	-.206

of the most expensive area of the carcass reduces its value when it is sold again. If it is in any case dissected into parts for retail or processing purposes even a larger part could be studied.

Considering material comprising large numbers of animals it seems unlikely that this could be performed in Finland under the present circumstances.

Table 33. Coefficients of determination obtained with live measurements on live and carcass weight and with live weight and live measurements on carcass weight and with initial weight and daily gain on live and carcass weight.

	Live weight		Carcass weight	
	1-yr.	1/2-yr.	1-yr.	1/2-yr.
I a. Live measurements	87	82	84	80
I b. and live weight			90	87
II. Initial weight and daily gain	96	93	86	77

Table 34. Coefficients of determination obtained with different trait groups on carcass composition.

	1-year olds						1/2-year olds					
	Meat		Fat		Bone		Meat		Fat		Bone	
	kg	%	kg	%	kg	%	kg	%	kg	%	kg	%
I a. Live measurements	78	26	37	16	64	45	78	20	37	9	78	45
I b. and live weight	85	26	39	16	65	45	84	20	36	9	81	48
II. Initial weight and daily gain	79	—	24	—	52	—	71	—	29	—	56	—
III a. Carcass quality scores	47	29	17	11	19	41	59	34	24	12	11	54
III b. and carcass weight	97	29	34	15	63	44	97	34	24	12	74	54
IV a. Live and carcass weight, dressing-% and by-products	96	23	62	50	59	32	97	27	48	35	67	31
IV b. Carcass weight, bowel and kidney fats and cutting results of flanks ¹⁾	—	—	60	47	—	—	—	—	70	59	—	—
V a. Cutting results of fore and hind shanks	95	22	29	15	68	55	—	—	—	—	—	—
V b. and carcass weight	97	24	35	15	68	56	—	—	—	—	—	—
						65 ²⁾						
V a. Cutting results of fore shanks	—	—	—	—	—	—	96	56	30	7	82	70
V c. and carcass weight	97	21	35	—	62	43	98	56	30	7	83	70
V d. Cutting results of hind shanks and carcass weight	97	24	41	—	65	51	—	—	—	—	—	—
VI. Carcass weight and cutting results of fore quarters	98	76	86	77	83	88	99	76	73	67	95	81
VI. Carcass weight and cutting results of hind quarters	98	82	87	89	85	88	99	82	86	81	95	91

¹⁾ cutting results of flanks only for 1/2-year olds.

²⁾ R²-value for the experiments where the shanks were dissected in a similar manner.

Summary

Tables 33 and 34 present the coefficients of determination for live weight, carcass weight and carcass composition obtained with different trait groups.

Procedure with which the estimates mentioned in the Tables 33 and 34 can be obtained (excluding those obtained with cutting results of fore and hind quarters):

1. Determination of live weight with weighing.
2. Measuring of heart girth, width of chest, height at withers natural length and width of hind cannon on a live animal to estimate live and carcass weight.
3. Determination of carcass weight with weighing.
4. Giving carcass quality scores paying special attention to fleshiness.
5. Weighing the head, heart, liver and kidney fats and eventual bowel fats to estimate the quantity and percentage of carcass fat.
6. Dissection of one flank into meat and fat selection to estimate the quantity and percentage of carcass fat.
7. Dissection of one fore or hind shank into meat and bone selection to estimate the quantity and percentage of bone and percentage of meat.

REFERENCES

- ALLEN, D. M., MERKEL R. A., MAGEE, W. T. & NELSON, R. H. 1968. Variation in some beef carcass compositional characteristics within and between selected weight and fat thickness ranges. *J. An. Sci.*, 27: 1239—1246. Ref. ABA 37: 1251.
- * — MERKEL, R. A., MAGEE, W. T. & NELSON, R. H. 1969. Relationship of physically separable muscle, fat and bone from the left side of steer carcasses to yields of retail cuts, fat trim and bone of the right side. *J. An. Sci.*, 28: 311—315.
- BLACKWELL, R. L., KNOX, J. H., SHELBY, C. E. & CLARK, R. T. 1962. Genetic analysis of economic characteristics of young Hereford cattle. *J. An. Sci.*, 21: 101—107.
- de BOER, H. BERGSTRÖM, P. L. & de ROOY, H. 1969. Methods of evaluating carcass quality in cattle. EAAP Congr. Helsinki 1969.
- du BOSE, L. E., CARTWRIGHT, T. C. & COOPER, R. J. 1967. Predicting steak and roast meat from production and carcass traits. *J. An. Sci.*, 26: 688—693.
- BRACKELSBERG, P. O., HALE, N. S., COWAN, W. A. & KINSMAN, D. M. 1968. Relationship of sectional characteristics to beef carcass composition. *J. An. Sci.*, 27: 39—44.
- * — & WILLHAM, R. L. 1968. Relationship among some common live and carcass measurements and beef carcass composition. *J. An. Sci.*, 27: 53—57.
- BRÄNNÄNG, E. 1966. En studie i avkommebedömningsteknik för tillväxt- och slaktegenskaper hos kombinerande mjölkkötttraser. *Lantbrukshögk. meddel.* 101/1968.
- * — & NILSSON, K. 1969. Principles of cattle and sheep carcass evaluation. EAAP Congr. Helsinki 1969.
- BUSCH, D. A., DINKEL, C. A. & MINYARD, J. A. 1969. Body measurements, subjective scores and estimates of certain carcass traits as predictors of edible portion in beef cattle. *J. An. Sci.*, 29: 557—566. Ref. ABA 38: 1097.
- BUSS, G. 1968. Comparison of the slaughter results for young fattening bulls of German Black Pied, Spotted Mountain and German Yellow breeds in Thuringen. *Arch. Tierz.* 11: 53—72. Ref. ABA 37: 2262.
- BUTTERFIELD, R. M. 1965. The relationship of carcass measurements and dissection data to beef carcass composition. *Res. vet. Sci.*, 6: 24—32. Ref. ABA 34: 73.
- CALLOW, E. H. 1962. The relationship between the weight of a tissue in a single joint and the total weight of the tissue in a side of beef. *Anim. Prod.* 4: 37—46.

- COLE, J. W., ORME, L. E. & KINCAID, C. M. 1960. Relationship of loin eye area, separable lean of various beef cuts and carcass measurements to total carcass lean in beef. *J. An. Sci.*, 19: 89—99.
- CUNDIFF, L. V., CHAMBERS, D., STEPHENS, D. F. & WILLHAM, R. L. 1964. Genetic analysis of some growth and carcass traits in beef cattle. *J. An. Sci.*, 23: 1133—1138.
- * —, MOODY, W. G., LITTLE, J. E. (Jr), JONES, B. M. (Jr) & BRADLEY, N. W. 1967. Predicting beef carcass cutability with live animal measurements. *J. An. Sci.*, 26: 210, Abstr. 42.
- DIETERT, W. 1969. Untersuchungen über verschiedene Prüfungsverfahren auf Mastleistung und Schlachtkörperwert beim Rind. Georg-August-Univ. Göttingen 1969.
- DINKEL, C. A., WILSON, L. L., TUMA, H. J. & MINYARD, J. A. 1965. Rations and percents as measures of carcass traits. *J. An. Sci.*, 24: 425—429.
- DUMONT, B. L., QUELTE, Le, P. & ARNOUX, J. 1961. A biometrical study of beef cattle. I. Variability in the anatomical composition of the carcass of Charolais cattle. II. Estimation of the weight of the musculature in Charolais cattle. *Ann. Zootech.* 10: 149—154, 321—326. Ref. ABA 30: 2387.
- EPLEY, R. J., HEDRICK, H. B., STRINGER, W. C. & HUTCHESON, D. P. 1970. Prediction of weight and percent retail cuts of beef using five carcass measurements. *J. An. Sci.*, 30: 872—879.
- FIELD, R. A., SCHOONOVER, C. O. & NELMS, G. E. 1966. Relationship between carcass weight and muscle, fat and bone in bull carcasses. *J. An. Sci.*, 25: 588, Abstr. 22.
- GOTTSCH, H. A., MERKEL, R. A. & MACKINTOSH, D. L. 1961. Relationship of muscle, fat, bone and some physical measurements to beef carcass cutability. *J. An. Sci.*, 20: 917, Abstr. 66.
- GRAVIR, K. 1967. Studies on Different Body Measurements as Estimators of Live and Carcass Weight in Young NRF Bulls. *Acta Agric. Skand.* 17: 217—227.
- HAMMOND, J. 1958. Zuwachs und Fleischproduktion. *Handbuch der Tierzucht* I. pp. 197—235.
- HARVEY, W. R. 1966. Least squares analysis of data with unequal subclass numbers. *ARS* 20—8. Agric. Res. Service, U. S. Dept. of Agric. 157 p.
- HENDERSON, D. W., GOLL, D. E. & KLINE, E. A. 1966 a. Relationships of muscling and finish measurements from three different groups of beef carcasses with carcass yield. *J. An. Sci.*, 25: 323—328.
- * —, GOLL, D. E. & KLINE, E. A. 1966 b. Measures of carcass yield and tenderness of two muscles in four groups of beef carcasses. *J. An. Sci.*, 25: 329—333.
- HINKS, C. J. M. & BECH ANDERSEN, B. 1968. Phenotypic aspects of growth and carcass quality in veal calves. *Anim. Prod.* 10: 331—340.
- HUTH, F. W. 1965. Typfragen und Mastmethoden beim Rind. *Z. Biol.* 82: 122—138.
- JENSEN, B. 1967. Effect of age on muscle development in cattle. *Beretn. NJF.* 13 kongr. No. 4, II: 370.
- JOHANSON, I & HILDEMAN, S. E. 1953. The relationship between certain body measurements and live and slaughter weight in cattle. *ABA* 22: 1—17.
- LAINE, J. J. 1971. Amerikkalaisen kauppiaan näkemys edullisesta naudanlihan hankinnasta. *Suomen Liha- ja Karjatalous* 1971, 1: 4—6.
- LEVY, D., HOLZER, Z. & VOLCANI, R. 1968. The effect of age and live weight of feed conversion and yield of saleable meat of intact Israeli Friesian male calves. *Anim. Prod.* 10: 325—330.
- LILJEDAHL, L-E. 1965. Estimation of the Weight of Total Muscle Tissue in Beef Carcasses. *Lantbrukshögsk. Ann.* 31: 231—236.
- LINDHÉ, B. 1968. Crossbreeding for Beef with Swedish Red and White Cattle. I. Performance under varying field conditions. *Lantbrukshögsk. Ann.* 34: 1—465.
- * — & HENNINGSSON, T. 1968. Crossbreeding for Beef with Swedish Red and White Cattle. II. Growth and feed efficiency under standardized conditions together with detailed carcass evaluation. *Lantbrukshögsk. Ann.* 34: 517—550.
- LINDSTRÖM, U & MAIJALA, K. 1970. Evaluation of performance test results for A. I. Bulls. *Acta Agric. Skand.* 20: 207—218.

- MARTIN, T. G. & STARKENBURG, R. T. 1965. Genetic correlations between beef and dairy traits in dual-purpose cattle. *World Rev. An. Prod. Special Issue 1965* 1: 45—52.
- MATHER, R. E., RANDEL, P. F. & RIMM, A. A. 1959. Relative errors in some body measurements of dairy cattle. *J. Dairy Sci.*, 43: 1890—1891.
- NEIMAN-SØRENSEN, A., NIELSEN, E. O. & LEGATES, J. E. 1965. The interrelations between growth, body measurements, and carcass quality and meat characteristics in the production. *World Rev. An. Prod. Special Issue 1965*, 1. 41—44.
- NIELSEN, E. O. 1962. Undersøgelser over kødproduktionsegenskaber ved de danske afkomprøver med tyre. Mimeogr.
- ORME, L. E., COLE, J. W. & KINCAID, C. M. 1959 a. Predicting total carcass lean in mature beef using weights of certain entire muscles. *J. An. Sci.*, 18: 1479. Abstr. 50.
- ORME, L. E., PEARSON, A. M., BRATZLER, L. J., MAGEE, W. T. & WHEELER, A. C. 1959 b. The muscle-bone relationship in beef. *J. An. Sci.*, 18: 1271—1281.
- * —, PEARSON, A. M., MAGEE, W. T. & BRATZLER, L. J. 1959 c. Relationship of live animal measurements to various carcass measurements in beef. *J. An. Sci.*, 18: 991—999.
- PIRCHNER, F. 1965. Züchtungsfragen in der Fleischerzeugung. *Z. Biol.* 82: 139—153.
- RUOHOMÄKI, H. 1967. Charolaisristeytyskokeiden tuloksia. Ref. Results of Charolais trials in Finland. *Ann. Agric. Fenn.* 6: 115—126.
- * — & VARO, M. 1967. Effect of Crossing with charolais on the beef yield of slaughter animals based on results gained in the cutting of carcasses. *Acta Agr. Fenn.* 109, 1: 155—163.
- SCHMITTEN, F. 1968. Über die Einflüsse von Endgewicht und Haltungform auf die Schlacht-Körperzusammensetzung und Fleischqualität von Mastkälbern. *Mitt. Tierhalt.*, No. 114. Ref. *ABA* 37: 2293.
- SCHÖN, L. 1969. Examination of the gross tissue composition of slaughter cattle. *Bayer. landw. Jb.*, 46: 771—827.
- SEEBECK, R. M. & TULLOH, N. M. 1966. The representation of yield of dressed carcass. *Anim. Prod.* 8: 281—288.
- SKJERVOLD, H. 1958. Registrering av kjøttproduksjonsegenskapene hos storfe. *Norges Landbrukshøgsk. Melding nr. 123.*
- SWIGER, L. A., GREGORY, K. E., SUMPTION, L. J., BREIDENSTEIN, B. C. & ARTHAUD, V. H. 1965. Selection indexes for efficiency of beef production. *J. An. Sci.*, 24: 418—424.
- TALLIS, G. M., KLOSTERMAN, E. W. & CAHILL, V. R. 1959. Body measurements in relation to beef type and to certain carcass characteristics. *J. An. Sci.*, 18: 108—115.
- THORTON, J. W. & HINER, R. L. 1965. Volume of beef round related to carcass composition. *J. An. Sci.*, 24: 301—304.
- TOUCHBERRY, R. W. & LUSH, J. L. 1950. The accuracy of linear body measurement of dairy cattle. *J. Dairy Sci.*, 33: 72—80.
- VARO, M. 1969. Über Methoden zur Bewertung der Schlachtkörperqualität bei lebenden und geschlachteten Rindern. *EAAP Congr. Helsinki 1969.*
- WEBER, F. 1957. Die statistischen und genetischen Grundlagen von Körpermessungen am Rind. *Z. Biol.* 69: 225—260.
- WENIGER, J. H. 1965. Zusammensetzung und Bewertung des Schlachtkörpers. *Z. Biol.* 82: 199—217.
- WILSON, L. L., DINKEL, C. A., TUMA, H. J. & MINYARD, J. A. 1964. Live-animal prediction of cutability and other beef carcass characteristics by several judges. *J. An. Sci.* 23: 1102—1107.
- WISMER-PEDERSEN, J. 1969. Undersøgelser af kødkonsistens og muskelfibre. *Afkomprøver for kødproduktion II.* 372. beretn. fra førsøgslab. s. 72—103.
- WITT, M. 1965. Rind und Schwein als Fleischproduzenten. *Z. Biol.* 82: 103—121.

SELOSTUS

Nuoren lihanaudan teurasominaisuuksien arvioimisesta

HILKKA RUOHOMÄKI

Maatalouden tutkimuskeskus, Kotieläinjalostuslaitos, 01301 Vantaa 30

Tutkimuksen tarkoituksena oli selvittää, voidaanko elävän eläimen mitoilla arvioida riittävällä luotettavuudella elopainoa, teuraspainoa ja ruohon koostumusta — lihan, rasvan ja luiden kilo- ja prosenttimääriä — ja voidaanko ruhon koostumusta arvioida riittävällä luotettavuudella muilla keinoin kuin paloittelemalla koko ruho. Aineiston muodosti yht. 205 n. vuoden vanhaksi kasvatettua härkää ja sonnia sekä 174 n. $\frac{1}{2}$ -vuotiaaksi kasvatettua sonnivasikkaa. Pienimmän neliösumman menetelmällä lasketuilla korjaustermeillä korjattiin kukin ominaisuus yksilöittäin kokeiden sisällä iän, ruokinnan ja rodun suhteen. Korjatuista arvoista laskettiin askeltavalla regressioanalyysillä ne ominaisuudet tai ominaisuusyhdistelemät, jotka parhaiten selittivät edellä mainittuja arvioitavia ominaisuuksia.

Elopainoille saatiin elävän eläimen mitoilla 1-vuotiailla 87 %:n ja $\frac{1}{2}$ -vuotiailla 82 %:n sekä teuraspainoille vastaavasti 84 %:n ja 80 %:n selitysasteet. Parhaat elävän eläimen mitat näiden ominaisuuksien sekä lihan kilomäärän arvioimisessa olivat rinnan ympäryksen, rinnan leveys ja pituus. Lihan kilomäärää arvioi parhaiten teuraspaino, selitysaste molemmissa ikäryhmissä 96 %. Muut ominaisuudet lisäsivät teuraspainolla saatua selitysastetta 1—2 %-yksikköä. Lihan prosenttimäärän selitysaste jäi pieneksi 1-vuotiailla, mutta $\frac{1}{2}$ -vuotiailla saatiin etupotkien paloittelutuloksilla 56 %:n selitysaste. Rasvan kilo- ja prosenttimäärää arvioivat 1-vuotiailla parhaiten teuraspaino ja elimet, selitysasteet 62 % ja 50 %, kun taas $\frac{1}{2}$ -vuotiailla vastaavat selitysasteet olivat 70 % ja 59 %, jotka saatiin munuaisrasvoilla ja kupeiden paloittelutuloksilla. Luiden kilo- ja prosenttimäärälle saatiin 1-vuotiailla potkien paloittelutuloksilla 68 %:n ja 55 %:n selitysasteet sekä $\frac{1}{2}$ -vuotiailla etupotkien paloittelutuloksilla vastaavasti 82 %:n ja 70 %:n selitysasteet. Etu- ja takapään paloittelutuloksilla saadut selitysasteet olivat suurempia kuin muilla ominaisuusryhmillä saadut. Teurasarvostelupisteillä saadut selitysasteet jäivät edellä selostettuja pienemmiksi.

Taulukoissa 33 ja 34 esitettyjen eri ominaisuusryhmillä saatujen selitysasteiden — lukuunottamatta etu- ja takapään paloittelulla saatuja selitysasteita — saavuttaminen edellyttää seuraavia toimenpiteitä:

1. Elopainon toteaminen punnitsemalla.
2. Elävän eläimen rinnan ympäryksen, rinnan leveyden, pituuden, säkäkorkeuden ja takasään leveyden mittaaminen elopainon ja teuraspainon arvioimiseksi.
3. Teuraspainon toteaminen punnitsemalla.
4. Teurasarvostelussa lihakuuden arvioiminen.
5. Pään, sydämen, maksan ja munuaisrasvojen punnitseminen rasvan kilo- ja prosenttimäärän arvioimiseksi.
6. Toisen kuvekappaleen leikkaaminen liha- ja rasvalajitelmiin rasvan kilo- ja prosenttimäärän arvioimiseksi.
7. Toisen etu- tai takapotkan leikkaaminen liha- ja luulajitelmiin lihan prosenttimäärän sekä luiden kilo- ja prosenttimäärän arvioimiseksi.