

Faunal communities of the field stratum and their succession in reserved fields

HEIKKI HOKKANEN and MIKKO RAATIKAINEN

University of Jyväskylä, Department of Biology

Yliopistonkatu 9, 40100 Jyväskylä 10, Finland

Abstract. The arthropod of the field stratum and small mammal fauna of 51 reserved fields were studied in Central Finland in 1974. In high summer the arthropod density was on the average 210 individuals/m², of which *Auchenorrhyncha* formed 42 %, *Hymenoptera* 18 %, *Coleoptera* 14 %, and others 26 %. Arthropods were more abundant in fields that had been reserved after leys than in fields after open cultivations. The abundance of most arthropod taxa increased as the time in reservation increased. However, the abundance of most pests of any significance decreased with years.

Sorex araneus L. and *Microtus agrestis* L. formed almost 90 % of the small mammal fauna of the fields. Their abundance was on the average much less than in the Scots pine seed orchards nearby.

The leafhopper faunas were divided into three communities. One appeared in young (1st-3rd year) fields, and the other two in older (2nd-6th year) fields, one in dry and one in moist fields. The *Apion* fauna was divided into four communities. The dependence of the described arthropod communities on the vegetational communities of the fields was weak, although some relationships were observable.

1. Introduction

To decrease the area under cultivation, the Field Reservation Act (216/1969) was enacted in 1969. By the end of 1974 about 8 % of the cultivated area in Finland had been reserved (ANON. 1975). Research on the flora and fauna of reserved fields was started in the University of Jyväskylä in 1973. Several papers have been published on the subject, e.g. TÖRMÄLÄ and HOKKANEN (1976), TÖRMÄLÄ and RAATIKAINEN (1976), HOKKANEN and RAATIKAINEN (1977), TÖRMÄLÄ (1977). The purpose of this study is to identify the most important animal taxa and faunal communities of the field stratum, as well as their succession in reserved fields.

Relatively little information about the fauna of reserved or abandoned fields in Finland is presented in the literature, the works of GYLLENBERG (1969), KOPONEN (1972), TÖRMÄLÄ and RAATIKAINEN (1976) and TÖRMÄLÄ (1977) are almost the only ones from Finland. Much more is known about the fauna of cultivated fields, although most works deal only with one distinct group of animals. E.g. MARKKULA and MYLLYMÄKI (1958), RAATIKAINEN (1971), HUHTA

and RAATIKAINEN (1974), LAITINEN and RAATIKAINEN (1975), and RAATIKAINEN and VASARAINEN (1976) present information from Finland; similar studies have been made e.g. in Sweden (JÜRISOO 1964) and Poland (ANDRZEJEWSKA 1971, KAJAK et al. 1971, OLECHOWICZ 1971, etc).

2. Material and methods

2.1. Study area

The study area consisted of the following communes: Jyväskylä, the rural commune of Jyväskylä, Petäjävesi and Uurainen (about 62° N and 25° E) in Central Finland. 20 reserved farms were selected at random, and from each 2–3 homogenous fields again at random. Thus 51 fields were obtained, which were reserved as follows: 13 in 1969 (= 6th year fields), 5 in 1970, 6 in 1971, 11 in 1972, 10 in 1973 and 6 in 1974 (= 1st year fields). Further details about the fields and the study area are given in HOKKANEN and RAATIKAINEN (1977).

2.2. Sampling

Arthropods were sampled in early summer (4–11. VI 1974) with standard sweep-nets described by HEIKINHEIMO and RAATIKAINEN (1962). Four samples, 15 sweeps each, were taken from each field between 10 a.m. and 4 p.m. The samples were not taken if it was rainy or if there was a gale, or the vegetation was wet. The animals were killed with diethyl ether, stored in a dry place, and sorted later by hand according to species (the most numerous or otherwise important species) or group.

In high summer (16–24. VII 1974) quantitative samples were taken with Burkard suction apparatus from a circular metal enclosure with an area of 0.10 m² placed on the ground. A suction time of 1.5 minutes was used. 4 randomly located samples were taken from each field, they were treated in the same way as the sweep-net samples. The nomenclature of *Auchenorrhyncha* is according to LINNAVUORI (1969), and that of the *Apions* according to HANSEN (1965)

The abundance of small mammals was studied with the small quadrat-method described by MYLLYMÄKI et al. (1971). Trapping was conducted between 21–26. VIII. 1974 so that one quadrat was located per field. The whole material consisted of the catch from 1224 trap-nights. The nomenclature of small mammals is according to SIIVONEN (1974).

2.3. Treatment of data

Faunal communities were differentiated by means of the modified Sørensen's quotient of similarity:

$$QS = 100 \times \frac{\sum 2c}{\sum (a + b)}, \text{ where}$$

a = the amount of a species in sample 1

b = the amount of the same species in sample 2

c = the smaller of these two values (a and b)

The following symbols for the levels of significance are used in all the tests: p* < 0.05, p** < 0.01, p*** < 0.001.

3. Results

3.1. Taxa, their occurrence and abundance

3.1.1. Arthropods

The 202 sweep-net samples which were taken, contained a total of 13 640 arthropod specimens. Table 1 shows the number of individuals in the different arthropod taxa/60 sweeps, according to different age classes and previous use

Table 1. Number of arthropods in sweep-net samples according to the age and previous use of the fields. 1-6 = age as a reserved field in years, O = fields reserved after open cultivations, L = fields reserved after ley or pasture, n = number of samples.

Taxon	Number of individuals/60 sweeps											All fields
	01	02	03	04	05	06	L1	L2	L3	L4-5	L6	
<i>Araneida</i>	2.5	22.3	14.9	12.0	23.7	21.5	11.5	12.3	16.0	20.0	25.3	17.0
<i>Phalangida</i>	—	—	0.1	—	—	—	—	—	—	—	—	0.0
<i>Ephemeroptera</i>	—	—	0.6	—	0.3	0.2	0.5	—	2.0	—	—	0.3
<i>Plecoptera</i>	—	0.8	0.1	—	—	—	—	—	—	—	—	0.1
<i>Orthoptera</i>	0.8	0.3	0.1	2.5	2.7	1.2	—	0.2	1.0	0.8	1.7	1.0
<i>Thysanoptera</i>	5.3	3.8	7.3	42.0	25.3	2.2	1.0	154.3	125.3	8.8	4.1	36.0
<i>Homoptera</i>												
<i>Auchenorrhyncha</i>	1.8	10.8	14.4	27.0	40.4	64.2	18.5	42.7	34.5	72.3	111.0	44.1
<i>Balclutha punctata</i> .	0.3	2.5	10.7	4.5	22.7	4.7	8.0	9.0	0.3	3.0	3.1	6.0
<i>Diplocolenus abdo-</i>												
<i>minalis</i>	—	2.3	—	3.5	3.7	5.5	—	10.7	8.5	6.3	22.0	6.7
<i>Doliotettix pallens</i> .	—	2.3	0.9	9.0	2.0	38.2	2.5	7.2	15.1	42.3	47.0	17.5
<i>Javesella</i> spp	1.0	—	0.3	3.0	1.3	2.0	2.5	4.2	0.3	1.8	3.4	1.9
<i>Megadelphax sordi-</i>												
<i>dula</i>	—	3.0	0.7	5.5	4.4	4.5	3.5	6.8	9.1	4.8	28.2	7.5
<i>Stiroma bicarinata</i> .	—	0.5	—	—	1.0	6.2	—	0.2	—	10.8	3.9	2.2
<i>Psyllina</i>	5.5	2.5	6.1	4.8	39.3	3.8	1.5	3.8	4.0	3.5	3.3	6.2
<i>Heteroptera</i>	4.0	12.8	3.5	10.1	64.7	10.3	21.0	8.9	12.1	22.8	10.2	13.6
<i>Coleoptera</i>	5.6	19.0	17.7	29.3	68.0	67.8	28.0	33.5	137.3	74.5	45.6	46.5
<i>Apion</i> spp.	0.3	3.5	3.0	3.5	5.0	2.7	1.0	5.7	1.5	9.8	5.4	3.9
<i>Chaetocnema</i> spp .	1.0	0.5	1.3	—	—	—	—	2.7	—	0.3	—	0.6
<i>Longitarsus</i> spp. ...	0.3	1.8	2.0	—	0.3	—	—	1.2	—	0.3	—	0.6
<i>Phyllotreta</i> spp. ...	1.0	1.0	0.9	—	—	0.3	1.5	1.8	0.3	—	—	0.6
<i>Sitona</i> spp.	0.3	1.8	0.6	—	3.0	0.7	1.5	0.2	—	1.3	1.9	0.9
<i>Galerucella tenella</i> .	—	—	—	5.5	9.3	3.3	15.0	0.2	1.3	52.3	7.9	7.3
<i>Polydrosus mollis</i> ..	—	2.0	2.3	12.0	3.7	50.2	—	7.7	128.0	1.8	8.1	19.7
<i>Agriotes obscurus</i> ...	—	—	—	0.3	—	—	—	0.2	—	—	—	0.0
<i>Corymbites incanus</i>	—	—	1.4	—	—	—	—	0.3	—	—	—	0.2
<i>C. pectinicornis</i>	—	0.3	—	—	—	—	—	0.2	—	0.3	—	0.1
<i>Dolopius margina-</i>												
<i>tus</i>	—	—	—	0.3	0.3	0.2	—	0.5	—	—	—	0.1
<i>Hymenoptera</i>	3.0	14.0	12.1	26.3	100.0	37.7	16.5	61.5	31.0	59.0	28.7	34.3
<i>Lepidoptera</i>	—	—	0.1	1.3	1.0	0.5	—	0.5	0.3	0.5	0.4	0.4
<i>Diptera</i>	30.3	54.0	39.7	45.8	83.3	72.5	152.0	154.2	39.8	58.0	35.0	65.6
<i>Amaurosoma armil-</i>												
<i>latum</i>	—	0.3	0.6	1.8	—	0.5	1.5	2.2	1.0	1.3	0.6	0.9
<i>A. flavipes</i>	0.5	8.8	4.1	3.5	4.0	1.3	1.0	6.0	7.3	5.0	1.1	3.8
<i>Insecta, larvae</i>	0.5	0.8	1.1	4.8	5.0	2.3	5.0	1.7	2.5	4.3	2.1	2.4
n	16	16	28	15	12	23	8	24	16	16	28	202

Table 2. Number of arthropods in suction samples. Explanations as in Table 1.

Taxon	Number of individuals/m ²											All fields
	O1	O2	O3	O4	O5	O6	L1	L2	L3	L4-5	L6	
<i>Araneida</i>	4.4	12.5	8.2	15.6	41.7	14.6	8.8	12.9	20.6	20.0	23.6	16.1
<i>Phalangida</i>	—	0.6	0.4	—	—	—	—	—	0.6	—	—	0.2
<i>Ephemeroptera</i>	—	—	0.4	—	—	—	—	—	—	—	—	0.0
<i>Plecoptera</i>	—	0.2	—	—	—	—	—	—	—	—	0.7	0.1
<i>Orthoptera</i>	—	—	—	—	—	—	—	—	0.6	—	0.4	0.1
<i>Thysanoptera</i>	6.9	—	—	—	2.5	—	—	0.4	3.1	—	0.4	1.0
<i>Homoptera</i>												
<i>Auchenorrhyncha</i> , lar- vae	3.1	33.1	19.3	54.4	57.5	38.3	46.3	30.8	80.0	68.1	73.9	44.9
<i>Agallia</i> spp	0.6	5.6	1.8	0.6	4.2	3.6	1.3	—	1.9	2.5	2.9	2.3
<i>Auchenorrhyncha</i> , adults	6.9	41.3	27.9	47.5	33.3	27.5	17.5	25.8	51.9	96.9	79.3	42.8
<i>Aphrodes bifasciatus</i>	0.6	1.9	1.1	1.9	5.8	—	—	3.8	5.6	—	1.4	1.9
<i>Arthaldeus pascuel- lus</i>	—	0.8	0.6	1.3	—	0.8	1.0	0.8	5.0	4.3	3.7	1.7
<i>Dicranotropis ha- mata</i>	—	—	—	3.1	1.7	0.8	—	—	3.8	4.4	0.7	1.2
<i>Diplocolenus abdo- minalis</i>	1.3	11.9	5.7	15.0	9.2	12.1	6.3	4.2	15.6	31.3	27.1	13.1
<i>Doratura stylata</i> ...	—	—	—	—	0.8	0.4	—	—	—	3.1	4.3	0.9
<i>Elymana sulphurella</i>	—	0.6	4.3	—	—	0.4	—	0.4	0.6	2.5	5.0	1.7
<i>Evacanthus inter- ruptus</i>	0.6	6.3	6.8	10.6	3.3	1.7	3.8	4.2	6.3	5.0	3.6	4.7
<i>Paluda flaveola</i> ...	—	—	—	10.6	9.2	3.8	—	1.7	3.8	29.4	17.1	7.0
<i>P. preysleri</i>	—	0.6	—	1.3	0.8	2.5	—	—	—	4.4	5.0	1.5
<i>Philaenus spuma- rius</i>	1.9	1.9	1.1	1.9	—	0.4	1.3	3.8	0.6	1.3	0.7	1.4
<i>Stiroma bicarinata</i> .	—	1.3	0.4	6.3	0.8	1.7	1.3	0.4	1.3	5.0	2.5	1.8
<i>Streptanus sordidus</i>	0.6	3.1	1.4	3.1	—	0.4	5.0	1.3	4.4	3.8	7.5	2.8
<i>Xanthodelphax fla- veola</i>	—	0.6	0.4	0.6	9.2	0.4	—	—	—	1.9	3.6	1.4
<i>Aphidina</i>	0.6	—	—	—	3.1	—	1.3	0.4	3.1	—	0.7	0.7
<i>Psyllina</i>	—	0.6	1.4	—	0.8	—	—	—	1.9	—	0.7	0.5
<i>Heteroptera</i>	1.9	12.5	3.9	23.8	32.5	18.8	15.1	14.6	25.1	36.9	24.7	18.2
<i>Coleoptera</i>	33.1	20.0	16.8	21.2	29.2	42.1	26.3	14.6	14.4	84.4	24.3	28.6
<i>Apion</i> spp	—	8.8	7.5	8.8	10.8	6.7	2.5	10.8	3.1	24.4	12.9	9.1
<i>A. apricans</i>	—	3.8	2.1	0.6	—	—	1.3	2.5	—	0.6	4.3	1.6
<i>A. cerdo</i>	—	—	0.4	2.5	1.7	—	—	—	—	1.9	0.4	0.5
<i>A. curtirostre</i>	—	—	0.7	1.3	0.8	2.1	1.3	—	0.6	1.3	0.7	0.8
<i>A. facetum</i>	—	—	—	1.9	5.8	0.4	—	—	—	6.9	1.1	1.2
<i>A. flavipes</i>	—	0.6	0.4	—	0.8	—	—	0.8	—	0.6	2.1	0.6
<i>A. simile</i>	—	1.9	1.8	0.6	0.8	1.3	—	5.8	0.6	1.9	0.4	1.6
<i>A. viciae</i>	—	—	1.1	1.9	0.8	0.8	—	0.4	1.3	7.5	1.4	1.4
<i>Hymenoptera</i>	25.0	14.3	13.2	9.4	65.0	20.0	17.5	25.4	20.0	70.6	110.7	37.8
<i>Formicoidea</i>	3.1	4.4	5.7	1.3	15.8	5.4	7.6	17.2	6.9	32.5	100.7	21.0
<i>Lasius niger</i>	—	3.8	2.5	—	3.3	0.4	—	6.7	—	0.6	1.4	1.9
<i>Myrmica laevinodis</i>	3.1	0.6	3.2	1.3	11.7	5.0	6.3	1.7	5.6	22.5	98.9	18.3
<i>Lepidoptera</i>	—	2.0	—	—	—	1.7	—	—	—	2.5	1.1	0.7
<i>Diptera</i>	20.0	22.5	20.7	16.3	13.3	6.3	12.5	13.8	12.5	14.4	10.4	14.6
<i>Insecta</i> , larvae	—	3.8	5.0	8.1	5.0	2.1	6.3	5.0	1.3	1.3	2.5	3.5
n	16	14	27	16	12	22	8	20	14	16	28	193

of the fields. The 193 suction samples contained 4 280 specimens. According to Table 2, the most abundant group was *Auchenorrhyncha*. Second in abundance was *Hymenoptera*, mainly because of *Myrmica laevinodis*.

The numbers of individuals of the most important species or families of *Auchenorrhyncha* and *Coleoptera* are given in Tables 1 and 2. Table 3 shows also the frequencies of some pest groups in sweep-net samples.

Table 3. Frequencies (%) of some pest-groups in sweep-net samples (early June). Explanations as in Table 1.

Taxon	01	02	03	04	05	06	L1	L2	L3	L4-5	L6	All fields
<i>Chaetocnema</i> spp.	25	13	25	—	—	—	—	33	—	6	—	11
<i>Longitarsus</i> spp.	6	31	25	—	8	—	—	29	—	6	—	11
<i>Phyllotreta</i> spp.	19	25	21	—	—	9	13	17	19	—	—	11
<i>Sitona</i> spp.	6	31	14	—	33	17	25	4	—	31	36	18
<i>Elatерidae</i>	—	6	21	13	8	4	—	30	—	6	—	9
<i>Galerucella tenella</i>	—	—	—	40	50	17	25	5	13	50	32	19
<i>Polydrosus mollis</i>	—	19	25	53	33	57	—	45	25	31	50	33
<i>Amaurosome</i>												
<i>armillatum</i>	—	6	14	27	—	13	13	50	19	31	14	18
<i>A. flavipes</i>	13	94	46	60	58	35	25	55	81	69	21	49
n	16	16	28	15	12	23	8	24	16	16	28	202

3. 1. 2. Small mammals

The total catch of small mammals was 167 individuals, 13.6 per 100 trap-nights. The species composition and other data are given in Table 4. *Sorex araneus* and *Microtus agrestis* formed about 90 % of the total catch, other species occurred only accidentally. On 16 % of the quadrats the catch of small mammals was zero individuals, one individual was caught on 14 % of the quadrats, two on 22 %, three on 2 %, four on 18 %, five on 10 %, six on 6 %, seven on 2 %, eight on 4 %, and ten, eleven or twelve individuals per quadrat each on 2 % of them.

Table 4. The catches of small mammals. n = number of captured individuals, F% = frequency-% in quadrats, \bar{x} = average number of individuals per quadrat, N = number of individuals per 100 trap-nights.

	n	F%	\bar{x}	N
<i>Sorex araneus</i>	96	80	1.9	7.8
<i>Microtus agrestis</i>	53	38	1.0	4.3
<i>Clethrionomys glareolus</i>	6	6	0.1	0.5
<i>Micromys minutus</i>	3	6	0.1	0.3
<i>Sorex isodon</i>	3	6	0.1	0.3
<i>Neomys fodiens</i>	2	4	0.0	0.2
<i>Arvicola terrestris</i>	2	2	0.0	0.2
<i>Apodemus flavicollis</i>	2	2	0.0	0.2
Total	167	84	3.3	13.6

3. 2. Factors affecting abundances

Tables 1–3 and Fig. 1 show the changes in the abundances of the main taxa as a function of the age of the field (as a reserved field). According to the results of the sweep-net samples, the abundances of *Auchenorrhyncha* ($r = .587^{***}$), *Araneida* (.443**), *Hymenoptera* (.430**), *Coleoptera* (.395**), and *Heteroptera* (.364*) increase with time. The only negative correlations observed, were: *Phyllotreta* spp. (–.498***), *Chaetocnema* spp. (–.385**), and *Longitarsus* spp. (–.283*). Results of the suction samples also showed significant positive correlations with age for *Auchenorrhyncha* (.440**), *Araneida* (.432**), and *Heteroptera* (.356*), and a negative one for *Diptera* (–.402**) (Fig. 1).

As to the previous use of the fields, leafhoppers and bugs were significantly more abundant on fields that had been reserved after leys or pastures than on fields after open cultivations. No taxon was significantly more abundant after open cultivations than after leys.

The moisture conditions of the fields were evaluated, and some groups showed differences in abundance according to the wetness of the field. For arthropods, all the correlations showed greater abundance on drier fields, whereas small mammals were more abundant on moist fields (.334*). *Coleoptera* (.374**), *Thysanoptera* (.336*), *Heteroptera* (.342*) and some *Auchenorrhyncha* species (*Dicranotropis hamata* .421**, *Doratura stylata* .347**, *Elymana sulphurella* .290*) showed a greater tendency towards drier conditions than the average.




Fig. 1. Abundances of the main Arthropod taxa in mid-summer, according to the age of the fields as a reserved field.

3.3. Communities

3.3.1. Description

The leafhopper data of the sweep-net and the suction samples and the *Apion* data of the suction samples were treated with the Sørensen's quotient of similarity method to differentiate seasonal aspects and communities. Three early summer aspects could be recognized from the leafhopper fauna of the sweep-net samples; their species composition, dominance relations, and succession are given in Table 5 and Fig. 2.

1) *Balclutha punctata* aspect appeared mainly on 1st–3rd year fields, which were situated usually on coarse mineral ground. *Balclutha*, which hiberna-

Table 5. Dominance relations in the early summer aspects of leafhopper communities. A = *Balclutha punctata* aspect, B = *Megadelphax sordidula* aspect, C = *Doliotettix pallens* aspect, n = number of leafhoppers on the average.

	A	B	C
n/60 sweeps	17.0	24.7	128.9
Dominance %			
<i>Balclutha punctata</i>	64	19	3
<i>Delphacodes venosus</i>	0	—	0
<i>Diplocolenus abdominalis</i>	5	27	11
<i>Doliotettix pallens</i>	10	19	69
<i>Javesella</i> spp.	15	3	2
<i>Megadelphax sordidula</i>	6	30	9
<i>Stiroma bicarinata</i>	1	1	6




Fig. 2. The occupation of fields of different ages by the early summer aspect of leafhopper communities, as a percentage of the maximum number of fields. A = *Balclutha punctata* aspect, B = *Megadelphax sordidula* aspect, C = *Doliotettix pallens* aspect.

tes as an adult, formed 64 % of the leafhopper fauna of the aspect, other important species were *Javesella* spp. and *Doliotettix pallens*. This aspect occupied about 35 % of the reserved fields in the study area.

2) *Megadelphax sordidula* aspect appeared on 2nd–6th year fields, which were situated usually on mineral ground. Moisture conditions in these fields were mostly intermediate or dry. *Diplocolenus abdominalis*, *Doliotettix pallens* and *Balclutha punctata* are the other three dominant species (D % > 15). This aspect occurred on about 16 % of the fields.

3) *Doliotettix pallens* aspect shows the greatest abundance of leafhoppers of the aspects described, the number of leafhoppers per 60 sweeps was about 5–10 fold in comparison with the other aspects. It appeared on older, 3rd–6th year fields, favouring moist fields on organic soils. Besides *Doliotettix pallens*, no other dominants occurred in this aspect, second in abundance was *Diplocolenus abdominalis*. This aspect occupied about 31 % of the fields.

The leafhopper fauna of high summer (suction samples) was also divided into three fairly distinct midsummer aspects, whose species composition, dominance relations and succession are given in Table 6 and Fig. 3.

Table 6. Dominance relations in the high summer aspect of leafhopper communities. A = *Evacanthus interruptus* aspect, B = *Diplocolenus abdominalis* aspect, C = *Paluda flaveola* aspect, n = number of adult leafhoppers on the average.

	A	B	C
n/m ²	18.3	31.3	76.1
Dominance %			
<i>Agallia brachyptera</i>	—	—	0
<i>A. nymphs</i>	7	2	4
<i>Aphrodes bicinctus</i>	—	—	0
<i>A. bifasciatus</i>	4	14	2
<i>Arthaldeus pascuellus</i>	10	7	6
<i>Balclutha punctata</i>	—	—	1
<i>Chlorita</i> sp.	—	—	0
<i>Deltocephalus pulicaris</i>	3	—	—
<i>Dicranotropis hamata</i>	—	8	1
<i>Diplocolenus abdominalis</i>	18	43	35
<i>Doliotettix pallens</i>	—	—	2
<i>Doratura stylata</i>	—	—	1
<i>Elymana sulphurella</i>	3	1	1
<i>Eupteryx</i> sp.	—	—	0
<i>Evacanthus interruptus</i>	37	9	3
<i>Javesella forcipata</i>	—	—	1
<i>Macrosteles cristatus</i>	—	—	0
<i>Megadelphax sordidula</i>	—	2	1
<i>Muellerianella nymphs</i>	—	—	1
<i>Notus flavipennis</i>	—	—	1
<i>Paluda flaveola</i>	—	5	20
<i>P. preysleri</i>	—	—	4
<i>Philaenus spumarius</i>	14	—	2
<i>Streptanus sordidus</i>	4	2	6
<i>Stiroma bicarinata</i>	3	7	5
<i>Xanthodelphax flaveola</i>	—	—	2
<i>X. straminea</i>	—	—	0

1) *Evacanthus interruptus* aspect. This aspect appeared only on the youngest, 1st—2nd year fields, occupying about 12 % of the studied fields. Other common species were *Diplocolenus abdominalis*, *Philaenus spumarius*, and *Arthaldeus pascuellus*.

2) *Diplocolenus abdominalis* aspect was common on older, 2nd—6th year fields, situated on relatively dry mineral ground. *Diplocolenus* was the only




Fig. 3. The occupation of fields of different ages by the high summer aspect of leafhopper communities, as a percentage of the maximum number of fields. A = *Evacanthus interruptus* aspect, B = *Diplocolenus abdominalis* aspect, C = *Paluda flaveola* aspect.

dominant species, other common species were *Aphrodes bifasciatus*, *Evacanthus interruptus*, *Dicranotropis hamata*, and *Stiroma bicarinata*. This aspect occupied about 16 % of the fields.

3) *Paluda flaveola* aspect was typical on 3rd—6th year moist fields, which were most often situated on organic soils. The aspect was very diverse, and the number of leafhoppers/m² was clearly the greatest. *Paluda flaveola*, *P. preysleri*, and *Diplocolenus abdominalis* are probably the most common species of this aspect, others are e.g. *Streptanus sordidus* and *Arthaldeus pascuellus*. This aspect occupied about 26 % of the fields in high summer.

The leafhopper communities could be detected by studying the occurrences of the described aspects in the same fields at different times (Table 7). The *Balclutha punctata* aspect of early summer was replaced mainly by the *Evacanthus interruptus* aspect in high summer, and these aspects are parts of the *Evacanthus interruptus* community. Similarly the *Megadelphax sordidula* aspect changed to the *Diplocolenus abdominalis* aspect and these are parts of the *Diplocolenus abdominalis* community. The *Doliotettix pallens* aspect changed

Table 7. The relations between early and high summer aspects of the leafhopper communities, as a percentage of the maximum theoretical number of common fields in each comparison. The numbers in brackets indicate the number of fields occupied by this aspect.

	<i>Evacanthus interruptus</i> (6)	<i>Diplocolenus abdominalis</i> (8)	<i>Paluda flaveola</i> (13)
<i>Balclutha punctata</i> (17)	50	38	16
<i>Megadelpfax sordidula</i> (15)	17	63	39
<i>Doliotettix pallens</i> (8)	0	0	63

Table 8. Dominance relations in the *Apion* communities. A = *Apion simile* community, B = *Apion viciae* community, C = *Apion facetum* community, D = *Apion apricans* community, n = number of individuals on the average.

	A	B	C	D
n/m ²	13.7	7.3	20.5	20.0
Dominance %				
<i>Apion apricans</i>	33	—	—	50
<i>A. cerdo</i>	—	5	19	—
<i>A. curtirostre</i>	2	10	5	4
<i>A. ervi</i>	—	5	—	8
<i>A. facetum</i>	—	—	51	4
<i>A. flavipes</i>	6	5	—	25
<i>A. marchicum</i>	4	—	—	—
<i>A. simile</i>	43	5	7	—
<i>A. viciae</i>	4	50	10	—

to the *Paluda flaveola* aspect in high summer, and these are parts of the *Paluda flaveola* community.

The coleopteran genus *Apion* showed almost identical distribution in the sweep-net and the suction samples, and therefore no early and high summer aspects of the communities were separated. The QS method was applied to the data of high summer, and four different communities were identified (Table 8 and Fig. 4):

1) *Apion simile* community appeared mainly on young, 1st—3rd year reserved fields. These fields were most often intermediate in moisture, and were situated on coarse mineral ground. This was the most common of the *Apion* communities, appearing on 20 % of the fields.

2) *Apion viciae* community appeared on 3rd—6th year fields, which were classified as wet or intermediate as to moisture, and which were often situated on organic soils. The abundance of the *Apion* species was distinctly lower in these fields than in the drier fields, only 7.3 individuals/m² on the average. 10 % of the fields were representative of this community.

3) *Apion facetum* community was typical of old, 5th—6th year fields, which were dry or intermediate as to moisture. This community occupied about 14 % of the fields.

4) *Apion apricans* community was another community of old, dry fields situated on mineral ground. This community occupied about 6 % of the fields, all of which were 6th year fields.

About one third to one half of the fields ranked outside the presented classifications, partly because of the scarcity of animals in samples (especially in those from 1st year fields), and partly because their leafhopper and *Apion* communities were mixtures of the described communities, and thus could not be clearly differentiated with this method.




Fig. 4. The occupation of fields of different ages by the *Apion* communities, as a percentage of the maximum number of fields. A = *Apion simile* community, B = *Apion viciae* community, C = *Apion facetum* community, D = *Apion apricans* community.

3.3.2. Relations to vegetational communities

The vegetational communities of the studied fields have been described earlier (HOKKANEN and RAATIKAINEN 1977). Table 9 shows the relations between them, the faunal aspects of the communities, and the faunal communities described in this work. Of the early summer aspects of leafhopper communities only the *Megadelphax sordidula* aspect showed a fairly clear affinity towards a vegetational community, the *Phleum pratense* community. The *Doliotettix pallens* aspect appeared almost equally on all older fields, seemingly independently of the vegetation communities. The high summer aspects of the leafhopper communities were more clearly associated to vegetational communities. The *Evacanthus interruptus* aspect was best represented on fields of the *Phleum* type. The *Diplocolenus abdominalis* aspect also favoured the *Phleum* type, but was frequent in the *Elytrigia* type fields as well. The greatest affinity ob-

Table 9. The relations between vegetational and faunal communities, as a percentage of the maximum theoretical number of common fields in each comparison. The numbers in brackets indicate the number of fields occupied by the community.

	<i>Galeopsis</i> type (4)	<i>Phleum</i> type (8)	<i>Elytrigia</i> type (7)	<i>Anthoxanthum</i> type (11)	<i>Deschampsia</i> type (5)
<i>Balclutha punctata</i> (17)	0	25	14	18	20
<i>Megadelphax sordidula</i> (15)	0	50	29	36	20
<i>Doliotettix pallens</i> (8)	0	0	43	25	40
<i>Evacanthus interruptus</i> (6) .	25	50	17	17	0
<i>Diplocolenus abdominalis</i> (8)	0	38	29	13	0
<i>Paluda flaveola</i> (13)	0	0	29	36	80
<i>Apion simile</i> (10)	0	50	14	20	0
<i>Apion viciae</i> (7)	0	0	14	14	29
<i>Apion facetum</i> (5)	0	0	20	40	20
<i>Apion apricans</i> (3)	0	0	0	67	0

served was that of the *Paluda flaveola* aspect to *Deschampsia* type fields; almost all *Deschampsia* fields were occupied by this leafhopper aspect.

All the *Apion* communities coincided with a vegetational community. The *Phleum* type was occupied by the *Apion simile* community — no other *Apion* communities were observed in those fields. The *Apion apricans* community appeared in *Anthoxanthum* type fields, as well as the *Apion facetum* community, but the latter was also observed in *Elytrigia* and *Deschampsia* fields, although to a lesser extent. The *Apion viciae* community favoured *Deschampsia* type fields, but occurred also in others except in the youngest successional communities.

4. Discussion

The occurrence of animals harmful to surrounding cultivations in reserved fields has been of great concern. Among the pests of cultivated plants, as listed by VAPPULA (1965), the following insect species or taxa were met in this study: *Philaenus spumarius*, *Evacanthus interruptus*, *Javesella* spp., *Dicranotropis hamata*, *Corymbites pectinicornis*, *Galerucella tenella*, *Phyllotreta* spp., *Longitarsus* spp., *Chaetocnema* spp., *Apion flavipes*, *A. apricans*, *Polydrosus mollis*, *Sitona* spp., *Amaurosoma flavipes*, and *A. armillatum*. According to RAATIKAINEN (1970), some leafhoppers are vectors of cereal viruses or mycoplasma, causing at times severe damage to cultivations. Of these species the following were met in reserved fields: *Macrosteles cristatus*, *Javesella pellucida*, *J. obscurella*, *Dicranotropis hamata* and *Megadelphax sordidula*. The proportion of these species in the total leafhopper fauna was about 20 % in reserved fields, whereas in Finnish oats the proportion of such species is about 90 % (RAATIKAINEN and VASARAINEN 1976).

The reason for the scarcity of most of the important pests in the reserved fields is probably the small proportion of suitable host plants, in comparison

with e.g. cereals or timothy, and the competition of other species that may be better adapted to the conditions prevailing in reserved fields. The explanation may partly lie also in the distinct year-to-year fluctuations in the abundance of insects; according to MARKKULA (1975) in the summer of 1974 most pest groups were somewhat less abundant than the average. According to TÖRMÄLÄ and RAATIKAINEN (1976), the sampling times in this study coincided fairly well with the maximum abundances of arthropods, except for *Coleoptera*, which has its maximum later in the summer.

The populations of the field vole, *Microtus agrestis*, were fairly strong in the summer of 1974 (MARKKULA 1975), and especially strong in Central Finland, which can be seen from the severe damage in the winter of 1974/75 (TEIVAINEN 1975). In the study of MYLLYMÄKI and PAASIKALLIO (1976), the catch with the small quadrat method in seed orchards on forest ground was a total of 5.3 individuals per quadrat, 3.7 of them being *Microtus agrestis*. In the summer of 1974 the average catch in Central Finland was about 7–10 individuals per quadrat (Myllymäki, pers. com.). The catch from reserved fields was considerably lower, exactly 3.3 individuals per quadrat, and only 1.1 of them were *Microtus agrestis*. On about 12 % of the fields the catch of small mammals equalled the average from Scots pine seed orchards (at least 7 ex./SQ).

TÖRMÄLÄ and RAATIKAINEN (1976) describe three seasonal aspects of leafhopper communities from a reserved field. According to their timing, the seasonal aspects described in this study represent the early summer aspect and the mid-late summer aspect. The early summer aspect of TÖRMÄLÄ and RAATIKAINEN (1976) represents clearly the *Doliotettix pallens* aspect described in this study, but the mid-late summer aspect does not yet rank clearly into the *Paluda flaveola* aspect, because the samples were taken so early.

KONTKANEN (1950) gives information about the seasonal aspects of leafhopper communities as well as their biotope requirements. Also he distinguished communities living in dry, fresh and wet biotopes. Although the species composition in his study area differed widely from ours, the community he describes from fresh biotopes (the *Doliotettix pallens* community) in the early summer aspect corresponds very well with our *Doliotettix pallens* aspect. Of his high summer communities the one living on moist — fresh biotopes (the *Philaenus spumarius* — *Arthaldeus pascuellus* — *Elymana sulphurella* -community) resembles best the leafhopper fauna of reserved fields.

RAATIKAINEN and VASARAINEN (1976) describe the leafhopper communities of Finnish oat fields. None of their communities are similar to the communities on reserved fields, because the species composition is quite different.

The occurrence and the true nature of the so-called faunal communities have been widely discussed for a long time (see e.g. PRICE 1975). However, certain species do show a tendency to occur more or less together, and can thus be considered to form a community. The reason for the co-existence may be e.g. the same food plants, the same physical environment requirements, similar pressure from the enemies (predators, parasites, etc), or some other factor.

RAATIKAINEN and VASARAINEN (1976) present information about the ecological requirements, e.g. host plants, of many leafhopper species, and RAATIKAINEN et al. (1977) show the dependence of *Philaenus spumarius* on certain

plant community types. MARKKULA and MYLLYMÄKI (1958) list the host plants for many *Apion* species and show that different *Apion* faunas exist in different types of leys. KONTKANEN (1950) discussed the relation between leafhopper communities and vegetational communities, but he could not clearly show their mutual dependence. Although there is a definite relationship and a need for co-existence between e.g. a leafhopper and its host plant, these relations differ on the one hand, and overlap on the other hand, so much from one species to another that the dependence of an animal community on a vegetational community becomes much looser than the actual species-to-species relations. The data of this study likewise support this view, for although both the animal and the vegetational communities could be described relatively distinctly, their mutual dependence was rather weak, and their occurrences overlapped over a wide range.

From the changes that could be ascertained with years in the fauna, both at the species, the aspect, and the community level, it can be concluded that succession proceeds in a reserved field towards the conditions of natural meadows, and that the most drastic changes take place within a few years from reservation — after that the changes get slower and succession proceeds less vigorously. During the first years most of the important pest species decrease and the species of natural meadows increase their share. Especially important is the increasing proportion of predators (like *Araneida*, *Hymenoptera*) at the expense of their prey (e.g. *Diptera*). Similar changes were observed also in the study of the Vistula Valley in Poland (e.g. KAJAK et al. 1971).

Acknowledgements. We are grateful to Mr. Timo Törmälä, Ph. Lic., who took part in the field and laboratory work. The financial support from the National Research Council for Sciences is greatly appreciated.

REFERENCES

- ANDRZEJEWSKA, L. 1971. Productivity investigation of two types of meadows in the Vistula Valley. VI. Production and population density of leafhopper (*Homoptera* — *Auchenorrhyncha*) communities. *Ekol. Pol.* 19: 151—172.
- ANON. 1975. Monthly review of agricultural statistics 1975, 1: 1—56.
- GYLLENBERG, G. 1969. The energy flow of *Chorthippus parallelus* (Zett.) (*Orthoptera*) population on a meadow in Tvärminne, Finland. *Acta Zool. Fenn.* 123: 1—74.
- HANSEN, V. 1965. Biller XXI. Snudebiller. Danmarks fauna 69: 1—524. København.
- HEIKINHEIMO, O. & RAATIKAINEN, M. 1962. Comparison of suction and netting methods in population investigations concerning the fauna of grass leys and cereal fields, particularly in those concerning the leafhopper *Calligypona pellucida* (F.). *Publ. Finn. State Agric. Res. Board* 191: 1—31.
- HOKKANEN, H. & RAATIKAINEN, M. 1977. Yield, vegetation and succession in reserved fields in Central Finland. *J. Scient. Agric. Soc. Finl.* 49: 221—238.
- HUHTA, V. & RAATIKAINEN, M. 1974. Spider communities of leys and winter cereal fields in Finland. *Ann. Zool. Fenn.* 11: 97—104.
- JÜRISOO, V. 1964. Agro-ecological studies on leafhoppers (*Auchenorrhyncha*, *Homoptera*) and bugs (*Heteroptera*) at Ekensgård farm in the province of Hälsingland, Sweden. *Stat. Växtskyddsanst. Medd.* 13, 101: 1—147.
- KAJAK, A., BREYMEYER, A. & PETAL, J. 1971. Productivity investigation of two types of meadows in the Vistula Valley. XI. Predatory arthropods. *Ekol. Pol.* 19: 223—233.

- KONTKANEN, P. 1950. Quantitative and seasonal studies on the leafhopper fauna of the field stratum on open areas in North Karelia. *Ann. Zool. Soc. 'Vanamo'* 13, 8: 1–91.
- KOPONEN, T. 1972. Peltomyyräpopulaation rakenteesta. *Metsäntutkimuslaitos. Metsänviljelyn Koeaseman Tied.* 4: 1–20.
- LAITINEN, T. & RAATIKAINEN, M. 1975. Seasonal aspects of beetle fauna occurring in oats in Western-Finland. *Ann. Agric. Fenn.* 14: 203–209.
- LINNAVUORI, R. 1969 a. Nivelkärsäiset IV. Kaskaat 1. *Anim. Fenn.* 12: 1–244. Porvoo–Helsinki.
- » — 1969 b. Nivelkärsäiset IV. Kaskaat 2. *Anim. Fenn.* 13: 1–312. Porvoo–Helsinki.
- MARKKULA, M. 1975. Viljelykasvien tuhoeläimet 1974. *Koetoin. ja Käyt.* 32: 9.
- » — & MYLLYMÄKI, S. 1958. The composition of the *Apion* (*Col., Curculionidae*) population of grassland legumes and some wild leguminous plants. *Ann. Ent. Fenn.* 24: 97–124.
- MYLLYMÄKI, A. & PAASIKALLIO, A. 1976. Scots pine seed depredation by small mammals, as revealed by radioactive tagging of the seeds. *Ann. Agric. Fenn.* 15: 89–96.
- » — , PAASIKALLIO, A., PANKAKOSKI, E. & KANERVO, V. 1971. Removal experiments on small quadrats as a means of rapid assesment of the abundance of small mammals. *Ann. Zool. Fenn.* 8: 177–185.
- OLECHOWICZ, E. 1971. Productivity investigation of two types of meadows in the Vistula Valley. VIII. The number of emerged *Diptera* and their elimination. *Ekol. Pol.* 19: 183–195.
- PRICE, P. W. 1975. *Insect ecology.* 514 p. New York.
- RAATIKAINEN, M. 1970. Viljojen virukset ja kaskaat. (Summary: Virus diseases of cereals and leafhoppers). *Luonnon Tutkija* 75: 65–74.
- » — 1971. Seasonal aspects of leafhopper (*Hom., Auchenorrhyncha*) fauna in oats. *Ann. Agric. Fenn.* 10: 1–8.
- » — & VASARAINEN, A. 1976. Composition, zonation and origin of the leafhopper fauna of oatfields in Finland. *Ann. Zool. Fenn.* 13: 1–24.
- » — , HALKKA, O., VASARAINEN, A. & HALKKA, L. 1977. Abundance of *Philaenus spumarius* in relation to types of plant community in the Tvärminne archipelago, southern Finland. *Biol. Res. Rep. Univ. Jyväskylä* 3: 3–38.
- SIIVONEN, L. 1974. Pohjolan nisäkkäät. 196 p. Helsinki.
- TEIVAINEN, T. 1975. Myyräkanta kehitymässä huippuun Keski-Suomessa. *Metsälehti* 42, 31: 8.
- TÖRMÄLÄ, T. 1977. Effects of mowing and ploughing on the primary production and flora and fauna of a reserved field in Central Finland. *Acta Agric. Scand.* 27: 253–264.
- » — HOKKANEN, H. 1976. Pakettipeltojen ja niiden reuna-alueiden linnustosta Keski-Suomessa. *Keski-Suomen Linnut* 1: 44–47.
- » — & RAATIKAINEN, M. 1976. Primary production and seasonal dynamics of the flora and fauna of the field stratum in a reserved field in Middle Finland. *J. Scient. Agric. Soc. Finl.* 48: 363–385.
- VAPPULA, N. A. 1965. Pests of cultivated plants in Finland. *Acta Ent. Fenn.* 19: 1–239.

Ms received January 3, 1978

Pakettipeltojen kenttäkerroksen eläinyhteisöistä ja niiden seuraannosta Keski-Suomessa

HEIKKI HOKKANEN ja MIKKO RAATIKAINEN
Jyväskylän yliopisto, Biologian laitos

Tutkimuksen tarkoituksena oli selvittää, mitä eläimiä ja eläinyhteisöjä pakettipelloilla esiintyy ja miten eläinten runsaudet ja eläinyhteisöt muuttuvat paketointiaikana. Erityistä huomiota kiinnitettiin tuhoeläimiin.

Tutkimuskohteina oli 20 tilan 51 peltolohkoa, jotka poimittiin otannalla Jyväskylästä ja sen ympäristökunnista. Kultakin lohkolta otettiin v. 1974 kesäkuun alussa neljä 15 haavinvedon näytettä ja heinäkuun puolivälissä neljä 0,10 m²:n imunäytettä kenttäkerroksen niveljalkaisista. Elokuun lopulla suoritettiin lohkoilla pikkunisäkkäiden ruutupyynti 1 224 loukkuyönä.

Keskikikesällä pakettipeltojen kenttäkerroksesta tavattiin keskimäärin 210 niveljalkaista/m². Niistä oli kaskaita 42 %, pistiäisiä 18 %, kovakuoriaisia 14 % ja muita yhteensä 26 %. Nurmen jälkeen paketoituilla pelloilla oli eläimiä runsaammin kuin avoviljelyksen jälkeen paketoituilla pelloilla. Useimpien ryhmien runsaudet kasvoivat peltosten pakettisaatolion kasvaessa, mutta lähes kaikkien merkittävien tuholaisien runsaudet kuitenkin pienenevät peltosten vanhetessa.

Pikkunisäkkäistä runsaimmat olivat metsäpäästäinen (57 % kaikista) ja peltomyyrä (32 %). Niiden tiheydet olivat kuitenkin keskimäärin paljon pienemmät pakettipelloilla kuin läheisillä männyn siemenviljelyksillä.

Kaskasaineistoista erottui kolme yhteisöä. Yksi niistä esiintyi nuorilla, (1.—3. vuotta pakettisaatolion pelloilla), yksi vanhoilla (2.—6. vuoden) kuivahkoilla ja yksi vanhoilla kosteilla pelloilla.

Alku- ja keskikikesän kausiyhteisöt vastasivat esiintymispaikoiltaan toisiaan niin hyvin, että kyseessä selvästikin olivat samojen yhteisöjen vuodenaikaisaspektit. Kaskasyhteisöjen lisäksi kuvattiin neljä nirppuyhteisöä. Minkään kuvatussa eläinyhteisön riippuvuus peltosten kaskiyhteisöistä ei ollut kovin selvä, vaan monet yhteisöt esiintyivät näennäisen itsenäisesti usealla eri kasvillisuustyypillä.

Tulokset osoittavat, että pakettipellot eivät yleensä ole merkittäviä tuhoeläinten leviämiskeskuskoita, koska vahingollisten lajien tiheydet ovat niissä alhaisia. Ne voivat kuitenkin olla joidenkin lajien, esim. timoteikärpästen, osalta tärkeitä suoja- ja lisääntymispaikkoja.

Eniten haittaa lienee peltomyyrästä, joiden tiheydet pakettipelloilla vaihtelevat pellon kosteuden, kasvillisuuden yms. mukaan. Huippuarvot eivät kuitenkaan saavuttaneet esimerkiksi männyn siemenviljelyksiltä todettuja arvoja. Paikallinen merkitys voi kuitenkin joskus olla huomattava.