

## Digestibility of amino acids in pig diets containing Eurolysine bacterial protein or Pekilo protein, with special reference to a gas chromatographic method used in amino acid determination

MATTI NÄSI<sup>1</sup> AND LEA HUIDA<sup>2</sup>

<sup>1</sup> Department of Animal Husbandry, University of Helsinki, 00710 Helsinki 71

<sup>2</sup> Institute of Animal Husbandry, Agricultural Research Centre, 36100 Jokioinen

**Abstract.** The apparent faecal digestibilities of amino acids were determined in growing pigs on diets in which 50 % or 100 % soybean meal supplement was replaced by Eurolysine bacterial protein or Pekilo protein. The trial was conducted as two 3 × 3 Latin squares with six castrated pigs. Compared with the soybean meal supplement, the Eurolysine diet had lower ( $P < 0.05$ ) digestibilities of arginine, leucine, methionine, phenylalanine and tyrosine. Compared with the Pekilo diet the Eurolysine diet had lower digestibilities ( $P < 0.05$ ) of alanine, aspartic acid, glutamic acid, glycine, leucine, proline and serine. Lysine digestibility was higher on the Eurolysine diet than on the Pekilo diet.

A gas chromatographic method used in amino acid determination is described and evaluated.

### Introduction

The value of Eurolysine bacterial protein and Pekilo protein as protein supplements in diets for growing pigs and laying hens was assessed in two experiments (NÄSI 1982, a, b) by replacing conventional protein sources at levels ranging from 0 to 100 %. In digestibility trials with growing pigs the diets supplemented with Eurolysine were found to have lower digestibilities than the diets with soybean meal or Pekilo, but combined with barley Eurolysine proved to have a high biological value. Its high lysine content (8.3 g/16 g N) makes it a suitable protein source for growing pigs. In a production trial with laying hens Eurolysine and Pekilo were used to replace soybean-fishmeal supplementation at several different levels. The performance of the hens deteriorated as the replacement level rose. In poultry feeding the sulphur-containing amino acids are of importance, and Eurolysine and Pekilo proved to be deficient in this respect.

The purpose of this study was to investigate the amino acid digestibilities of the diets containing Eurolysine bacterial protein and Pekilo protein, and to assess the precision of a gas chromatographic method in amino acid determination.


## Materials and Methods

The digestibility trial was carried out with six Landrace pigs weighing 37–65 kg. The total collection method was used and the experiment was designed as two  $3 \times 3$  Latin squares, the soybean supplement being replaced with Eurolysine bacterial protein or Pekilo protein at two levels 50 % and 100 %. The basic feed was barley milled with a 3-mm sieve. The details of the experimental procedure have been described in a paper by NÄSI (1982 a).

Amino acid analyses were performed on the diet ingredients and faeces, using a Hewlett Packard Model 5710 A gas chromatograph equipped with dual-flame ionisation detectors and a Hewlett Packard Integrator Model 3380 A. The amino acids were determined from their *n*-heptafluorobutyric (N-HFB) *n*-propyl ester derivatives, according to the procedure described by MARCH (1975). The columnpacking material, 3 % SE 30 on Gas Chrom Q 100/120 mesh, was obtained from Applied Science Lab. Glass columns of 3 m  $\times$  2 mm i.d. were used. The columns were conditioned at 250 °C overnight with a nitrogen flow rate of 30 ml/min.

Feed and faeces samples, 14 mg, were hydrolysed in SVL schrew cap tubes (10 ml) with 5 ml of 6 N HCl (constant boiling) at 110 °C for 20 h. The acid to protein ratio was 1000/l. Dry, extra pure nitrogen was bubbled into the mixture 5 min before the tube was sealed. After hydrolysis 250  $\mu$ l of 3 mmol pipercolic acid solution was added to the hydrolysate and the mixture was filtered through a Whatman GF/A glass microfibre paper. The filtrate was evaporated in a rotary evaporator at 30°C and the residue was taken up in 1 ml of 0.1 N HCl. The solution was cleaned up by passing it through an ion-exchange column, 40  $\times$  5 mm, made of Dowex 50 W  $\times$  8, 200–400 mesh. Before the cation-exchange resin was put into the column, it was placed in a Erlenmeyer flask, and washed successively with 7 N NH<sub>4</sub>OH, double distilled water (to neutral pH), 3 N HCl and double distilled water (to neutral pH). Impurities from the sample were washed out of the column after the sample solution by passing through 4 ml of 0.01 N HCl followed by 2 ml of water. Then the amino acid fraction was eluted with 0.5 ml of 2 N NH<sub>4</sub>OH followed by 4 ml of 7 N NH<sub>4</sub>OH and 1 ml of water and collected in an evaporation flask (50 ml). The standard amino acid solution was purified daily by ion exchange in the same way. After evaporation in a rotary evaporator under reduced pressure at 30°C, the *n*-heptafluorobutyric *n*-propyl esters were derived from the amino acids. Figure 1 shows a gas chromatogram of the *n*-heptafluorobutyric *n*-propyl derivatives of 18 amino acids. Each amino acid peak and the pipercolic acid peak represent approximately 0.05  $\mu$ mol. Pipercolic acid is used as internal standard. The temperature was raised from 90°C to 230°C at a rate of 4°/min; the temperature of the

Figure 1. Gas chromatogram of the N-HFB n-propyl derivatives of 18 amino acids. Each amino acid peak and the pipelicolic peak represents approximately 0.05  $\mu$ mol. Pipelicolic acid is used as internal standard. The temperature was raised from 90°C to 230°C at a rate of 4°C/min; the detector and injection block temperature was 250°C. The flow rate of the carrier  $N_2$  was 30 ml/min.



detector and injection block was 250°. The flow rate of the carrier  $N_2$  was 30 ml/min.

The precision of the gas-liquid chromatographic analysis was estimated from five independent analyses performed during one month on a standard amino acid solution and from seven analyses of hydrolysates obtained by repeated hydrolysis of the same wheat sample during a period of three months.

## Results and Discussion

Table 1 shows the precision of the ion-exchange cleanup, derivation and gas-liquid chromatographic analysis obtained with the standard amino acid solution. The mean values of the relative molar response (RMR) with respect to pipelicolic acid are calculated from the five analyses. Table 2 presents the mean values and standard deviations of the amounts of amino acids in the seven independent hydrolysates of the same wheat sample. The standard deviations are small and in most cases their relative values are under 5%. The precision of the method can be considered sufficient for practical analytical use. Figure 2 shows the amino acids separated from hydrolysed faecal protein in the form of their N-HFP n-propyl esters.

The amino acid composition of the diet ingredients are shown in Table 3. The lysine content of Eurolysine protein exceeded those of soybean meal and Pekilo by 2–3 g. For pigs, lysine is the essential amino acid that is most limiting in many foods of plant origin, especially cereals. When lysine supplementation is considered for feeds in which lysine is limiting, it is important to obtain an accurate estimate of its availability in these feeds. The

Table 1. Precision of the ion exchange cleanup, derivation and gas-liquid chromatographic analysis performed on a standard amino acid solution.

Amino acid	RMR mean	SD	RSD %
Alanine	0.63	0.02	3.3
Argine	1.31	0.03	2.1
Aspartic acid	1.05	0.02	1.4
Cystine	1.08	0.07	6.5
Glutamic acid	1.10	0.03	2.6
Glycine	0.58	0.02	4.2
Histidine	0.72	0.03	3.8
Isoleucine	0.79	0.04	5.1
Leucine	1.01	0.01	1.3
Lysine	1.15	0.02	1.6
Methionine	0.86	0.02	1.8
Phenylalanine	1.40	0.04	2.5
Proline	0.93	0.02	2.3
Serine	0.91	0.04	4.5
Threonine	0.92	0.05	5.6
Tyrosine	1.48	0.05	3.6
Valine	0.84	0.03	3.7

Table 2. The mean values and standard deviation of the amounts of amino acids in seven independent hydrolysates of the same wheat sample (g/16 g N).

Amino acid	Mean	SD	RSD %
Alanine	3.6	0.14	3.9
Arginine	4.3	0.24	5.6
Aspartic acid	4.9	0.19	3.9
Glutamic acid	28.6	1.07	3.7
Glycine	3.9	0.09	2.3
Histidine	2.1	0.25	11.9
Isoleucine	3.3	0.20	6.1
Leucine	6.6	0.20	3.0
Lysine	2.6	0.14	5.4
Methionine	1.2	0.09	7.5
Phenylalanine	4.9	0.24	4.9
Proline	10.4	0.33	3.2
Serine	4.5	0.13	2.9
Threonine	2.8	0.18	6.4
Tyrosine	3.5	0.29	8.3
Valine	4.2	0.15	3.6

content of threonine, the second limiting amino acid in pig diets, was almost the same as in the other protein sources. Methionine was low in both microbial proteins.

Estimates of the average digestibilities of individual amino acids in different diets are presented in Table 4. The apparent digestibilities of




Figure 2. Amino acid separated from hydrolysed faecal protein in the form of their N-HFB n-propyl esters. The peaks represent approximately 30  $\mu$ g of faecal protein.

arginine, leucine, methionine, phenylalanine and tyrosine were lower with Eurolysine than with soybean meal supplementation the difference being statistically significant ( $P < 0.05$ ). Compared with the Pekilo diet, the Eurolysine diet had lower digestibilities for alanine, aspartic acid, glutamic acid, glycine, leucine, proline and serine ( $P < 0.05$ ).

In the Eurolysine diet lysine digestibility was lower ( $P > 0.05$ ) than in the soybean meal diet but higher than in the Pekilo diet.

Table 3. Amino acid composition of the experimental feeds.

Amino acid g/16 g N	Eurolysine	Pekilo	Soybean meal	Barley
Alanine	7.0	5.6	4.4	4.1
Arginine	3.4	4.6	6.6	4.4
Aspartic acid	7.1	7.1	11.1	5.6
Glutamic acid	9.7	11.5	17.4	20.2
Glycine	3.7	4.2	4.3	4.0
Histidine	0.9	1.2	2.1	2.2
Isoleucine	3.6	3.5	5.0	3.6
Leucine	5.8	6.1	7.7	6.7
Lysine	8.3	5.5	5.9	3.5
Methionine	0.6	0.5	1.0	1.2
Phenylalanine	3.0	3.3	5.0	4.6
Proline	2.6	3.5	5.1	9.4
Serine	3.0	3.7	5.3	4.1
Threonine	3.7	3.7	4.3	3.6
Tyrosine	2.1	2.7	3.5	3.0
Valine	4.5	4.3	5.2	5.0

Table 4. Apparent digestibilities of amino acids in diets containing soybean meal, Eurolysine or Pekilo as protein supplement.

Amino acid	Protein supplement				
	Soybean meal 100 %	Soybean meal 50 % Eurolysine 50 %	Eurolysine 100 %	Eurolysine 50 % Pekilo 50 %	Pekilo 100 %
Alanine	73.6	74.7	66.9	69.9	74.4
Arginine	90.7 <sup>a</sup>	87.4 <sup>b</sup>	81.0 <sup>c</sup>	79.7 <sup>c</sup>	84.6 <sup>b</sup>
Aspartic acid	82.8	76.7	71.1 <sup>b</sup>	72.5 <sup>a</sup>	72.7 <sup>a</sup>
Glutamic acid	89.5	86.7	82.7 <sup>b</sup>	84.4 <sup>ab</sup>	85.7 <sup>a</sup>
Glycine	78.3	76.2	69.4 <sup>b</sup>	73.5 <sup>ab</sup>	76.1 <sup>a</sup>
Histidine	95.5	90.5	90.2	91.8	94.7
Isoleucine	84.8	78.2	76.9	81.0	80.1
Leusine	83.6 <sup>a</sup>	81.0 <sup>ab</sup>	74.9 <sup>b</sup>	78.4 <sup>a</sup>	79.6 <sup>a</sup>
Lysine	82.2	81.6	77.9	74.8	74.8
Methionine	77.8 <sup>a</sup>	69.6 <sup>a</sup>	56.0 <sup>b</sup>	57.9	60.8
Phenylalanine	83.2 <sup>a</sup>	79.8 <sup>b</sup>	74.2 <sup>c</sup>	76.5	77.5
Proline	89.3	88.5	84.8 <sup>b</sup>	85.9 <sup>ab</sup>	87.2 <sup>a</sup>
Serine	85.9	81.9	77.2 <sup>b</sup>	79.2 <sup>ab</sup>	80.7 <sup>a</sup>
Threonine	82.2	76.9	75.3	77.2	77.3
Tyrosine	83.5 <sup>a</sup>	75.1 <sup>b</sup>	70.0 <sup>c</sup>	72.0	73.0
Valine	83.8	78.6	76.3	80.2	80.9

Differences between means with different letters were statistically significant (a, b, c  $P < 0.05$ ).

The digestibilities of various amino acids were lower in diets supplemented with Eurolysine bacterial protein than in diets with soybean meal and Pekilo protein. This is in good agreement with the results for the crude protein digestibilities. In the diets with Eurolysine as protein source the apparent digestibilities were similar to those presented by BRAUDE et al. (1977) for diets containing bacterial protein, but considerably lower than the results of experiments reported by D'MELLO et al. (1976).

The validity of the faecal analysis method for determination of the apparent digestibility of the amino acids is often questioned, partly because secretion of endogenous protein into the digestive tract and partly because of the presence of microorganisms in the hind gut. The microflora of the lower intestine may alter the concentrations of the different amino acids in the faeces (BRAUDE et al. 1977, JUST 1980). However, although possibly modified by microbial activity, the amino acid digestibilities obtained in the present analysis, gave useful information and the results are in good agreement with the data on protein digestibility, nitrogen balance and growth (NÄSI 1982 a).

## References

- BRAUDE, R. HOSKING, Z. D., MITCHELL, K. G., PLONKA, S. & SAMBROOK, I. E. 1977. Pruteen, a new source of protein for growing pigs. I. Metabolic experiment: utilization of nitrogen. *Livest. Prod. Sci.* 4: 79–89.
- D'MELLO, J. P. F., PEERS, D. G. & WHITTEMORE, C. T. 1976. Utilization of dried microbial cells grown on methanol in a semi-purified diet for growing pigs. *Br. J. Nutr.* 36: 403–410.
- JUST, A. 1980. Ileal digestibility of protein: applied aspects. Current concepts of digestion and absorption in pigs. *NIRD-HRI Tech. Bull.* no 3: 66–75. Reading.
- MARCH, J. F. 1975. A modified technic for the quantitative analysis of amino acids by gas chromatography using n-heptafluorobutyric n-propyl ester derivatives. *Anal. Biochem.* 69: 420–442.
- NÄSI, M. 1982 a. Nutritive value of Eurolysine bacterial protein and Pekilo protein for growing pigs. *J. Scient. Agric. Soc. Finl.* 54: 263–269.
- 1982 b. Performance of laying hens on diets containing Eurolysine bacterial protein or Pekilo protein. *J. Scient. Agric. Soc. Finl.* 54: 271–278.

Ms received July 19, 1982

## SELOSTUS

### **Eurolysinebakteerivalkuaista ja pekiloproteiinia sisältävien dieettien aminohappojen sulavuudet siällä**

Matti Näsi

*Helsingin yliopisto, kotieläintieteen laitos 00710 Helsinki 71*

Lea Huida

*Maatalouden tutkimuskeskus, kotieläinhoidon tutkimuslaitos 36100 Jokioinen*

Tutkimuksessa määritettiin aminohappojen näennäiset sulavuudet lihasioilla dieeteillä, joissa soijajauho korvattiin 50 % tai 100 % eurolysine bakteeriproteiinilla tai pekiloproteiinilla. Koe tehtiin kuudella leikolla  $3 \times 3$  latinalaisen neliön koemallin mukaan. Eurolysineä sisältävällä ruokinnalla oli arginiinin, leusiinin, metioniinin, fenyylalaniinin ja tyrosiinin sulavuudet alempia kuin soijaa sisältävällä dieetillä. Samoin alaniinin, asparagiinihapon, glutamiinihapon, glysiinin, leusiinin, proliinin ja seriinin sulavuudet olivat alempia eurolysine- dieetillä verrattuna pekilodieettiin. Lysiinin sulavuus oli eurolysiinidieetillä korkeampi kuin pekilodieetillä.

Tutkimuksessa selvitettiin kaasukromatografisen menetelmän soveltuvuutta rehujen aminohappomäärityksiin.