

Preliminary experiments into the use of *Streptomyces* spp. isolated from peat in the biological control of soil and seed-borne diseases in peat culture

RISTO TAHVONEN

Department of Plant Pathology, University of Helsinki SF-00710 Helsinki 71, Finland

Abstract. *Streptomyces* spp. isolates obtained from peat effectively inhibited the growth of *Alternaria brassicicola* Wiltshire, *Fusarium culmorum* (W. G. Smith) Sacc., *F. sulphureum* Schlechtendahl, *Pythium debaryanum* auct. non Hesse and *Rhizoctonia solani* Kühn on PDA medium, but *Streptomyces* spp. isolates from fine sand soil were not effective against *F. culmorum* compared to the strains isolated from peat. Treatment of cauliflower seeds with *Streptomyces* spp. isolated from peat effectively controlled damping-off caused by *A. brassicicola* and *R. solani* when the seedlings were grown on either disinfected or fresh peat. Spraying the seeding layer of the peat substrate with a suspension of *Streptomyces* reduced the mortality of barley sprouts and foot rot caused by *F. culmorum*, and damping-off on sugar beet caused by *P. debaryanum*.

Introduction

In a number of studies carried out in 1975-1979 on the inhibitory effect of light-coloured *Sphagnum fuscum* peat on plant pathogens (TAHVONEN 1982), it was found that some of the peat lots prevented or reduced the damage caused by a number of soil and seed-borne fungal pathogens. The phenomenon was shown to be due to the action of certain microbes in the peat, of which *Trichoderma viride* Pers. ex Fr. and *Streptomyces* spp. were the most effective against fungal pathogens grown on growth media. In an experiment into the control of damping-off in cabbage, the effect of treatment with *Streptomyces* was greater and of longer duration than that with *T. viride*.

Since, as far as is known, *Streptomyces* species isolated from light-coloured *Sphagnum* peat have not been tested in the biological control of fungal pathogens, further studies were carried out into the possibilities of using *Streptomyces* isolates obtained from peat as a biological control agent. Those pathogens and test plants which could be rapidly and easily used in testing and developing the methods and which, however, were important and/or typical soil and seed-borne fungal pathogens were selected for use in the study.

Material and methods

Streptomyces isolates obtained from peat lots supplied by peat producers and growers were cultivated on PDA medium containing prothiocarp and benomyl fungicides to inhibit the effect of interfering fungi (TAHVONEN 1982). The *Streptomyces* isolates were stored and grown on a medium (pH about 7) containing 4 g yeast extract, 10 g malt extract, 4 g glucose and 20 g agar / 1 000 ml water. In addition to transfers carried out in the normal way with a needle, large amounts of *Streptomyces* spp. were transferred by spraying spore suspensions onto the surface of the medium. This ensured that the surface of the petri dish would be covered with mycelia. The spore suspensions were prepared by scraping off all the mycelia growing in a petri dish ($\varnothing = 9$ cm) and mixing it with 50 ml of sterile water using an Ultra Turrax homogeniser.

The isolates obtained from peat were tested against *Alternaria brassicicola* Wiltshire, *Fusarium culmorum* (W. G. Smith) Sacc., *F. sulphureum* Schlechtendahl, *Pythium debaryanum* auct. non Hesse and *Rhizoctonia solani* Kühn growing on PDA medium (TAHVONEN 1982). *Streptomyces* spp. isolates obtained from fine sand soil, supplied by the Department of Microbiology, University of Helsinki, were tested against *F. culmorum* in order to compare their effectiveness with isolates obtained from peat.

The stock solution of *Streptomyces* needed for the seed and soil treatments was prepared by homogenising the mycelia scraped off the surface of the medium with either pure, autoclaved water or a nutrient solution (4 g yeast extract, 10 g malt extract, and 4 g glucose / 1 000 ml H₂O). Unless otherwise stated, the *Streptomyces* stock solution mentioned in the tables or figures was prepared by mixing Isolate No. 6 with water.

Streptomyces Isolate No. 6 contained 5.6×10^{10} – 1.3×10^{11} spores/ petri dish and Isolate No. 13 contained 4.6×10^9 – 6.0×10^{10} spores/dish. No counts were made on the other isolates. In the experiments shown in Tables 2, 3, 4, 5, 6 and 9, 2 dishes/100 ml were used, and in Fig. 4 and Tables 7 and 8 1 dish/100 ml. In the experiments shown in Fig. 4 and Table 8, the *Streptomyces* spp. homogenised in their own growing solution had been grown in a flat-bottomed flask containing the nutrient solution without agar. There were 1.5×10^9 spores/ml in this stock suspension.

In the seed treatment, the seeds were soaked for about 5 min. in the stock suspension of *Streptomyces* or in aqueous dilutions as shown in the tables, and then dried overnight between sheets of filter paper. In the soil treatment, the seeding layer was sprayed with the stock suspension or with aqueous dilutions as shown in the tables, at a level of about 100 ml/m².

Infection of the seeds was ensured by infecting cauliflower seeds with *A. brassicicola* or *R. solani* and barley seeds with *F. culmorum* or *Helminthosporium sativum* Pammel, King & Bakke. The seeds were immersed in a fungal suspension containing 2-week-old fungal mycelia grown on PDA medium in a petri dish ($\varnothing = 9$ cm) at a level of 1 dish/100 ml. The seeds were dried between filter paper in the laboratory. In the experiment in which soil was infected with *P. debaryanum* and *R. solani*, the fungal mycelium from

one PDA petri dish was mixed with 5 l of peat one week before sowing the sugar beet or cauliflower seeds.

The seeds were grown in plastic boxes (volume about 1 l) at a sowing density of 36 cauliflower seeds or 40 barley and sugar beet seeds/box. 3 or 4 replications of each were used. New peat was used as the growing substrate in all the experiments apart from that presented in Table 6, where steam-disinfected peat was used. The barley seedlings were grown for 3 weeks, the cauliflower seedlings for 3.5 to 4 weeks and the sugar beet seedlings for 6 weeks. Sprouting of the barley seeds was inventoried and the fresh-weight of the plants and degree of infection was determined at the end of the experiment using the scale 0–2, where 0 = healthy, 1 = slightly damaged foot of the stem, 2 = severe foot damage or dead. The number of cabbage and sugar beet seedlings which developed and the number of seedlings suffering from damping-off were determined. However, the results are expressed in the tables as the number of healthy plants/seeds sown so as to include the number of seedlings affected by belowground damping-off. Wherever necessary, the results have been subjected to analysis of variance and the $LSD_{t_{0.05}}$ between the means calculated.

Results

Most of the *Streptomyces* spp. isolates (35) obtained from the peat were effective or very effective in inhibiting the growth of the test fungi on PDA medium (Table 1). On the other hand, the *Streptomyces* spp. strains isolated




Fig. 1. Effect of *Streptomyces* spp. isolates obtained from peat and fine sand soil on the growth of *Fusarium culmorum* on PDA medium. Isolates in the top row made from peat and in the bottom row from fine sand soil. The *F. culmorum* transfer is in the centre of the dishes and the four *Streptomyces* transfers at the edge of the dishes.

Table 1. Inhibitory effect of *Streptomyces* isolates obtained from peat and fine sand soil against different fungi on PDA medium.

| Origin of <i>Streptomyces</i> isolate | Test fungus | Strength of inhibitory effect, number/number of isolates tested | | |
|---------------------------------------|--------------------------------|---|--------|--------|
| | | ineffective | medium | strong |
| Peat | <i>Alternaria brassicicola</i> | 0/19 | 5/19 | 14/19 |
| | <i>Fusarium culmorum</i> | 6/30 | 10/30 | 14/30 |
| | <i>F. sulphureum</i> | 2/24 | 3/24 | 19/24 |
| | <i>Pythium debaryanum</i> | 5/24 | 3/24 | 16/24 |
| | <i>Rhizoctonia solani</i> | 4/31 | 4/31 | 23/31 |
| Fine sand soil | <i>F. culmorum</i> | 16/20 | 4/20 | 0/20 |

Table 2. Effectiveness of different *Streptomyces* isolates in controlling damping-off caused by *Alternaria brassicicola* on cauliflower. Plants grown for 4 weeks.

| <i>Streptomyces</i> isolate No. | Seed treatment | | Soil treatment | |
|---------------------------------|----------------|-------------------|----------------|-------------------|
| | emergence-% | healthy plants, % | emergence-% | healthy plants, % |
| 1 | 75.9 | 63.9 | 46.2 | 23.1 |
| 2 | 85.1 | 75.8 | 51.7 | 32.2 |
| 3 | 58.3 | 54.4 | 32.4 | 15.6 |
| 4 | 87.7 | 82.0 | 39.7 | 15.6 |
| 5 | 72.2 | 61.0 | 52.8 | 20.3 |
| 6 | 81.4 | 71.1 | 52.8 | 35.0 |
| 7 | 36.1 | 19.4 | 37.8 | 15.6 |
| 8 | 65.6 | 52.8 | 40.6 | 23.1 |
| 9 | 61.1 | 48.9 | 30.6 | 19.4 |
| 10 | 72.2 | 55.5 | 56.4 | 32.2 |
| 11 | 55.6 | 45.3 | 35.0 | 22.2 |
| 12 | 75.0 | 61.9 | 59.2 | 32.2 |
| control | 14.7 | 5.6 | 14.7 | 5.6 |
| healthy seeds | 89.8 | 89.8 | 89.8 | 89.8 |

from mineral soil either did not inhibit at all or else only moderately inhibited the growth of *Fusarium culmorum* (W. G. Smith) Sacc. (Fig. 1 and Table 1).

In the seedling growing experiment, a number of *Streptomyces* spp. isolates, e.g. 1, 2, 4, 6, 10 and 12, effectively controlled damping-off caused by *Alternaria brassicicola* Wiltshire on cauliflower (Table 2). When the seed treatment was used, the control result was almost the same as for healthy seeds. However, the results obtained with the soil treatment were always significantly inferior to those for the seed treatment. Treatment of barley seed with *Streptomyces* spp. decreased the amount of damage caused by *F. culmorum* and *Helminthosporium sativum* Pamel, King & Bakke and increased the growth in fresh weight. Isolate No 2 reduced the incidence of damage and increased the growth in fresh weight of non-infected seeds where pathogens were found to occur naturally (Table 3). When the *Streptomyces* spp. treatment was given by spraying the seeding layer, the result for

Table 3. Effect of different concentrations of *Streptomyces* sp. isolates on the germination, disease index and fresh growth after 3 weeks of non-infected barley seed and barley seed infected with *Fusarium culmorum*, *Helminthosporium sativum* suspensions.

| <i>Streptomyces</i> seed treatment | Dilution made from stock <i>Streptomyces</i> suspension | | | | \bar{X} |
|---|---|--------------|--------------|--------------|--------------|
| | 1/1 | 1/2 | 1/5 | 1/10 | |
| sprouting-% / infection degree index, 0-2 / fresh yield, g/dish | | | | | |
| Seeds infected with <i>Fusarium culmorum</i> | | | | | |
| Isolate 1 | 72/0.62/28.3 | 68/0.69/20.8 | 66/0.73/28.5 | 76/0.60/31.7 | 70/0.66/27.3 |
| Isolate 2 | 62/0.45/23.8 | 40/0.76/16.2 | 50/0.48/24.1 | 60/0.60/26.0 | 54/0.57/22.5 |
| Isolate 3 | 58/0.98/14.9 | 66/0.99/15.9 | 56/0.78/16.2 | 48/0.87/16.2 | 58/0.91/15.8 |
| control | 52/1.07/12.4 | 52/1.07/12.4 | 52/1.07/12.4 | 52/1.07/12.4 | 52/1.07/12.4 |
| \bar{X} (isolate 1-3) | 64/0.68/22.3 | 58/0.81/17.6 | 57/0.66/22.9 | 61/0.69/24.6 | 61/0.71/21.9 |
| Seeds infected with <i>Helminthosporium sativum</i> | | | | | |
| Isolate 1 | 94/0.56/39.4 | 92/0.63/39.6 | 92/0.74/39.9 | 92/0.82/34.8 | 92/0.69/38.4 |
| Isolate 2 | 82/0.30/22.8 | 84/0.61/23.9 | 94/0.19/33.2 | 88/0.49/39.2 | 88/0.40/29.8 |
| Isolate 3 | 92/0.67/39.2 | 90/0.79/30.7 | 90/0.68/32.6 | 88/0.90/19.3 | 90/0.76/30.5 |
| control | 90/0.72/30.0 | 90/0.72/30.0 | 90/0.72/30.0 | 90/0.72/30.0 | 90/0.72/30.0 |
| \bar{X} (isolate 1-3) | 89/0.51/33.8 | 89/0.68/31.4 | 92/0.54/35.2 | 89/0.74/31.1 | 90/0.62/32.9 |
| Non-infected seeds | | | | | |
| Isolate 1 | 94/0.35/37.0 | 90/0.28/37.2 | 90/0.38/42.3 | 96/0.44/37.1 | 90/0.36/38.4 |
| Isolate 2 | 94/0.12/27.6 | 98/0.07/27.3 | 92/0.09/42.6 | 92/0.17/40.4 | 94/0.11/34.5 |
| Isolate 3 | 94/0.33/41.1 | 90/0.53/40.5 | 90/0.40/33.3 | 92/0.48/27.6 | 92/0.44/35.6 |
| control | 92/0.28/23.0 | 94/0.28/23.0 | 94/0.28/23.0 | 94/0.28/23.0 | 94/0.28/23.0 |
| \bar{X} (isolate 1-3) | 94/0.27/35.2 | 93/0.27/35.0 | 91/0.27/37.2 | 93/0.27/35.0 | 93/0.30/36.2 |

Table 4. Effect of different *Streptomyces* sp. isolates on the germination, degree of infection and fresh growth after 3 weeks of barley seeds infected with *Fusarium culmorum*. Soil and seed treatments carried out with different *Streptomyces* dilutions.

| Seed treatment | Dilution made from stock <i>Streptomyces</i> suspension | | | | \bar{X} |
|--|---|--------------|--------------|--------------|--------------|
| | 1/1 | 1/2 | 1/5 | 1/10 | |
| sprouting-% / infection degree index, 0-2 / fresh yield g/dish | | | | | |
| Isolate 1 | 60/1.09/8.6 | 60/0.96/8.3 | 73/0.88/10.2 | 78/0.96/14.5 | 68/0.97/10.4 |
| Isolate 2 | 78/0.65/14.1 | 76/0.70/12.7 | 63/0.81/6.5 | 75/0.83/11.2 | 73/0.75/11.1 |
| Isolate 3 | 71/0.92/8.5 | 40/1.06/3.4 | 73/0.81/11.6 | 77/0.75/13.5 | 65/0.89/9.3 |
| \bar{X} | 70/0.89/10.4 | 59/0.91/8.1 | 70/0.83/9.4 | 77/0.85/13.1 | 69/0.87/10.3 |
| Soil treatment | | | | | |
| Isolate 1 | 83/0.66/12.8 | 80/0.79/18.7 | 81/0.69/11.2 | 84/0.74/18.3 | 82/0.72/15.3 |
| Isolate 2 | 83/0.66/12.3 | 83/0.67/14.6 | 83/0.64/16.6 | 84/0.77/21.9 | 83/0.69/16.4 |
| Isolate 3 | 77/0.69/15.2 | 73/0.81/13.5 | 82/0.87/12.4 | 81/0.75/15.4 | 78/0.78/14.1 |
| \bar{X} | 81/0.67/13.4 | 79/0.76/15.6 | 82/0.73/13.4 | 83/0.75/18.5 | 81/0.73/15.3 |
| Control | 46/1.16/3.7 | | | | |
| Non-infected seeds | 89/0.29/18.7 | | | | |





Fig. 2. Effect of spraying the seeding layer with *Streptomyces* sp. on the damage caused by *Fusarium culmorum* on barley. Bottom dish = untreated seeds infected with *F. culmorum*, upper right-hand dish = untreated, uninfected seeds; upper left-hand dish = seeds infected with *F. culmorum* and seeding layer sprayed with *Streptomyces* suspension.

protection against *F. culmorum* improved significantly in comparison to the results for the seed treatment. Sprouting was at the same level as the uninfected seeds when the soil treatment was carried out, and the fresh weight of the shoots was of the same order of magnitude as for the non-infected barley and four times greater than that of the controls (Table 4). Diluting the stock solution to 10 % had no effect on the control results (Tables 3 and 4).

When the stock suspension of *Streptomyces* used for treating the cauliflower seeds against *A. brassicicola* and for the soil treatment against *F. culmorum* on barley, the nutrients used in the growth medium being added to the suspension, was tested, it was found that the stock solution could be diluted to at least 10^{-6} without any significant weakening of the control results (Tables 5 and 6). When dilutions were made with water, the control results deteriorated at dilutions below 10^{-2} . Mixing different *Streptomyces* isolates in the same stock solution had no effect on the control result.

When the *Streptomyces* sp. to be used as the stock suspension was grown in the nutrient solution and then homogenised in the original growing solution, and dilutions made with the suspension, it gave as good or better results in controlling seed-borne damping-off caused by *A. brassicicola* and *Rhizoctonia solani* Kühn on cauliflower, and in the control of *F. culmorum* on barley, than when the stock suspension was homogenised in a new nutrient medium (Fig 3 and Table 7).

HEALTHY SEEDLINGS



Δ=CONTROL, ○=STREPTOMYCES SP. HOMOGENIZED INTO ITS OWN GROWTH SOLUTION, ×=STREPTOMYCES SP. HOMOGENIZED INTO NEW GROWTH SOLUTION
 1= STREPTOMYCES SP. STOCK SUSPENSION, -1=10⁻¹ AND -3=10⁻³ DILUTION

Fig. 3. Effect of *Streptomyces* sp. mixed with the nutrient solution and with the original growing solution given as different concentrations on seed-borne damping-off caused by *Alternaria brassicicola* and *Rhizoctonia solani* on cauliflower.

Table 5. Effect of nutrient solution and dilutions on the effectiveness of *Streptomyces* sp. treatment of seeds in controlling damping-off caused by *Alternaria brassicicola*.

| | <i>Streptomyces</i> isolate No | Dilution made from stock <i>Streptomyces</i> suspension | | | | |
|--|--------------------------------|---|------------------|------------------|------------------|------------------|
| | | 1 | 10 ⁻¹ | 10 ⁻² | 10 ⁻⁴ | 10 ⁻⁶ |
| emergence-% / healthy plants, % | | | | | | |
| Aqueous <i>Streptomyces</i> suspension | 6 | 92.6/91.7 | 84.3/82.4 | 82.4/79.6 | 57.4/40.7 | 54.6/34.3 |
| | 13 | 91.7/89.8 | 89.8/85.2 | 59.3/49.1 | 48.1/37.0 | 40.7/26.9 |
| | 6+13 | 92.6/90.7 | 86.1/78.7 | 46.3/35.2 | 60.2/49.1 | 39.8/21.3 |
| | \bar{X} | 92.3/90.7 | 86.7/82.1 | 62.7/54.6 | 55.2/42.3 | 45.0/27.5 |
| Nutrient <i>Streptomyces</i> suspension | 6 | 91.7/87.0 | 89.8/89.8 | 75.9/65.7 | 80.6/65.7 | 76.9/69.4 |
| | 13 | 90.7/90.7 | 79.6/77.8 | 74.1/70.4 | 62.0/50.0 | 87.0/79.6 |
| | 6+13 | 90.7/88.0 | 85.2/80.6 | 64.8/61.1 | 80.6/70.4 | 82.4/77.8 |
| | \bar{X} | 91.0/88.6 | 84.9/82.7 | 71.6/65.7 | 74.4/62.0 | 82.1/75.6 |
| | control | | 44.4/29.6 | | | |

Table 6. Effect of nutrients added to the stock *Streptomyces* suspension and dilutions when carrying out soil treatment on the incidence of seed-borne foot rot disease caused by *Fusarium culmorum* in barley grown on steam-disinfected peat substrate.

| | Isolate | Dilution made from stock suspension | | | | | \bar{X} |
|---------------------------------------|-----------|-------------------------------------|-----------|-----------|-----------|-----------|-----------|
| | | 1 | 10^{-1} | 10^{-2} | 10^{-4} | 10^{-6} | |
| % healthy plants 3 weeks after sowing | | | | | | | |
| Aqueous stock suspension | 6 | 72.2 | 63.9 | 67.6 | 53.7 | 51.9 | 61.9 |
| | 13 | 79.6 | 65.7 | 68.5 | 52.8 | 60.2 | 65.4 |
| | 6+13 | 74.1 | 77.7 | 67.6 | 59.3 | 64.8 | 68.7 |
| | \bar{X} | 75.3 | 69.1 | 67.9 | 55.3 | 58.9 | 65.3 |
| Nutrient stock suspension | 6 | 82.4 | 74.1 | 74.1 | 75.0 | 57.4 | 72.6 |
| | 13 | 73.1 | 79.6 | 78.7 | 71.3 | 75.0 | 75.5 |
| | 6+13 | 72.2 | 73.2 | 69.4 | 70.4 | 70.4 | 71.1 |
| | \bar{X} | 75.9 | 75.6 | 74.1 | 72.2 | 67.6 | 73.1 |
| Control | \bar{X} | 53.7 | | | | | |
| Healthy seeds | \bar{X} | 78.7 | | | | | |

Table 7. Effect of *Streptomyces* suspensions homogenised in the nutrient solution and in the original growing solution when carrying out soil treatment on the incidence of seed-borne foot rot caused by *Fusarium culmorum* on barley.

| Stock <i>Streptomyces</i> solution | Dilution made from stock suspension | | | |
|--|-------------------------------------|-----------|-----------|---------|
| | 1 | 10^{-1} | 10^{-3} | control |
| Homogenised in fresh nutrient solution | | | | |
| fresh yield, g/dish | 18.7 | 18.8 | 16.6 | 11.9 |
| degree of infection, 0-2 | 0.53 | 0.67 | 0.53 | 0.67 |
| Homogenised in original growing solution | | | | |
| fresh yield, g/dish | 24.4 | 21.9 | 14.1 | 11.9 |
| degree of infection, 0-2 | 0.15 | 0.32 | 0.62 | 0.67 |

LSD_{10,05}: fresh yield = 5.2 g, degree of infection = 0.28

Table 8. Effect of the *Streptomyces* sp. treatment on the incidence of soil-borne damping-off caused by *Rhizoctonia solani* on cauliflower growing on peat substrate.

| Stock <i>Streptomyces</i> suspension | Seed treatment | Seed treatment + soil treatment, different dilutions | | | | |
|--------------------------------------|----------------|--|-----------|-----------|-----------|---------|
| | | 1 | 10^{-1} | 10^{-2} | 10^{-3} | control |
| Aqueous suspension | | | | | | |
| emergence-% | 75.0 | 77.1 | 79.9 | 76.4 | 77.1 | 77.1 |
| healthy plants, % | 70.8 | 47.2 | 61.1 | 15.3 | 64.6 | 31.3 |
| Nutrient suspension | | | | | | |
| emergence-% | 78.5 | 78.5 | 84.0 | 83.3 | 84.7 | 77.1 |
| healthy plants, % | 63.9 | 51.4 | 42.4 | 66.0 | 56.9 | 31.3 |

F values for healthy plants: aqueous suspension = 5.35**
nutrient suspension = 0.70.




Fig. 4. Effect of *Streptomyces* seed treatment on soil-borne damping-off caused by *Rhizoctonia solani* on cauliflower. Upper row = seeds treated with *Streptomyces*, bottom row = untreated seeds.



Fig. 5. Effect of treating the substrate with *Streptomyces* on damping-off caused by *Pythium debaryanum* on sugar beet. Upper row = sown layer treated with *Streptomyces*, bottom row = no treatment.

Treating the seeds with *Streptomyces* effectively controlled soil-borne damping-off caused by *R. solani* on cauliflower (Table 8). Combined seed and soil treatment with an aqueous solution of *Streptomyces* increased the incidence of damping-off in comparison to the seed treatment alone. Damping-off reached peak values when the soil was sprayed with a 10^{-2} dilution after sowing. The result was better when both greater and smaller concentrations were used. The same trend was evident with the nutrient suspension treatment, but the result was not statistically significant.

Spraying the seeding layer with *Streptomyces* significantly decreased the amount of soil-borne damping-off caused by *Pythium debaryanum* auct. non Hesse on sugar beet (Table 9).

Table 9. Effect of *Streptomyces* spp. isolates given as seed and soil treatment at different dilutions on soil-borne damping-off caused by *Pythium debaryanum* on sugar beet. Peat infected with *Pythium* one week before sowing.

| | Dilution made from stock <i>Streptomyces</i> suspension | | | | \bar{X} |
|----------------|---|------|--------------------|------|-----------|
| | 1/1 | 1/5 | 1/10 | 1/20 | |
| | % of healthy seeds 6 weeks after sowing | | | | |
| Soil treatment | | | | | |
| Isolate 2 | 36.2 | 40.0 | 39.0 | 42.8 | 39.5 |
| Isolate 6 | 47.6 | 53.3 | 42.9 | 43.8 | 46.9 |
| | \bar{X} | 41.9 | 46.7 | 41.0 | 43.3 |
| Seed treatment | | | | | |
| Isolate 2 | 21.0 | 29.5 | 13.4 | 19.1 | 20.6 |
| Isolate 6 | 30.5 | 19.1 | 16.2 | 7.6 | 18.4 |
| | \bar{X} | 25.8 | 24.3 | 14.8 | 13.4 |
| Control | \bar{X} | 24.8 | (emergence 61.1 %) | | 19.6 |

F values: treatment = 64.6**, isolates = 0.7,
dilutions = 1.7, combined effect 2.8

Discussion

In the studies carried out by BROADBENT et al. (1971), 10–47 % of the *Streptomyces* isolates were antagonistic to fungal pathogens, and in the studies carried out by COOPER and SHILTON (1949) 30 %. Between 75 and 80 % of the isolates made from peat were antagonistic while all the isolates made from fine sand soil were ineffective. This would indicate that the *Streptomyces* species isolated from peat are strong antagonists in comparison to the *Streptomyces* strains isolated from other types of soil. This may explain, in part, the strong inhibitory effect shown by certain peat lots against plant pathogens since only some of the peat lots were found to have large *Streptomyces* populations (TAHVONEN 1982). Since the *Streptomyces* isolates

tested here have not yet been identified in full, it is difficult to compare them with the results obtained by KNAUSS (1976). However, the antagonistic properties of the *Streptomyces* isolates obtained from peat are obviously at least as strong as the *Streptomyces* isolates obtained in many other studies (COOPER and SHILTON 1949, JOHNSON 1954, RANGASWAMI and ETHIRAJ 1962, BROADBENT et al. 1971, KNAUSS 1976, TURHAN 1981 a).

Peat appears to be a natural organic substrate for *Streptomyces* spp. where they are able to reproduce and spread, where the production of antibiotics is high and where the damage caused by fungal pathogens is restricted and inhibited effectively. The roots of growing plants apparently further stimulate the activity of *Streptomyces* spp. (COOPER and SHILTON 1949, RANGASWAMI and VEDYASEKAREN 1963), which certainly has a beneficial effect on the control of pathogens. Indications of plant roots having a stimulating effect have also been found with peat (TAHVONEN 1982).

The success of the biological control of *Alternaria*, *Fusarium*, *Helminthosporium* and *Pythium* fungi using *Streptomyces* isolates from peat appears to continue throughout the seedling stage at least, i.e. for 3–6 weeks. This is further evidence to show that this microbe thrives well in peat. The control result in new, non-disinfected peat is also good, the natural microflora still being present in the peat. In general, this sort of biological control result has been obtained from disinfected or partly disinfected substrates (JOHNSON 1954, LOCKWOOD 1958, BROADBENT et al. 1971, SCHER and BAKER 1980). However, TURHAN (1981 b) has obtained good results with *Streptomyces* treatment in the control of soil-borne pathogens in non-disinfected substrates. In the study carried out by BROADBENT et al. (1971), it was found that the pathogen-inhibiting effect of soil containing a number of different antagonistic species was not as high as soil where the number of antagonistic fungi was lower. Peat presumably contains only two significant antagonists, *Trichoderma viride* Pers. ex Fr. and *Streptomyces* spp. (TAHVONEN 1982). This may explain the good results obtained in biological control with *Streptomyces* spp. treatment in non-disinfected soil, too.

The experiments carried out here show rather reliably that the most common seed and soil-borne fungi causing damping-off can be controlled biologically in peat substrates by treating either the seeds or the soil with *Streptomyces* isolates obtained from peat. The control result in most of the experiments came close to the level of the healthy control plants. Even the soil-borne fungi, *Rhizoctonia solani* Kühn and *Pythium debaryanum* auct. non Hesse, could be controlled or restricted. This is usually extremely difficult or impossible to do by ordinary control measures.

Biological control methods for fungal pathogens in peat cultures require extensive studies in order to identify the best isolates and application techniques for the control of different pathogens, especially in long-term cultures. The number of factors involved in the preservation and growth of *Streptomyces* spp. isolates presupposes further extensive studies. A good example of the complexity of these phenomena is the experiment into the control of soil-borne *R. solani*, where seed treatment alone gave the best result and when seed treatment was combined with a soil surface treatment in

which different dilution levels of the stock solution were used, the effectiveness of the biological control of damping-off was lost completely in the case of certain dilutions.

Acknowledgements: Pirkko Harju, M.Sc. and Kaija Karhunen, M.Sc. participated in carrying out the practical aspects of the experiments. Lahja Pesonen, Pentti Heinänen and Tauno Koivunen, staff members of the Department of Plant Pathology, have all helped in establishing and carrying out the experiments. Kemira Oy have assisted by paying the salaries of some of the assistants and providing materials. I would like to extend my sincere gratitude to all of the above mentioned persons.

References

- BROADBENT, P., BAKER, K. F. & WATERWORTH, Y. 1971. Bacteria and Actinomycetes antagonistic to fungal root pathogens in Australian soils. *Aust. J. Biol. Sci.* 24: 925-944.
- COOPER, W. E. & SHILTON, S. J. P. 1949. Antibiosis of Actinomycetes strains to *Pythium arrhenomanes*, and *Rhizoctonia solani*. *Phytopath.* 39: 5.
- JOHNSON, L. F. 1954. Antibiosis in relation to *Pythium* root rot of sugarcane and corn. *Phytopath.* 44: 69-73.
- KNAUSS, J. F. 1976. In vitro antagonistic activity of several *Streptomyces* spp. against species of *Pythium* and *Phytophthora*. *Pl. Dis. Rep.* 60: 846-850.
- LOCKWOOD, J. L. 1958. *Streptomyces* spp. as a cause of natural fungitoxicity. *Phytopath.* 48: 395.
- RANGASWAMI, G. & ETHIRAJ, S. 1962. Antibiotic production by *Streptomyces* sp. in unamended soil. *Phytopath.* 52: 989-992.
- & VEDYASEKAREN, P. 1963. Antibiotic production by *Streptomyces* spp. in corn rhizosphere. *Phytopath.* 53: 995-997.
- SCHER, F. M. & BAKER, R. 1980. Mechanism of biological control in a *Fusarium*-suppressive soil. *Phytopath.* 70: 412-417.
- TAHVONEN, R. 1982. The suppressiveness of Finnish light coloured *Sphagnum* peat. *J. Scient. Agric. Soc. Finl.* 54: 357-369.
- TURHAN, G. 1981 a. A new race of *Streptomyces ochraceiscleroticus* in the biological control of some soil-borne plant pathogens I. Effects of the isolate C/2-9 on some of the most important soil-borne fungi in vitro. *Z. Pfl. krankheiten und Pfl. schutz* 88: 373-381.
- 1981 b. A new race of *Streptomyces ochraceiscleroticus* in the biological control of some soil-borne plant pathogens II. In vivo studies on the possibilities of using C/2-9 against some important diseases. *Z. Pfl.krankheiten und Pfl. schutz* 88: 422-434.

Esitutkimuksia turpeesta eristettyjen *Streptomyces* spp. -isolaattien käytöstä maa- ja siemenlevintäisten tautien biologisessa torjunnassa turveviljelyssä

Risto Tahvonen

Helsingin yliopiston kasvipatologian laitos, 00710 Helsinki 71

Vuosina 1975–79 tehdyissä selvityksissä vaalean rahkaturpeen sienitautien estovaikutuksista oli todettu, että turpeesta eristetyt *Streptomyces* spp. -isolaatit, joita seuraavassa kutsutaan sädesieniksi, estivät tehokkaasti sienten kasvun ravintoalustalla ja pystyivät vähentämään merkittävästi taimipoltetta kukkakaalilla. Vuosina 1979–81 tehtiin Helsingin yliopiston kasvipatologian laitoksella alustavia tutkimuksia sädesienien käyttömahdollisuuksista biologisessa kasvitautien torjunnassa turvealustalla.

Sädesieni-isolaatteja testattiin ravintoalustatestein *Alternaria brassicicola* Wiltshire, *Fusarium culmorum* (W. G. Smith) Sacc., *Pythium debaryanum* auct. non Hesse ja *Rhizoctonia solani* Kühn -sieniä vastaan. Kasvatuskokeissa olivat testikasveina kukkakaali, ohra ja sokerijuurikas. Kukkakaalin siemenet saastutettiin *A. brassicicola* ja *R. solani* -sienellä tai kasvialustana käytetty turve saastutettiin viikko ennen kylvöä *R. solani* -sienellä. Ohralla käytettiin *F. culmorum* ja *Helminthosporium sativum* Pammel, King & Bakke -sienellä keinosaastrutettua siementä ja sokerijuurikkaalla kasvialusta oli saastutettu viikko ennen kylvöä *P. debaryanum* -sienellä. Siemeniä ja maata käsiteltiin sädesienestä tehdyillä vesi- ja ravinnesuspensioilla sekä näiden vesilaimennoksilla. Sädesienien taudintorjuntakykyä mitattiin taimettumisella ja taimipolteen määrällä (kukkakaali ja sokerijuurikas) sekä orastumisella, tyvitaudin runsaudella ja tuorepainolla (ohra).

Useimmat turpeesta eristetyistä 35 sädesieni-isolaatista estivät ravintoalustalla testisienien kasvua, mutta hietamaasta eritetyt 20 sädesieni-isolaattia, jotka oli saatu Mikrobiologian laitokselta Viikistä, eivät olleet lainkaan tehokkaita testattua *F. culmorum* -sientä vastaan.

Kukkakaalin siementen peittäminen sädesienisuspensioilla esti tai vähensi merkittävästi *A. brassicicola* ja *R. solani* -sienen aiheuttamaa taimipoltetta. Maan käsittely sädesienisuspensioilla oli tehokkaampi kuin siemenkäsittely ohran *F. culmorum* -tyvitaudin torjunnassa. Saastuttamattomien ohran siementen sädesienikäsittely lisäsi oraiden tuorepainoa verrattuna käsittelemättömiin. Kylvökerroksen sädesienikäsittely vähensi merkittävästi *P. debaryanum* -sienen aiheuttamaa taimipoltetta sokerijuurikkaalla.

Nyt saadut tulokset antavat viitteitä siitä, että turveviljelyssä on mahdollista hallituissa kasvihuoneolosuhteissa torjua tai ainakin rajoittaa sienitautien tuhoja biologisesti. Biologisen torjunnan käyttömuotoina olisivat siementen peittäuskäsittelyt ja mahdollisesti kylvökerroksen käsittelyt desinfioidun tai uuden turpeen sienitautien estovaikutuksen palauttamiseksi tai kohottamiseksi. Käytännön sovellutukset vaativat kuitenkin vielä laajoja jatkotutkimuksia tärkeimmillä kasvihuoneissa viljeltävillä kasveilla.