

Cooked meat products made of coarsely ground pork: the main bacterial strains of bacterial flora, their heat resistance and effect on spoilage

ESKO PETÄJÄ and EERO PUOLANNE

PETÄJÄ, E. & PUOLANNE, E. 1993. **Cooked meat products made of coarsely ground pork: the main bacterial strains of bacterial flora, their heat resistance and effect on spoilage.** Agric. Sci. Finl. 2: 413-421. (Dept. Food Technol., Meat Sect., FIN-00014 University of Helsinki, Finland.)

This study was conducted to investigate the bacterial flora of the surface layer and the core of meat products made of coarsely ground pork at the moment of spoilage when stored at 7°C or 4°C. The dominating strains were isolated, their heat resistance was studied in APT-broth, on APT-agar and in coarsely ground cured pork, and their growth after heating and effect on spoilage were followed in coarsely ground cured pork.

The first signs of spoilage appeared in the surface layer of the products. The strains were coccoid lactic acid bacteria with counts ranging from 3.5 to 7.8 log cfu (colony forming units)/g. They survived only accidentally after heating for 15 minutes at 72°C in APT-broth. The core of the products contained only coccoid lactic acid bacteria or only pseudomonads or both as the main bacterial strains. The counts ranged from 2.6 to 6.0 log cfu/g. Most of the strains isolated from the core survived after heating for 30 minutes at 72°C in APT-broth in at least three tests out of six. The most noticeable result of the study was the occurrence of heat-resistant pseudomonads in the core. It must be pointed out that all pseudomonads found survived after heating for 60 minutes at 72°C in APT-broth, and often after heating for 15 minutes at 72°C in coarsely ground cured pork (core 72°C). The cfu number of the two most heat-resistant streptococcus strains decreased only 1 log unit over 15 minutes at 72°C in coarsely ground cured pork. The numbers of inoculated pseudomonads decreased but those of streptococci rose by a maximum of 1 log unit when the experimental porks were kept at 4°C after heating. This indicates that streptococci and pseudomonads probably do not constitute a serious spoilage factor in cooked meat products, but spoilage is generally effected by bacteria which have contaminated the surface layer of the products after heat treatment.

Key words: keepability, spoilage, cooked meat, heat resistance

Introduction

The manufacture and consumption of food products made of coarsely ground meat has grown over the last 10 years in Finland. These products sometimes spoil quite quickly so that not only the surface layer but also the core is spoiled. There is not much

information available on the keeping times or the microbial flora of these kinds of products, or about whether the bacteria which can survive heat treatment in the core spoil the product. These products can be assumed to resemble bologna-type cooked sausages and luncheon meat in many respects. In these products high bacterial counts may be found

although the products would not always have spoiled. PARADIS and STILES (1978) have found 60% of commercial bologna type sausages contain bacteria over 10^6 cfu/g. According to ALLEN and FOSTER (1960) and HILL et al. (1976), 4-week-old bolognas contain over 10^7 bacteria/g. DUISCHAEVER (1977) reported over 10^6 bacteria/g in 50% of commercial luncheon meats and HILL et al. (1976) found that 71% of luncheon meats stored for 3 weeks at 4 - 7°C contained over 10^6 bacteria/g. According to many scientists (REUTER 1969, MANTTEL and BECK 1977, DAELMAN and HOOF 1975) lactobacilli are the biggest bacterial group in cooked meat products. However, DOWDELL and BOARD (1968) and GARDNER (1966, 1968) report that *Microbacterium thermosphactum* is the most important bacterial group causing spoilage in cooked sausage. GARDNER (1966) also found that cooked sausage may contain gram-negative bacteria, pseudomonads and achromobacteria, although gram-negative bacteria are generally assumed to die during the processing of cooked meat products (HALLENBACH and POTTER (1981) and PALUMBO et al. (1974).

The purpose of this study was to find out the main bacterial types in the surface layer and core of cooked meat products made of coarsely ground pork, their survival after heat treatment in APT-broth, on APT-agar, and in coarsely ground cured pork, as well as their growth after heating and their effect on spoilage during storage in coarsely ground cured pork. Various methods were used since heat resistance is known to vary in different media.

Material and methods

Product samples

Seven samples of coarsely ground pork products were received directly from the manufacturers. The samples were 2-4 weeks old and they were kept at 4°C and 7°C up to the moment when the first signs of spoilage, smell and/or slime, were detected. Different samples were investigated for the two temperatures. At the moment when the first signs of

spoilage appeared the samples were examined microbiologically.

Microbiological examination of product samples

The product was peeled and the surface layer sample was taken aseptically at 1-2 mm in thickness. The core sample was taken from the centre of product after cutting the product in two parts so that nothing came into contact with the sampling area. The following examinations were made:

- Total count of bacteria on plate count agar (Difco 0479, 2-4 d at 30°C), on APT-agar (Merck 10453, 3 d at 30°C) and on blood agar (Orion, Espoo, Finland, 2 d at 37°C). Because the bacteria to be counted may not grow in all media, three different agars for aerobic total plate count were used to ensure the growth of the dominating strains.
- Total count of anaerobic bacteria on SPS-agar (Merck 10235, 2 d at 37°C in anaerobic jar).
- Number of lactic acid bacteria on Rogosa-agar (Merck 5463, 4 d at 30°C).
- Number of *Brochothrix thermosphacta* on STAA-agar (GARDNER 1966, 2 d at 22°C).
- Number of staphylococci and micrococci on Baird-Parker-agar (Lab 85 and X085, 2 d at 37°C).
- Number of coliforms on VRB-agar (Lab 31, 1 d at 37°C).

The results were tested by variance analysis (Epistat program) in relation to storage temperature and surface layer and core.

Isolation of the main bacterial types

The surface layer and core of all product samples contained only a few strains in high numbers, and these were isolated. The main colonies were selected from the APT-agar dishes on which the samples had been plated. The dishes with the highest dilution were used. The colonies to be isolated were transferred to APT-agar and incubated for 2

days at 30°C. One typical colony of each strain was removed to APT-agar for storing the culture.

Survival of the isolated strains after heating on solid and in liquid medium

Solid medium

The survival of the strains after heating on a solid medium was studied with a Gradiplate W10-temperature-gradient incubator (Biodata Oy, Helsinki, Finland) with which it is possible to achieve a stepless temperature gradient within a range of 0-10°C. The operating principle is as follows: The bacteria are grown in cuvettes of 65*98*12 mm, into which the agar is poured. The limit temperature for growth or survival is determined by measuring the distance of the growth limit from the lower temperature sensor and by calculating the respective temperature. The survival of the strains was investigated by growing them on APT-agar (Merck 10453) for 30 minutes at a temperature range of 50 - 70°C. The inoculation was carried out by rolling a drop (0.05 ml) of APT-broth culture onto the surface of the agar. Two strains were inoculated in the same cuvette. The 18-hour-old APT-broth culture was used for inoculation. Table 2 shows the ranges of inoculated cell count. Three determinations were made for each strain.

Liquid medium

The survival of the strains after heating in APT-broth at 72°C was investigated side by side with the experiments on solid medium. The 5 ml APT broth was heated in a test tube to 72°C and inoculated with 0.05 ml of 18-hour-old APT-broth culture. The ranges of the inoculated cell counts are presented in Table 2. Different tubes were prepared for each heating period and temperature. The inoculated test tubes were heated for 15, 30, 60, 90 and 120 minutes. After heat treatment they were incubated at 30°C for 2 days and the growth (turbid broth) was recorded. Six experiments were made for each strain.

Survival of some isolated *Pseudomonas* and *Streptococcus* strains after heating in coarsely ground cured pork

The survival of four *Pseudomonas* strains (1b₂, 4b, 6b, 7b₂) and two *Streptococcus* strains (3b₁, 7b₁) was also examined in coarsely ground cured pork after heating at 72°C. One experimental batch contained 150 g pork, 0.67 g glucose and 15 g water. The following additives were used: NaCl (2%), NaNO₂ (0.012%), phosphates (0.15% P₂O₅) and Na-ascorbate (0.04%). The bacterial inoculum was added as APT-broth culture (15 ml). The target number of colony forming units (cfu) in the inoculum was 10⁷/g meat. Control batches were made without bacterial inoculum. Instead, 15 ml of water was added to the control batch. The coarse grinding of the meat and mixing of additives and bacteria were done in a Moulinex mixer (Moulinex, France) to a particle size of 0.5-1 cm (diameter).

Both experimental and control meat batches were packed in the 10 ml glass tubes in which they were heated. Different tubes were prepared for each heating period. The pseudomonas tubes were heated in a water bath for 5, 15 and 30 minutes after the temperature had been raised to 72°C. The respective heating periods for *Streptococcus* 3b₁ tubes were 0.5, 2, 5, and 15 minutes and for *Streptococcus* 7b₁ tubes 0.5, 2, 5, 15, 30 and 60 minutes. After heating the tubes were cooled for 10 minutes at 10°C in a water bath and for 1 hour at 22°C. The following microbiological determinations were made for the non-heated and heated coarsely ground meats, as well as for the meats heated for 5 minutes and stored for 2-3 weeks at 6°C: total plate count on APT-agar, the number of pseudomonads on GSP-agar, staphylococci and micrococci on Baird-Parker-agar, lactic acid bacteria on Rogosa-agar, and *Brochothrix thermosphacta* on STAA-agar. The manufacturers of the media as well as the incubations were the same as those in the microbiological studies of the meat product samples.

The number of colony forming units of the inoculums was determined on APT-agar by incubating for 2 days at 30°C. The survival of the inoculums after heating in APT-broth at 72°C (0.05 ml in 5 ml APT-broth) was also tested side by side with

Table 1. Bacterial flora of cooked meat products made of coarsely ground pork at the moment of spoilage. The bacterial numbers are presented as logarithms of colony forming units (cfu)/g.

Storage temperature	4°C				7°C			
	Surface layer		Core		Surface layer		Core	
	X	R	X	R	X	R	X	R
Total plate count								
-Plate count agar	7.0A	4.9-7.8	5.5B	5.2-5.9	6.0AB	4.2-7.5	3.6C	2.6-4.7
-APTS-agar	7.3A	4.9-8.5	5.3B	4.6-5.7	6.2AB	4.6-7.7	4.3C	3.0-6.6
-Blood agar	6.4A	5.0-7.7	5.2B	4.6-6.0	5.7AB	3.5-6.9	3.4C	2.5-4.4
Total count of anaerobes	5.4A	3.8-6.3	3.9B	2.7-4.8	2.5AB	1.0-4.8	1.3C	1.0-2.7
Lactic acid bacteria	4.8A	1.0-6.7	2.1B	1.0-3.4	4.8AC	2.9-6.9	2.9ABC	1.0-4.6
<i>Brochothrix thermosphacta</i>		< 2.0-6.6		< 2.0-4.2	4.3	2.7-5.3		< 2.0-3.5
Staphylococci + micrococci	2.6	2.3-3.0		< 2.0-2.9		< 2.0-5.7		< 2.0-4.8
Coliforms		< 1.0		< 1.0-1.3		< 1.0		< 1.0

X = mean

R = range

¹⁾ Means within the same horizontal line not followed by the same capital letter are significantly different from each other ($p < 0.05$).

the coarsely ground pork experiments. Thermal resistance experiments with coarsely ground pork were made three times for *Pseudomonas* strains and four times for *Streptococcus* strains.

Results and discussion

Bacterial flora in the product samples

The samples stored at 4°C were 7 - 10 weeks old and the samples stored at 7°C 4 weeks old at the moment of spoilage. Only the surface layer showed signs of spoilage. The total number of aerobically growing bacteria in the surface layer ranged from 3.5 to 8.5 log cfu (colony forming units)/g at the moment of spoilage (Table 1). At the moment of spoilage the core of the products contained 4.6 - 6.0 log cfu/g when the products had been stored at 4°C and 2.5 - 6.6 log cfu/g when they had been stored at 7°C. The mean of bacterial counts at 7°C was also significantly lower ($p < 0.05$) than at 4°C. The means of the bacterial counts both at 7°C and at 4°C were significantly lower ($p < 0.05$) in the core than in the surface layer. KORKEALA and LINDROTH(1987) have previously also shown that the core of ring sausage contains less bacteria than the surface layer.

The counts for anaerobically growing bacteria ranged from 1.0 to 6.3 log cfu/g the counts being higher ($p < 0.05$) in the surface layer than in the core of the product, and also higher ($p < 0.05$) after storing at 4°C than after storing at 7°C (Table 1). The number of lactic acid bacteria growing on Rogosa-agar ranged from 1.0 to 7.0 log cfu/g, and the surface layer contained significantly ($p < 0.05$) higher counts than the core at 4°C. Most samples contained *Brochothrix thermosphacta* and staphylococci/micrococci. Their counts were also higher in the surface layer than in the core of the products. Apart from one core sample, none of the samples contained coliforms over 1.0 log cfu/g.

Lactic acid bacteria growing on Rogosa-agar, *Brochothrix thermosphacta*, staphylococci/micrococci or anaerobically growing bacteria were not found to form the main bacterial types of the bacterial flora of the products. However, the bacterial flora of the product samples contained 1 - 3 main bacterial types which grew on the APT-agar.

Isolated bacteria

Bacterial strains representing the main bacterial types of the flora were isolated both from the surface layer (12 strains) and the core (12 strains) of

Table 2. Main bacterial strains of surface layer (a series) and core (b series) of cooked meat products made of coarsely ground pork, their survival after heat treatment at 72°C in APT-broth and their highest survival temperature on APT-agar (Gradiplate method).

Product/ main strains	gram + or	cell - shape	Range of bacterial counts (log cfu) inoculated in 5 ml APT-broth	Survival at 72°C in APT-broth (positive tests of 6)					Highest survival temperature (°C) on APT-agar range of 3 tests)
				15	30	60	90	120	
a series									
1/1	+	coccob.	6.1-6.8	2	0	0	0	0	< 45.5-49.1
1/2	+	coccob.	7.0-7.5	2	0	1	1	0	< 46.4-52.9
2/1	+	coccob.	7.3-7.4	1	0	0	0	0	< 46.2-51.1
2/2	+	coccob.	7.5-8.2	1	0	0	0	0	48.8 (1 test)
3/1	+	coccob.	7.4-7.7	0	0	0	0	0	47.4-49.0
3/2	+	coccob.	7.5-8.3	1	0	1	0	0	< 45.5 (1 test)
3/3	+	cocci	7.2-8.2	0	0	0	0	0	47.3-49.1
4/1	+	cocci	6.6-8.2	2	1	0	0	0	< 46.2-52.6
4/2	+	coccob.	4.8-7.3	1	0	0	0	0	55.3-58.2
5/1	+	cocci	3.7-6.2	0	0	0	0	0	47.3-47.3
7/1	+	cocci	5.7-7.2	0	0	0	0	0	48.1-50.2
7/2	+	coccob.	4.2-6.8	0	0	0	0	0	47.8-55.0
b series									
1/1	+	cocci	6.5-7.1	3	2	0	0	1	49.0-50.8
1/2 (1b2)	-	rods	6.8-6.9	6	6	5	5	4	48.5-52.7
2/1	+	cocci	6.0-7.5	3	2	0	0	0	< 45.5 (1 test)
2/2	+	cocci	6.0-7.9	5	4	0	0	0	46.8-51.3
3/1	+	cocci	5.7-7.4	6	6	6	5	2	50.3-56.6
3/2	+	cocci	6.9-8.2	5	4	0	0	0	< 46.4 (1 test)
3/3	+	cocci	5.9-7.1	1	1	2	2	2	48.5-50.3
4/1 (4b)	-	rods	7.2-7.8	6	4	3	2	0	50.5-58.9
5/1	-	rods	6.1-6.9	2	2	2	0	0	51.0-60.0
6/1 (6b)	-	rods	6.9-8.2	4	3	2	2	0	50.1-56.7
7/1	+	cocci	7.7-7.9	6	6	6	6	0	54.8-60.6
7/2 (7b2)	-	rods	6.5-7.1	3	4	3	3	3	58.8-62.2

the products (Table 2). The strains isolated from the surface layer of the products were gram-positive cocci or coccibacilli also growing in chains. The core of two products also contained gram-positive cocci as the main bacterial types (5 pcs.), which were isolated. The core of two products contained gram-positive coccus and gram-negative rod as the main bacterial strains, whereas the core of three products contained only gram-negative rod as a main strain. These 7 strains were also isolated.

The gram-positive coccus strains isolated were of the same type of lactic acid bacteria as those found by KORKEALA et al. (1988) to be the main spoilage organisms in cooked sausages. Five strains

also formed slime on APTS-agar (APT-agar + 1 % saccharose). The two most heat-resistant strains proved to be streptococci. It must be pointed out that the cores of the products contained gram-negative rods, although gram-negative bacteria are generally destroyed by heat treatment (HALLENBACH and POTTER 1981, PALUMBO et al. 1974). However, GARDNER (1966) also reports that cooked sausage may contain pseudomonads and achromobacteria. The gram-negative rod strains isolated were oxidative and they were, therefore, regarded as pseudomonads. They also grew on GSP-*Pseudomonas*-agar.

Table 3. Survival of some *Pseudomonas* strains after heat treatment at 72°C in APT-broth and in coarsely ground cured and cooked pork (3 test series). The strains were isolated from the core of cooked meat products made of coarsely ground pork.

Strain	Range of bacterial counts (log cfu) inoculated in 5 ml APT-broth	Survival at 72°C in APT-broth (postive tests of 6)					Range of bacterial counts (log cfu/g) inoculated in pork	Survival at 72°C in coarsely ground pork (log cfu/g)			
		Minutes						Minutes			
		5	15	30	60	90		0	5	15	30
1b ₂	6.8-8.7	-	4	4	4	4	7.1-8.0	7.0	¹⁾	1	2
4b	6.7-7.8	5	4	3	4	2	7.0-7.2	6.0	1	1	2
6b	6.4-7.1	6	4	4	4	2	7.4-7.7	5.8	2	2	0
7b ₂	7.3-8.2	6	6	4	5	2	7.6-8.5	7.2	3.4	3.6	2

¹⁾ Number of samples containing over 100 cfu/g. Total number of samples = 3.

Survival of isolated bacterial strains after heat treatment in APT-broth and on APT-agar

Seven strains from the surface layers survived after heating for 15 minutes at 72°C in either one or two of the six tests conducted (Table 2). Only one surface layer strain survived after heating on APT-agar at temperatures over 55°C. The highest survival temperatures of most surface layer strains on APT-agar varied around 50°C.

Most of the strains isolated from the core of coarsely ground pork products survived after heating for 30 minutes at 72°C in APT-broth in at least three of the six tests (Table 2). Two gram-positive cocci and one gram-negative rod survived after heating for 30 minutes at 72°C in all six tests. These three strains also survived after heating for 90 minutes in at least five of the tests. It is noteworthy that all gram-negative rods isolated from the core of coarsely ground pork products were heat-resistant to the extent that all of them survived after heating for 60 minutes at 72°C in APT-broth. The isolated strains survived heating on APT-agar at distinctly lower temperatures than after heating in APT-broth. Only three strains exhibited the highest survival temperature of over 60°C in some of the experiments.

Survival of four isolated *Pseudomonas* strains after heating at 72°C in coarsely ground cured pork

Of the *Pseudomonas* strains isolated in this study from the core of the cooked products made of coarsely ground pork, strain 7b₂ survived best after heating in coarsely ground cured pork (Table 3). The mean number of colony forming units decreased from log 7.2/g to log 3.6/g during heating for 15 minutes. After heating for 30 minutes inoculated pseudomonads were found in two of the three tests over log 2.0 cfu/g. After heating strains 1b₂, 4b and 6b in coarsely ground cured pork for 15 minutes each strain were found over log 2.0 cfu/g in 1 - 2 tests of the three.

The total plate counts of bacteria in the heat-treated coarsely ground experimental porks ranged between log 3.1 and 5.6 cfu/g. The control porks (not inoculated) contained pseudomonads and staphylococci/micrococci over 2.0 cfu/g only accidentally. They did not contain lactic acid bacteria over log 1.0 cfu/g or *Brochothrix thermosphacta* over log 2.0 cfu/g. The non-heated coarsely ground, cured pork did not exceed these values either.

When the inoculated pork samples which had been heated for 5 minutes at 72°C were incubated

Table 4. Bacterial flora of coarsely ground cured pork inoculated with four *Pseudomonas* strains after heating for 5 minutes at 72°C and incubating for 2 days at 30°C or 2-3 weeks at 4°C (3 test series).

	After heating Strains				2 days at 30°C Strains				2-3 weeks at 4°C Strains			
	1b ₂	4b	6b	7b ₂	1b ₂	4b	6b	7b ₂	1b ₂	4b	6b	7b ₂
Total plate count, APT-agar	¹⁾ X 4.1 s 0.8	4.4 1.4	4.9 0.7	5.1 0.1	5.3 1.0	6.4 1.8	6.3 1.5	5.2 0.1	5.2 0.4	5.3 1.8	4.2 0.3	6.8 2.0
Pseudomonads, inoculated	X 2.6 s 0.9	²⁾ 1	2	3.4 1.3	2	2	0	1	0	0	0	1
Staphylococci + micrococci	2	1	2	-	2	2	2	2	2	2	1	2
Lactic acid bacteria	³⁾ 0	0	0	0	0	0	0	0	0	0	0	0

¹⁾ log cfu/g

²⁾ Number of samples containing over 100 cfu/g. Total number of samples = 3.

³⁾ Number of samples containing over 10 cfu/g.

for two days at 30°C, the cfu number of strains 1b₂ and 4b rose, whereas the cfu number of strains 6b and 7b₂ decreased (Table 4). The total plate count for the porks rose to over log 6.0 cfu/g.

When the inoculated pork samples, which had been heated for 5 minutes at 72°C, were stored for 2 - 3 weeks at 6°C, the numbers of all inoculated strains decreased (Table 4). The total plate count rose distinctly only in the tests in which strain 7b₂ was used.

The incubation experiments both at 30°C and at 6°C indicate that even if pseudomonads survive after heat treatment, they are probably not a very serious factor causing spoilage in cooked meat products made of coarsely ground meat or in cooked meat products in general.

Survival of two most heat-resistant *Streptococcus* strains after heating at 72°C in coarsely ground cured pork

The mean cfu numbers of the porks before heating were 6.8 log cfu/g (strain 3b₁) and 7.6 log cfu/g (strain 7b₁) (Table 5). The decrease for both strains was about 1 log unit during heating for 2 minutes. The thermal death diminished when the heating

was continued, and was about 2 log units for 3b₁ and 0.5 log unit for 7b₁ over the next 13 min. Coarsely ground experimental porks contained staphylococci and micrococci over 2.0 log cfu/g only accidentally, while pseudomonads and *Brochothrix thermosphacta* never exceeded 2.0 log cfu/g. When the heat-treated and cooled experimental coarsely ground cured experimental porks were stored at 6°C for 2-3 weeks the cfu number of strain 3b₁ rose by 1 log unit and that of strain 7b₁ by 0.5 log unit. However, these bacteria probably do not constitute a spoilage problem because their cfu number decreases by a few log units during cooking and rises only slightly during storage.

Conclusion

The core of the pork products were found to contain heat-resistant streptococci and pseudomonads as the main bacterial strains, but they were not the cause of spoilage of the products. The bacteria which contaminated the products after heat treatment caused the spoilage in the surface layer of the products. However, the reasons for eventual core spoilage were not found in this investigation.

Table 5. Thermal death of *Streptococcus* strains 3b1 and 7b1 at 72°C in APT-broth (6 test series) and in coarsely ground cured pork (CCP) (4 test series).

Heating time, min	Cfu/ml in APT-broth				Cfu/g in CCP			
	3b1		7b1		3b1		7b1	
	X	s	X	s	X	s	X	s
0	6.3	0.9	6.5	0.2	6.8	0.5	7.6	0.5
0.5	4.2	0.8	5.2	0.5	6.3	1.0		
1	4.1	0.8	5.0	0.4				
1.5	4.0	1.7	5.0	0.3				
2	3.5	1.3	4.8	0.5	5.7	0.7	6.8	0.9
5	1.7	1.9	4.3	1.0	5.2	0.5	6.5	0.6
15	1.2	1.4	3.7	1.1	4.0	1.2	6.0	1.3
30	¹⁾ 4		2.1	0.9			5.5	1.1
60	1		2				5.3	1.4
120	0		1					

Cfu = colony forming unit

X = mean of colony forming units

s = standard deviation of mean

¹⁾ Number of samples containing colony forming units

References

- ALLEN, J. R. & FOSTER, E. M. 1960. Spoilage of vacuum-packed sliced processed meats during refrigerated storage. *Food Res.* 25: 19-25. (Ref. Paradis & Stiles 1978.)
- DAELMAN, W. & VAN HOOF, J. 1975. Einfluss des pH-Wertes, der Verwendung von Polyphosphat und der Lagerung auf die bakteriologische Beschaffenheit von Brühwurst und Brühwurstaufschnitt. *Archiv für Lebensmittelhygiene* 26: 213-217.
- DOWDELL, M. & BOARD, R. 1968. A microbiological survey of British fresh sausage. *J. Appl. Bacteriol.* 31: 378-396.
- DUITSCHAEVER, C. L. 1977. Bacteriological evaluation of some luncheon meats in Canadian retail market. *J. Food Prot.* 40: 382-384.
- GARDNER, G. 1966. A selective medium for the enumeration of *Microbacterium thermosphactum* in meat and meat products. *J. Appl. Bact.* 29: 455-460.
- 1968. Effects of pasteurization or added sulphite on the microbiology of stored vacuum-packaged baconburgers. *J. Appl. Bacteriol.* 31: 462-478.
- HALLENBACH, C. H. & POTTER, N. N. 1981. Effects of nitrite and sorbate on bacterial populations in frankfurters and thuringer cervelat. *J. Food Prot.* 44: 341-346.
- HILL, W. M., REAUME, J. & WILCOX, J. C. 1976. Total plate count and sensory evaluation as measures of luncheon meat shelf life. *J. Milk Food Techn.* 39: 759-762.
- KORKEALA, H. & LINDROTH, S. 1987. Differences in microbiological growth in the surface layer and at the centre of vacuum-packed cooked ring sausage. *Int. J. Food Microbiol.* 4: 105-110.
- , SUORTTI, T. & MÄKELÄ, P. 1988. Ropy slime formation in vacuum-packed cooked meat products caused by homofermentative lactobacilli and a *Leuconostoc* species. *Int. J. Food Microbiol.* 7: 339-347.
- MANTEL, T. & BECK, G. 1977. Zur microbiologischen Situation vorverpackter Brühwurst. *Fleischwirtsch.* 57: 245-247.
- PALUMBO, S. A., HUHTANEN, C. N. & SMITH, J.-L. 1974. Microbiology of frankfurter's process: *Salmonella* and natural aerobic flora. *Appl. Microbiol.* 27: 724-732.
- PARADIS, D. C. & STILES, M. E. 1978. A study of microbial quality of vacuum packaged sliced bologna. *J. Food Prot.* 10: 811-815.
- REUTER, G. 1969. Untersuchungen zur Mikroflora von Vorverpackten Aufgeschnittenen Brüh- und Kochwurst. *Proc. 15th Eur. Meet. Meat Res. Workers, Helsinki.* p. 124-134.

Manuscript received April 1993

Esko Petäjä
Eero Puolanne
Department of Food Technology
Meat Section
FIN-00014 University of Helsinki, Finland

SELOSTUS

Keitettyjen karkeahienonnetusta lihasta valmistettujen lihavalmisteiden bakteeriflooran hallitsevat bakteerikannat, niiden lämpöresistenssi ja vaikutus pilaantumiseen

ESKO PETÄJÄ ja EERO PUOLANNE

Helsingin yliopisto

Keitettyjen karkeahienonnetusta lihasta valmistettujen lihavalmisteiden pintaosan ja sisäosan bakteeriflooraa tutkittiin pilaantumishetkellä, kun valmisteita oli säilytetty 7°C tai 4°C:ssa. Flooraa hallitsevat bakteerikannat eristettiin ja niiden lämpöresistenssiä tutkittiin APT-liemessä, APT-agarilla ja karkeahienonnetussa suolatussa sianlihassa.

Valmisteiden pintaosan bakteeriflooraa hallitsevat kannat olivat kokkimaisia maitohappobakteereja lukumäärien vaihdella välillä 3,5-7,8 log pmy (pesäkkeen muodostava yksikkö)/g. Valmisteiden sisäosan bakteerifloora sisälsi valtaosin kokkimaisia maitohappobakteereja ja pseudomonadeja tai pseudomonadeja. Kaksi lämpöresisteintä maitohappobakteeria olivat streptokokkeja. Sisäosan bakteerien kokonaislukumäärät vaihtelivat välillä 2,5-6,0 log pmy/g.

Valmisteiden pintaosasta eristetyt bakteerikannat säilyivät lisääntymiskykyisinä vain satunnaisesti, kun niitä lämpökäsiteltiin 15 min 72°C:ssa APT-liemessä. Useimmat valmisteiden sisäosasta eristetyistä kannoista säilyivät lisääntymiskykyisinä vähintään kolmessa kokeessa kuudesta, kun niitä läm-

pökäsiteltiin 30 min 72°C:ssa APT-liemessä. On merkille pantavaa, että kaikki sisäosasta eristetyt pseudomonadikannat säilyivät lisääntymiskykyisinä, kun niitä lämpökäsiteltiin 60 min 72°C:ssa APT-liemessä, ja usein, kun niitä lämpökäsiteltiin karkeahienonnetussa sianlihassa 5 min 72°C:ssa. Inokuloitujen pseudomonadien lukumäärä laski, kun lämpökäsiteltyjä koelijoja säilytettiin 4°C:ssa. Siten ne eivät todennäköisesti ole vakava pilaantumistekijä karkeahienonnetusta lihasta valmistetuissa lihavalmisteissa tai keitettyissä lihavalmisteissa yleensä.

Lämpöresisteimpien streptokokkien lukumäärä (pmy) laski APT-liemessä jyrkemmin kuin karkeahienonnetussa sianlihassa, jossa lasku oli noin yksi log-yksikkö 15 minuutissa. Kun lämpökäsiteltyjä karkeahienonnettuja ja suolatuttuja sianlihoja säilytettiin 4 viikkoa 4°C:ssa, streptokokkien lukumäärä nousi enintään yhden log-yksikön. Kuitenkin myöskään streptokokit eivät aiheuttane pilaantumista, koska niiden lukumäärä laskee lämpökäsittelyn aikana selvästi ja nousee kylmävarastoinnin aikana vain vähän.