

Monitoring and predicting crop growth and analysing agricultural ecosystems by remote sensing

Tsuyoshi Akiyama¹ and Y. Inoue

National Institute of Agro-Environmental Sciences, 3-1-1, Kannondai, Tsukuba, Ibaraki 305, Japan

M. Shibayama

National Grassland Research Institute, Japan

Y. Awaya and N. Tanaka

Forestry and Forest Products Research Institute, Japan

LANDSAT/TM data, which are characterized by high spectral/spatial resolutions, are able to contribute to practical agricultural management. In the first part of the paper, the authors review some recent applications of satellite remote sensing in agriculture. Techniques for crop discrimination and mapping have made such rapid progress that we can classify crop types with more than 80% accuracy. The estimation of crop biomass using satellite data, including leaf area, dry and fresh weights, and the prediction of grain yield, has been attempted using various spectral vegetation indices. Plant stresses caused by nutrient deficiency and water deficit have also been analysed successfully. Such information may be useful for farm management. In the latter half of the paper, we introduce the Arctic Science Project, which was carried out under the Science and Technology Agency of Japan collaborating with Finnish scientists. In this project, monitoring of the boreal forest was carried out using LANDSAT data. Changes in the phenology of subarctic ground vegetation, based on spectral properties, were measured by a boom-mounted, four-band spectroradiometer. The turning point dates of the seasonal near-infrared (NIR) and red (R) reflectance factors might indicate the end of growth and the beginning of autumnal tints, respectively.

Key words: arctic area, biomass, environment, phenology, spectroradiometer, vegetation, yield

¹*Current address: Institute for Basin Ecosystem Studies, Gifu University, Yanagido, Gifu 501-11, Japan, e-mail: akiyama@green.gifu-u.ac.jp*

Table 1. Characteristics of earth observing satellites.

Satellite/ sensor	Launched/ yr/revolution	Wavelength range nm	Band	Ground resolution
NOAA/ AVHRR	U.S.A. (1978-) 1/2 d	C1: 580-680	Visible	1.1 km
		C2: 725-1100	Near-IR	1.1 km
		C3: 3550-3930	Mid-IR	1.1 km
		C4: 11.5-12.5 μ	Thermal	1.1 km
LANDSAT/ MSS	U.S.A. (1972-) 16(18) d	MSS4: 500-600	Visible	80 m
		MSS5: 600-700	Visible	80 m
		MSS6: 700-800	Near-IR	80 m
		MSS7: 800-1100	Near-IR	80 m
LANDSAT/ TM	U. S.A. (1984-) 16 d	TM1: 450-520	Visible	30 m
		TM2: 520-600	Visible	30 m
		TM3: 630-690	Visible	30 m
		TM4: 760-900	Near-IR	30 m
		TM5:1550-1750	Mid-IR	30 m
		TM7:2080-2350	Mid-IR	30 m
		TM6:10.4-12.5 μ	Thermal	120 m
SPOT/ HRV	France (1986-) 26 d	B1: 500-590	Visible	20 m
		B2: 610-680	Visible	20 m
		B3: 740-900	Near-IR	20 m
		B4: 510-730	Panchromatic	10 m
IRS-1A/ LISS-I	India (1988-) 22 d	B1: 450-520	Visible	72.5 m
		B2: 520-590	Visible	72.5 m
		B3: 620-680	Visible	72.5 m
		B4: 770-860	Near-IR	72.5 m
ERS-1/ SAR	ESA (1991-) 35 d	C : 5.3 GHz	Microwave	30 m
JERS-1/ OPS	Japan (1992-) 44 d	B1: 520-600	Visible	18 m
		B2: 630-690	Visible	18 m
		B3: 760-860	Near-IR	18 m
SAR		L:1.275 GHz	Microwave	18 m

Monitoring of crops and environment by earth observation satellites

Characteristics of satellite sensors and vegetation indices

Satellite sensors

More than two decades have passed since LANDSAT-1 was launched by NASA in 1972. At present, several kinds of satellites are orbiting and observing the surface of the earth (Table 1). By the end of this century, still more, new

satellites with special missions will be launched.

In the NOAA series, two satellites are maintained in polar orbit, one in a morning orbit and the other in an afternoon orbit. They provide a wide range of data, including sea surface temperature, cloud cover, data for land surface studies, temperature and humidity profiles, and ozone concentrations.

The LANDSAT and SPOT satellites provide high resolution imagery in the range of visible and infrared bands. They are used extensively for high resolution land surface studies.

The ERS series concentrates on global and regional environmental issues, making use of ac-

tive microwave techniques, which enable a range of measurements to be made of the land, sea, and ice surface independent of cloud cover.

The aim of JERS-1 is to observe the earth using optical sensors and a high resolution synthetic aperture radar. Land surveys and monitoring of various resources are the main applications of this satellite.

Spectral vegetation indices

Based on the fact that green vegetation absorbs red (R) wavelengths but reflects near-infrared (NIR) wavelengths, several spectral vegetation indices (VI) derived from satellite-borne data have been proposed. VI, including ratio vegetation index (RVI, NIR/R), difference vegetation index (DVI, $NIR-R$), normalized vegetation index (NDVI, $(NIR-R) / (NIR+R)$), are used the most by vegetation scientists. In addition, perpendicular vegetation index (PVI), soil adjusted vegetation index (SAVI), and K values have been proposed to eliminate background soil effects from vegetation.

Crop monitoring from space

In recent decades, many attempts have been made at extracting agro-environmental information at the regional level, and some useful technologies have been developed. As a result, the following information can be obtained by space-borne sensors with high spectral/spatial resolutions, such as LANDSAT /TM and SPOT/HRV: 1) crop inventory and planting acreage estimates, 2) leaf area and phytomass estimates and yield prediction, 3) crop stress detection, and 4) agro-environmental survey including soil, vegetation, water and atmosphere.

Crop discrimination and planting acreage estimation

From 1974 onwards, in the Large Area Crop Inventory Experiment (LACIE), jointly undertaken by NASA, NOAA and USDA, satellite remote sensing technology was applied, on an experimental basis, to forecast harvests in the major

wheat producing areas of the world using LANDSAT/MSS data. As a result, a 1977 real-time forecast of the wheat production of the Soviet Union indicated that the system could operate and could be applied to other areas and other crops (MacDonald and Hall 1980). This project was followed by AgRISTARS, which focused on the assessment of crop conditions from space (AgRISTARS 1983). Many attempts at crop discrimination and planting acreage estimation have been made world-wide for a variety of crops using LANDSAT/MSS, LANDSAT /TM and SPOT/HRV data.

Recent studies and classification accuracies are summarized in Table 2. Most of the studies report accuracies exceeding 80%. To illustrate, Table 3 shows crop discrimination results using LANDSAT/TM data in the Tokachi district of Japan. Here, the major crops were classified with 90% accuracy (Fukuhara et al. 1988). In general, the accuracy of crop discrimination is strongly dependent on the spectral/spatial resolution of the satellite sensor, the timely acquisition of data at suitable crop stages, and ample field size for detection purposes.

Biomass, leaf area and yield estimation

Several spectral vegetation indices (VI) have been proposed to estimate vegetational information, including biomass, yield and leaf area. RVI, which employs the ratio between NIR and R, is often applied for the estimation of aboveground dry weight, fresh weight and leaf area.

Estimates of pasture grass yield at the first cut were made over Tochigi prefecture in central Japan, using LANDSAT/MSS data collected on May 22, 1979. The analysis was conducted by attempting first a land-use classification to identify pasture and meadows. Secondly, a multiple regression analysis was performed to estimate the yield of the first cutting in each pasture plot. To design the regression model, 26 plots with known yield data were prepared in advance. A high correlation was observed between the actual forage yield at the first cutting and the estimated yield based on LANDSAT/MSS using the four bands (Akiyama et al. 1985).

Table 2. Results of crop discrimination studies using satellite data.

Country/ District	Satellite/ Sensor	Main crops	Accuracy	Authors
U.S.A./ Corn belt	Landsat/ MSS	Corn, Soybean	83.9%	Batista et al. 1985
U.S.A./ Wisconsin	Landsat/ MSS	Alfalfa, Corn, Oats	89.8%	Lo et al. 1986
Hungary	Landsat/ MSS	Corn, Alfalfa Sunflower etc.	90%	Csillag 1986
Argentine/ Buenos Ires	Landsat/ MSS&TM	Corn, Soybean Sorgham etc.	80%	Badhwar et al. 1987
U.S.A./ NY State	Landsat/ TM	3 Vegetables 4 Crops	>90% >75%	Williams et al. 1987
Japan/ Tokachi	Landsat/ TM	7 Crops	90.1%	Fukuhara et al. 1988
Hungary/ Kiskore	SPOT/XS Landsat/TM	Wheat, Soya Alfalfa	XS: 85-88% TM: 87-91%	Büttner et al. 1989
U.K./ East Anglia	SPOT/ HRV	10 Vegetation types 4 Vegetation types	71% 88%	Jewell 1989

Table 3. Results of crop classification studies using LANDSAT/TM data in Tokachi district, Japan (Fukuhara et al. 1988).

Crop name	Performance (%)								
	Su	Po	Ad	So	Co	Wh	Pa	Fo	Others
Sugar beet (Su)	<u>96.3</u>	0	0	0	0	0	1.7	0	2.0
Potato (Po)	0	<u>98.5</u>	0	0	0	0	0.5	0	1.0
Adzuki bean(Ad)	0	0	<u>97.3</u>	0.5	1.6	0	0	0	0.6
Soybean (So)	0	0	0	<u>100.0</u>	0	0	0	0	0
Corn (Co)	0	0	1.2	0	<u>59.5</u>	14.4	0	22.5	2.4
Wheat (Wh)	0	0	0	0	3.5	<u>95.1</u>	0	0.2	1.2
Pasture (Pa)	0	0	1.0	0	0	0	<u>88.0</u>	0	11.0
Forest (Fo)	0	0	1.0	0	6.1	0.6	0.5	<u>90.2</u>	1.6

Underlined numbers indicate the proportion of crops correctly classified (%).

Recent results of grain yield prediction for rice and wheat are presented in Table 4. A high accuracy was attained with a multiple regression model using 2 or 3 TM bands for rice. Meteorological damage caused by cold wind and flooding are also analysed effectively using satellite data. For example, rice damage caused by flooding was highly correlated with the turbidity of water (Yamagata and Akiyama 1988), which reflected in TM bands 2 and 3.

Crop stress detection

Plants experience various kinds of stresses during growth, including stresses caused by atmospheric factors, by rhizospheric factors, or biotic stresses caused by a variety of organisms. Moreover, in recent years, it has often been reported that toxic materials in the environment resulting from human activities can inflict damage on crops and forests, as in the case of acid rain. If farmers could detect various crop stresses in the

Table 4. Results of crop yield prediction using satellite data.

Crops	Country/ district	Satellite/ sensor	Date of data acquisition	Model used	Correlation, Standard deviation	Authors
Rice	Thailand/ Saraburi	LANDSAT/ TM	09/Dec/88 27/Dec/88	MRM (TM 1, 5, 7) Growth model	R ² = 0.85 to 0.92	Tennakoon et al. 1992
Rice	India/ Orissa	IRS-1A/ LISS-1	12,13/Oct/ 88	Area Weighted Average Ratio	SD: -2% to +14%	Patel et al. 1991
Rice	Japan/ Ishikari	LANDSAT/ TM	23/Sep/86	MRM (TM2, 3, 4, 5 & TM4/TM3)	R ² = 0.953	Mubekti et al. 1991
Cold wind damaged rice	Japan/ Ishikari	LANDSAT/ MSS	19/Sep/80	MRM (MSS5, 6)	R= 0.91	Miyama et al. 1983
Flood damaged rice	Japan/ Ibaraki	LANDSAT/ TM	6/Aug/86	MRM (TM2, 3)	R=0.972	Yamagata and Akiyama 1988
Wheat	Brazil/ Sao Paolo	LANDSAT/ TM	24/Jun/86 27/Jun/87	Meteo. model TM4/TM3	R ² = 0.65	Rudorff and Batista 1991
Wheat	Japan/ Ishikari	LANDSAT/ TM	27/Jun/86 29/May/90	MRM (TM2, 3)	R ² = 0.66 to 0.79	Shiga 1993
Wheat	India/ Sultanpur	LANDSAT TM	22/Feb/86	NDVI, RVI Layers method	Variance 0.1958	Singh et al. 1992
Wheat	India/ Haryana	LANDSAT/ TM	26/Jan/89	Linear yield-spectral index	Deviation +14% to -18%	Sharma et al. 1993
		IRS/1A/ LISS-1	18/Feb/89			

MRM: Multiple Regression Model
 R: Multiple regression coefficient
 R²: Coefficient of determination
 SD: Standard deviation

early stages of an infection or a nutrient deficiency before any symptoms appear, a timely intervention could prevent the damage to the yield and save the quality of the final product.

Water deficit in plants can be clearly detected by NIR and mid-infrared (MIR) wavelength reflectances, which indicate the thermal changes caused by stomatal movement. The same band can be used for the crop stress caused by soil-borne diseases, as it enables to detect the wilting of the leaves (Torigoe 1992).

Stresses due to meteorological factors do not always show up early. Spikelet sterility is often caused by low temperatures during the flowering stage of rice in the northern part of Japan. Usually, the fertile ears die off at ripening, while the sterile ears remain green at harvest time. Using LANDSAT /MSS imagery, Mi-

yama et al. (1983) were able to create a map of the cold damage to rice on the Ishikari plane of northern Japan.

Agro-environmental monitoring

Soil survey

Many scientists have classified soil types using satellite data. Information on organic matter and moisture content in the soil is particularly important for farmers. Fukuhara et al. (1980) were able to determine differences in the moisture contents of the soil using LANDSAT/MSS data in the Tokachi district in Japan. In the same area, Hatanaka et al.(1989) classified the organic matter content in the soil, using band 3 reflectance data from LANDSAT/TM obtained in

May, when the surface of the upland field was almost bare. The classification results were plotted on a 1/50,000 scale map, and used by officers of the agricultural extension services.

Land evaluation

It is possible for the developing countries to select the land suitable for agricultural development using satellite data. These data can provide several thematic maps relating biomass, soil moisture and present landuse. When these thematic maps are overlaid on existing meshed topographic or geographic maps, we can obtain value-added information on the productivity of the land (Akiyama et al. 1987).

In recent years, the need to quantify the function of agricultural land has become increasingly important in Japan. Attempts have been made to quantify the multiple functions of such land, for example the impact of the cultivation of paddy rice on soil conservation, landscape maintenance or the capacity for water purification. Remote sensing by a satellite is a prominent tool for such evaluations.

Arctic Science Project with Finland

International Cooperative Study on Observation of Variabilities in the Arctic Atmosphere, Hydrosphere and Biosphere, and their Interactions

One of our most serious concerns today is the fact that global environment change (chiefly deterioration) is taking place at an unprecedented speed, because of ever-growing human activity. We have come to realize that polar regions are very important areas for understanding processes of global change; no less so than the tropical and mid-latitude areas, where most of the human race live.

The ongoing environmental changes are not

only global in scale, but also varied, and involve diverse and complicated processes, calling for an interdisciplinary approach to address the issue. In this context, in 1990 the Science and Technology Agency of Japan organized a 5-year program "International Cooperative Study on Observation of Variabilities in the Arctic Atmosphere, Hydrosphere and Biosphere, and their Interactions" in cooperation with 15 governmental institutes, laboratories and universities.

The Arctic Science Project consists of three sections, namely, 1) Arctic oceanography and glaciology, 2) Atmospheric chemistry, 3) Arctic vegetation and agricultural environment. The Arctic vegetation team enrolled the assistance of the Technical Research Centre of Finland (VTT), Turku University and the Agricultural Research Centre of Finland. Here, we report on two topics investigated by the vegetation monitoring sub-teams in the Arctic Science Project.

Vegetation change monitoring in the boreal area

This part of the project was conducted by scientists in the Forestry and Forest Products Research Institute of Japan working together with VTT of Finland. An outline of the collaborative research is given below.

The study concentrated on the monitoring of vegetation change using satellite data. Seasonal and successional changes of spectra, as a basis for long term monitoring, were studied over the southern boreal forest zone using LANDSAT/TM images (Awaya et al. 1995).

Two monitoring methods were compared at a test site in Kevo, at the northern tip of Finland. The first was boundary detection using Laplacian filtering, and the second biomass change detection by subtraction between two normalized difference vegetation indices (NDVI) of LANDSAT/MSS images acquired in 1972 and 1987. A zonal shift of arctic vegetation was not captured using these methods, but damage to birch forests by the moth caterpillar were clearly detected. This phenomenon appeared to




Fig.1 Seasonal changes of spectra for different the tree species estimated using TM data (Awaya et al. 1995). The spectral changes, from early spring to late autumn, were drawn using eight TM images taken between 1985 and 1993. The numbers in the figure caption represent the year of planting of the trees.

be caused by a combination of the moth caterpillar and by temperatures lower than the average in the latter half of the 1960's. It suggested that the vegetation changes are occurring that are related both ecological processes, i.e. the damage and the recovery of the trees, and the effects of global climate change.

The seasonal spectral patterns of the southern boreal forests were clearly identified in the TM3, TM4 and TM5 of LANDSAT in Hokkaido, Japan. Though the spectra of the young spruce stands seemed to be affected by mixed broadleaved trees in TM3 and TM4, seasonal spectral patterns could be classified into two types, evergreen and deciduous trees, in TM5 (Fig.1, left). Forest types classified according to the major tree showed constant intensities in a normalized difference ratio of TM5 and TM3 (ND53) over the growing season (Fig.1, right). The successional spectral patterns of the spruce stands varied between seasons and channels. However, the relationship between the digital number and the stand age was expressed using an exponential function (not shown here).

Spectral properties of subarctic ground vegetation

This experiment was conducted by the National

Institute of Agro-Environmental Sciences, and the National Grassland Research Institute of Japan collaborating with VTT of Finland and Turku University (Shibayama et al. 1995).

Ground truth information for remote sensing, to assess the phenology of boreal plants, is needed in global scale climate change research activities. A boom-mounted, four-band spectroradiometer was installed to measure seasonal radiances in the green (520-600nm), red (630-690nm), near-infrared (765-900nm) and mid-infrared (1570-1730nm) spectral bands from boreal shrub canopies over the growing season in northernmost Finland at Utsjoki. It measured the spectra of four fixed ground plots, each one once per hour, from early June to mid-September between 0600 and 1800 hrs local time. A reflectance panel was also measured a few minutes before and after measurement of each plot. The plant phenology on each plot was also observed weekly during the experiment. Seasonal reflectance factors for each plot in each band were calculated based on a new calibration method. It involves a correction for the degradation of the reference panel. A hand-held spectroradiometer was also used to measure the plant canopies and the reference panel in the early autumn. The turning point dates of the seasonal near-infrared (765-900nm) and red (630-690nm) reflectance factors might indicate the end of

Akiyama, T. et.al.: Analysing agricultural ecosystems by remote sensing




Fig. 2. Seasonal patterns of daily reflectance factors at bands 560, 658, 833 and 1649nm, and the bi-band ratio of the 560 and 658 nm reflectance factors measured for the plant plots (Plots#1,#2) by the automated four-band spectroradiometer at Kevo. The lateral bars show the 95% confidence intervals of the estimated intersection points (the turning point dates of the radiometric variables) (Shibayama et al. 1995).

growth and the beginning of the autumnal tints, respectively. The ratio of the green (520–600nm) and red band reflectance factors, however, seemed to be more accurate in predicting these turning points (Fig.2).

Acknowledgements. We wish to express our gratitude to Dr. T. Häme, Dr. A. Salli and Dr. A. Lohi of the Technical Research Centre of Finland and Dr. M. Alanen and Dr. S. Neuvonen of the Kevo Subarctic Research Institute, University of Turku, for their diligent support in the Arctic Science Project. The authors are also deeply grateful to Professor T. Mela for his helpful suggestions.

References

- AgRISTARS Program Management Group** 1983. Agriculture and resources inventory surveys through aerospace remote sensing. Research Report, Fiscal Year 1982, APJ2-0393. p. 1–45.
- Akiyama, T., Miyama, K., Soemarman, H. & Setiyono, J.** 1987. Land evaluation system using Landsat data for agricultural development. Application of PATTERN method in North Sumatera. *Journal of the Japanese Agricultural Systems Society* 3(2): 74–89.
- , **Yasuda, Y., Emori, Y. & Miyama, K.** 1985. Grassland diagnosis by remote sensing techniques. 2. First cutting yield estimate of pasture using Landsat multispectral data. *The Journal of Japanese Society of Grassland Science* 31: 97–103. (in Japanese)
- Awaya, Y., Tanaka, N., Häme, T. & Lohi, A.** 1995. Vegetational change monitoring in the boreal area using remote sensing. Changes in the northern Finland and spectral characteristics of spruce. *Proceedings of International Arctic Science Symposium, Tsukuba, D:* 30–61.
- Badhwar G.D., Gargantini, C.E. & Redondo, F.V.** 1987. Landsat classification of Argentina summer crops. *Remote Sensing of Environment* 7: 265–281.
- Batista, G.T., Hixson, M.M. & Bauer, M.E.** 1985. LANDSAT MSS crop classification performance as a function of scene characteristics. *International Journal of Remote Sensing* 6: 1521–1533.
- Büttner, G. & Csillag, F.** 1989. Comparative study of crop and soil mapping using multitemporal and multispectral SPOT and Landsat Thematic Mapper data. *Remote Sensing of Environment* 29: 241–249.
- Csillag, F.** 1986. Comparison of some classification methods on a test-site. *International Journal of Remote Sensing* 7: 1705–1715.
- Fukuhara, M., Amano, T. & Miyaji, N.** 1988. Creation of cropping map using Landsat TM data. *Proceedings of 1988 Annual Meeting of the Japan Society of Photogrammetry and Remote Sensing, J-2:* 169–172. (in Japanese)
- , **Hayashi, S., Yasuda, Y., Emori, Y. & Iisaka, J.** 1980. Soil moisture analysis for soil mapping. *Proceedings of the Machine Processing of Remotely Sensed Data Symposium, Purdue University, Indiana, U.S.A.* p. 1–10.
- Hatanaka, T., Shiozaki, H., Fukuhara, M., Miyaji, N. & Saito, G.** 1989. Estimation of organic matter contents of upland soils with Landsat TM data. *Japanese Journal of Soil Science and Plant Nutrition* 69: 426–431. (in Japanese)
- Jewell, N.** 1989. An evaluation of multi-date SPOT data for agriculture and land use mapping in the United Kingdom. *International Journal of Remote Sensing* 10: 939–951.
- Lo, T.H.C., Scarpace, F.L. & Lillesand, T.M.** 1986. Use of multitemporal spectral profiles in agricultural land-cover classification. *Photogrammetric Engineering and Remote Sensing* 52: 535–544.
- MacDonald, R.B. & Hall, F.G.** 1980. Global crop forecasting. *Science* 208: 670–679.
- Miyama, K., Sato, H., Yasuda, Y. & Emori, Y.** 1983. An applied study on a remote sensing techniques to survey agricultural land. Survey technique for geographical distribution of damage to rice from cold weather using Landsat MSS data. *Transactions of JSIDRE* 105: 27–35. (in Japanese)
- Mubekti, Miyama, K. & Ogawa, S.** 1991. Study on rice yield distribution by using Landsat TM data. *The Hokkaido National Agricultural Experiment Station, Rural Development Research* 5: 101–113.
- Patel, N.K., Ravi, N., Navalgund, R.R., Dash, R.N., Das, K.C. & Patnaik, S.** 1991. Estimation of rice yield using IRS-1A digital data in coastal tract of Orissa. *International Journal of Remote Sensing* 12: 2259–2266.
- Rudorff, B.F.T. & Batista, G.T.** 1991. Wheat yield estimation at the farm level using TM Landsat and agrometeorological data. *International Journal of Remote Sensing* 12: 2477–2484.
- Sharma, T., Sudha, K.S., Ravi, N., Navalgund, R.R., Tomar, K.P., Chakravarty, N.V.K. & Das, D.K.** 1993. Procedures for wheat yield prediction using Landsat MSS and IRS-1A data. *International Journal of Remote Sensing* 14: 2509–2518.
- Shibayama, M., Salli, A., Häme, T., Iso-livari, L., Morinaga, S., Inoue, Y. & Akiyama, T.** 1995. Spectral properties of subarctic ground vegetation during the growth period. -Preliminary results of a 1994 experiment in the northernmost Finland. *Proceedings of the International Arctic Science Symposium, Tsukuba, Japan, D:* 6–29.
- Shiga, H.** 1993. Agricultural information system in Hokkaido: Integration of arable land information and application of remote sensing for land evaluation. *Journal of the Japanese Agricultural Systems Society* 9: 32–39. (in Japanese)
- Singh, R., Goyal, R.C., Saha, S.K. & Chhikara, R.S.**

Akiyama, T. et.al.: Analysing agricultural ecosystems by remote sensing

1992. Use of satellite spectral data in crop yield estimation surveys. *International Journal of Remote Sensing* 13: 2583–2592.

Tennakoon, S.B., Murty, V.V.N. & Eiumnoh, A. 1992. Estimation of cropped area and grain yield of rice using remote sensing data. *International Journal of Remote Sensing* 13: 427–439.

Torigoe, Y. 1992. Low input sustainable agriculture and information system: Developing integrated pest manage-

ment for soilborne disease. *Journal of the Japanese Agricultural Systems Society* 8: 149–157. (in Japanese)

Williams, V.L., Philipson, W.R. & Philpot, W.D. 1987. Identifying vegetable crops with Landsat Thematic Mapper data. *Photogrammetric Engineering and Remote Sensing* 53: 187–191.

Yamagata, Y. & Akiyama, T. 1988. Flood damage analysis using multi-temporal Landsat Thematic Mapper data. *International Journal of Remote Sensing* 9: 503–514.

SELOSTUS

Maatalousekosysteemien analysointi ja sadon ennustaminen kaukokartoituksen avulla

Tsuyoshi Akiyama, Y. Inoue, M. Shibayama, Y. Awaya ja N. Tanaka

National Institute of Agro-Environmental Sciences, National Grassland Research Institute ja Forestry and Forest Products Research Institute, Japani

Tämän artikkelin alussa arvioidaan muutamia satelliittikaukokartoituksen uusimpia sovelluksia maataloudessa ja lopussa esitellään japanilais-suomalainen yhteistyöhanke Arctic Science Project. LANDSAT/TM-satelliittikuva-aineistoa voidaan käyttää maataloustuotannon seurannassa, sillä kuvien spektri/spaatialinen tarkkuus on hyvä. Erittely- ja kartoitustekniikat ovat kehittyneet niin paljon, että viljalajit voidaan erottaa toisistaan 80 % tarkkuudella. Satelliittiaineiston avulla on arvioitu sadon biomassaa, esimerkiksi lehtialaa sekä kuiva- ja tuorepainoa, ja en-

nustettu viljasatoa erilaisia laajaspektrisiä kasvillisuusindeksejä käyttäen. Myös ravinteiden ja veden puutteen aiheuttamaa stressiä on onnistuttu analysoidaan. Arctic Science Projectissa seurattiin boreaalisia metsäalueita LANDSAT-satelliittiaineiston avulla. Subarktisen pohjakasvillisuuden spektriominaisuuksiin perustuvia fenologisia muutoksia mitattiin nelikanavaisella spektroradiometrillä. Käännekohtat vuodenaikaisissa lähes-infrapuna- ja punaheijastu- missa saattavat osoittaa kasvun päättymistä ja ruskan alkua.