

cluded in *Compsonannus* by Wagner, differs strikingly from the other species (head short and broad, basal margin of vertex distinctly keeled; 2nd antennal joint black with two pale rings; upper surface without dark dots; erect hairs on upper surface black; membrane without dark mottling; hind tibia with large black spots, tibial spines black) and is not congeneric with them. Its generic position will be treated elsewhere.

Acknowledgements. I am greatly indebted to Dr. A. Jansson and Prof. H. Strümpel for allowing me to examine type material preserved in the Helsinki and Hamburg Museums.

References

- Lindberg, H. 1958: Hemiptera Insularum Caboverdensium. — Comment. Biol. 19(1):1–246.
- Linnauori, R. E. 1986: Heteroptera of Saudi Arabia. — Fauna of Saudi Arabia 8:31–197.
- Reuter, O. M. 1902: Capsidae novae mediterraneae IV. — Öfv. Finska Vet. Soc. Förhandl. 44:51–70.
- Wagner, E. 1965: Die Gattung Atomoscelis, Reuter und ihre Verwandten in der Paläarktis (Hem. Het. Miridae). — Notulae Entomol. 45:74–92.
- 1968: Über einige Miriden aus Marokko (Heteroptera). — Notulae Entomol. 48:103–125
- 1971: Die Gattung *Compsonannus* Reuter, 1902 (Heteroptera, Miridae). — Notulae Entomol. 51:95–98.
- 1974: *Compsonannus ovatus* sp. n. (Heteroptera, Miridae). — Deutsche Entomol. Zeitsch. 21:165–166.

Received 4.V.1989

Erratum

Disney, R. H. L. 1990: Two new species and a revised key to *Rhynchomicropteron* (Diptera, Phoridae). — Entomol. Fennica 1:21–24.

Disney, R. H. L. 1990: A striking new species of *Megaselia* (Diptera, Phoridae) from Sulawesi, with re-evaluation of related genera. — Entomol. Fennica 1:25–31.

Disney, R. H. L. 1990: A key to *Diplonevra* males of the Australasian and Oriental Regions, including two new species (Diptera, Phoridae). — Entomol. Fennica 1:33–39.

Pages 22, 26, 34–37: For # read ♂
For ♂ read ♀