

## Nomenclatural note on the genus *Moissonia* Reuter (Hemiptera, Miridae, Phylinae)

R. E. Linnauori & M. M. Al-Safadi

Linnauori, R. E. & Al-Safadi, M. M. 1993: Nomenclatural note on the genus *Moissonia* Reuter (Hemiptera, Miridae, Phylinae). — Entomol. Fennica 4:233–234.

The taxonomic position of the genus *Moissonia* Reuter is discussed. *Ellenia* Reuter, *Marshalliella* Poppius, and *Melanotrichiella* Poppius are junior synonyms of *Moissonia*. *Marshalliella obscuricornis* Poppius and *Psallus labeculus* Odhiambo are synonyms of *Moissonia punctata* (Fieber).

R. E. Linnauori, Somersoja, FIN-21220 Raisio 22, Finland  
M. M. Al-Safadi, P.O. Box 392, Sana'a, Republic of Yemen

The article is based on field work made by the senior author in 1992 in the Republic of Yemen. A zoogeographically interesting synonymy between the genera *Moissonia* Reuter and *Ellenia* Reuter was detected.

### *Moissonia* Reuter, 1894

*Moissonia* Reuter 1894:148. Type species: *Agalliastes punctatus* Fieber.

*Ellenia* Reuter 1910:168. Type species: *Capsus cuneatus* Stål, syn. n.

*Marshalliella* Poppius 1914:75. Type species: *M. obscuricornis* Poppius, syn. n.

*Melanotrichiella* Poppius 1914:80. Type species: *M. anulicornis* Poppius, syn. n.

Reuter (1894:148) described the genus *Moissonia* with *Agalliastes punctatus* Fieber as the type species. The genus has remained enigmatic ever since. Wagner (1975:65–66), for example, misinterpreted the claw structure (Fig. 1b) regarding the fleshy parempodia as pulvilli, and located the genus close to *Campylomma* Reuter with which it is not actually related. *Moissonia punctata* was formerly known only from Spain (Malaga and Alicante) and Algeria (Oran). Prof. Attilio Cara-

pezza, Palermo, recently found the species in Sicily and kindly sent us several specimens. An examination of these surprisingly revealed that *Moissonia* is identical with the well-known intertropical genus *Ellenia* Reuter. The type species *M. punctata* is even a senior synonym of the common East African species *E. obscuricornis* (Poppius), which is distributed from South Africa to Ethiopia and Yemen, and, undoubtedly during the Pluvial period along the Red Sea hills, even to the Mediterranean area. The genera *Marshalliella* Poppius and *Melanotrichiella* Poppius were declared synonyms of *Ellenia* in Carvalho 1952:76 and Linnauori 1973:89–90 and 1975:87.

### *M. punctata* (Fieber, 1861)

Fig. 1a

*Agalliastes punctatus* Fieber 1861:311.

*Moissonia punctata* Reuter 1894:148.

*Marshalliella obscuricornis* Poppius 1914:76, syn. n.

*Psallus labeculus* Odhiambo 1959:518–521, syn. n.  
(synonymized with *Marshalliella obscuricornis* in Linnauori 1973:90).

Material studied: Ethiopia: Sululta, 2 exx, 11.VI.1963, Linnauori. Yemen: Sana'a, 1 ex, VII–IX.1991, A. van




Fig. 1. *Moissonia punctata* (Fieber): a) vesica (ex from Sana'a). — *M. kilimana* (Poppius): b) claw.

Harten. Italy: Sicily, Etna, Rif. Sapienza, alt. 1600 m, several exx, 24.VII.1990, A. Carapezza.

Redescription in Schuh 1974:158–159. Vesica illustrated in Fig. 1a.

Biology: Undoubtedly restricted to the Compositae genus *Senecio*. Schuh found it on *Senecio burchelli* and *S. angustifolius* in South Africa, Carapezza on *S. aetnensis* in Sicily.

Distribution: East African. A common species in South and East Africa.

## References

- Carvalho, J. C. M. 1952: On the major classification of the Miridae (Hemiptera). — An. Acad. Brasil. Sci. 24:31–110.
- Fieber, F. X. 1861: Die europäischen Hemipteren. Halbflügen (Rhynchota Heteroptera) II: 113–444. — Druck und Verlag von Carl Gerold's Sohn. Wien.
- Linnavuori, R. E. 1973: A collection of Heteroptera from Katanga, with remarks on some species from other parts of the Ethiopian Region. — Ann. Entomol. Fennici 39:70–94.
- 1975: Hemiptera of the Sudan, with remarks on some species of the adjacent countries. 4. Miridae and Iso-metopidae. — Ann. Zool. Fennici 12:1–118.
- Odhiambo, T. 1959: Notes on the East African Miridae (Hemiptera) XIII. Twelve new species of Psallus Fieber. — Ann. Mag. Nat. Hist. (13)2:513–544.
- Poppius, B. 1914: Die Miriden der Äthiopischen Region II. — Acta Soc. Sci. Fennicae 44:1–136.
- Reuter, O. M. 1894: Ad Cognitionem Capsidarum II. Capsidae palaearcticae. — Rev. d'Entomol. 13:128–152.
- 1910: Neue Beiträge zur Phylogenie und Systematik der Miriden nebst einleitenden Bemerkungen über die Phylogenie der Heteropteren-Familien. Mit einer Stammbaum. — Acta Soc. Sci. Fennicae 37(3):1–169.
- Schuh, R. T. 1974: The Orthotylinae and Phylinae (Hemiptera, Miridae) of South Africa with a phylogenetic analysis of the ant-mimetic tribes of the two subfamilies for the world. — Entomol. Americana 47:1–332.
- Wagner, E. 1975: Die Miridae Hahn, 1831, des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera, Heteroptera). Teil 3. — Entomol. Abh. Staatl. Mus. Tierkunde Dresden 40, Suppl., 483 pp.

Received 29.XII.1992