

# Karyotype dispersal of the common lizard *Zootoca vivipara* (Lichtenstein, 1823) in eastern and northeastern Fennoscandia

Larissa Kupriyanova, Martti Niskanen & Tuula A. Oksanen

*Kupriyanova, L., Zoological Institute, Russian Academy of Sciences, St. Petersburg 199034, Russia. E-mail: larissakup@zin.ru*

*\*Niskanen, M. & Oksanen, T., University of Jyväskylä, Department of Biological and Environmental Science, P.O. Box 35, FI-40014 University of Jyväskylä, Finland. E-mail: martti.e.a.niskanen@jyu.fi (\*corresponding author); tuula.a.oksanen@jyu.fi*

The wide-ranging Eurasian common lizard *Zootoca vivipara* (Lichtenstein, 1823) is remarkably uniform morphologically but highly varied in its karyotype. Previous studies have revealed two distinctly different chromosomal forms of *Z. v. vivipara* in the Baltic basin. Moreover, a zone of secondary contact between these forms has been localized on the southern Baltic Sea seashore. Intraspecific karyotype diversity for *Z. vivipara* and new zones of secondary contact have recently been suggested for other parts of the Baltic Sea seashore. We studied the karyotype of *Z. vivipara* in central, western and northern parts of Finland. All the individuals karyotyped represented the Russian form of *Z. v. vivipara* that differs from the western form of the subspecies located at the southern and western Baltic Sea seashore. Together with previous data sets, our results suggest intraspecific karyotype diversity in the northern and northwestern parts of Fennoscandia. The results give support to the hypothesis of *Z. vivipara*'s re-colonization of the Baltic Sea basin. Moreover, the results support the previous observations of Voipio (1961, 1968 and 1969) who has reported variability in the shield pattern of *Z. vivipara* in the same region.

## Introduction

The widely-ranged Eurasian common lizard *Zootoca vivipara* (Lichtenstein 1823) (family Lacertidae) is a squamate species with a huge distribution range from western Europe (the Pyrenees) throughout central, eastern and northern Europe up to eastern Asia (the Russian Far East, islands Sakhalin and Kunashir and northern Japan). The species is characterized by viviparous and oviparous reproduction in different populations (Braná & Bea 1987) and substantial geo-

graphic variation in body size and reproductive output (Horváthová et al. 2013). Despite such special characteristics *Z. vivipara* is remarkably uniform morphologically but polymorphic in its haplotype and karyotype. The species has 1) different diploid numbers:  $2n = 36/36$  in both sexes or  $2n = 35$  in female and  $2n = 36$  in male; 2) different size of female sex chromosomes: w microchromosome (m) or W macrochromosome (M); 3) different systems of sex chromosomes: Zw in female and ZZ in male or  $Z_1Z_2W$  in female and  $Z_1Z_1Z_2Z_2$  in male; 4) different morphology of

Table 1. Karyotype characteristics of subspecies and different forms of *Zootoca vivipara* (Lichtenstein 1823) and their distribution in Europe (Size of sex chromosomes: m = microchromosome, M = macrochromosome; Morphology of sex chromosomes a/A = acrocentric, ST = subtelocentric, SV = submetacentric; Mode of reproduction: O = oviparous, V = viviparous). A modification of Table 1. in Kupriyanova (2013).

N/N	2n ♂ / ♀	Sex chromosomes Size / System / Morphology			Mode of reproduction	Localities	Species, subspecies, chromosomal forms
<b>The first group of karyotype</b>							
1.	36A/36: 35A + 1a	m	Zw	a	O	Central, South- western Europe	<i>Z. vivipara</i> , now <i>Z. v. carniolica</i>
2.	36A/36: 35A + 1a	m	Zw	a	V	Central Europe	<i>Z. vivipara</i> , now <i>Z. v.</i> <i>vivipara</i> Hungarian form
<b>The second group of karyotype</b>							
3.	36A/35: 34A + 1A (A/ST)	M	Z <sub>1</sub> Z <sub>2</sub> W	A, A/ST	O	Western Europe, the Pyrenees	<i>Z. v. vivipara</i> Pyrenean form, now <i>Z. v. lousislantzi</i>
4.	36A/35: 34A + 1ST (A/ST)	M	Z <sub>1</sub> Z <sub>2</sub> W	ST, A/ST	V	Central Europe	<i>Z. vivipara</i> , now <i>Z. v.</i> <i>vivipara</i> Austrian form; <i>Z. v. pannonica?</i>
5.	36A/35: 34A + 1A	M	Z <sub>1</sub> Z <sub>2</sub> W	A	V	Asia, Eastern Europe, Baltic region	<i>Z. vivipara</i> , now <i>Z. v. vivipara</i> Russian form
6.	36A/35: 34A + 1SV	M	Z <sub>1</sub> Z <sub>2</sub> W	SV	V	Western, Central Europe, Baltic region	<i>Z. vivipara</i> , now <i>Z. v. vivipara</i> Western form

w and W sex chromosomes: acrocentric (a, A), subtelocentric (ST) or submetacentric (SV) and 5) some differences in cytogenetic and molecular structure of w and W sex chromosomes: heterochromatic amount and some other features (Table 1).

From all these karyotype characteristics six/seven separate chromosomal forms have been recognized among oviparous and viviparous females from different populations in Europe and in Asia. Among them, two new oviparous subspecies; *Z. v. carniolica* (Mayer, Böhme, Tiedemann & Bischoff 2000) and *Z. v. lousislantzi* (Arribas 2009). Two recent studies even suggest that *Z. v. carniolica* may be approaching species status (Lindtke et al. 2010, Cornetti et al. 2014). Subspecies *Z. v. vivipara* may be subdivided into four viviparous chromosomal forms, three of which are closely related, although the taxonomy of the latter is still questionable, they can be easily recognized by their 2n and some other karyotype characteristics (Table 1.).

In addition, several molecular and chromosomal studies have discussed the geographical distribution of different haplogroups (Heulin et al. 1999, 2011, Surget-Groba et al. 2001, 2006, Velekei et al. 2014) and chromosomal forms (Kupriyanova 1990, Kupriyanova & Böhme 1997, Kupriyanova et al. 2005, 2006, 2007, Odierna et al. 2001, Puky et al. 2004) of *Z. vivipara*. Chromosomal studies have shown that described subspecies and separate chromosomal forms of *Z. v. vivipara* have their distinct distribution ranges in Europe and in Asia (Table 1). In central Europe, subspecies and forms occur in allopatric, parapatric and sometimes mosaic populations. Some of them appear to inhabit small areas while others are relict and rare within one country. However, the western form of *Z. v. vivipara* and the Russian form of *Z. v. vivipara* occupy a vast territory in Europe and Asia. It has been indicated that the Russian form has in its female karyotype 34 acrocentric (A) chromosomes and 1 acrocentric (A) W sex chromosome (35 chromosomes in total). The latter has short arms at some metaphase plates

and it is close to subtelocentric (ST). Therefore, it is sometimes indicated as A/ST. Chromosomal formula is: ♀ $2n = 35: 34A + 1A$ , where W is A (or A/ST). The western form has in its female karyotype 34 acrocentric (A) chromosomes and 1 submetacentric (SV) W sex chromosome (the same 35 chromosomes in total). Chromosomal formula is: ♀ $2n = 35: 34A + 1SV$ , where W is SV. The males of all forms of *Z. v. vivipara* and of subspecies *Z. v. louisiantzi* have in their karyotype 36 acrocentric chromosomes: ♂ $2n = 36A$  with 4 acrocentric  $Z_1Z_1Z_2Z_2$  sex chromosomes (Table 1.).

Based on the karyotype (Kupriyanova 1990, 2004, Kupriyanova & Rudi 1990, Kupriyanova & Böhme 1997, Kupriyanova et al. 2007, Odierna et al. 1998, 2001) and the Mt DNA data (Surget-Groba et al. 2006, Velekei et al. 2014), the Russian form has been discovered from the eastern Carpathian throughout Russia up to Sakhalin island and northern Japan, whereas the western form has also been found in the populations in the eastern and western Carpathians as well as in central and in western Europe up to the Pyrenees. So far, the highest karyotype diversity has been discovered among the populations in the Carpathian Basin, due to which a centre of evolution of different chromosomal forms of *Z. vivipara* has been assumed to occur there (Kupriyanova & Böhme 1997, Odierna et al. 1998). Additionally, from all the available biogeographical and chromosomal data, it can be concluded that the Russian form is the most primitive one, whereas the western form has been derived from it (Kupriyanova 1990, Kupriyanova & Rudi 1990, Odierna et al. 1998).

Based on the karyotype markers, many specimens of western form *Z. v. vivipara* have been further identified in the western part of the Baltic region (Denmark, north of Germany, south of Sweden), while specimens of Russian form could be found in its eastern part (Estonia, north-west of Russia, south-east of Finland). Therefore, it was predicted that the southern Baltic Sea is a zone of secondary contact between these two chromosomal forms (Kupriyanova 1997).

In previous chromosomal studies, both the western and the Russian form of *Z. vivipara* have been identified on the limited territory of this seashore, namely the Kaliningrad oblast [Königsberger Gebiet] in western Russia (Kupriyanova et al. 2007, Kupriyanova & Melashchenko 2011).

Therefore, intraspecific karyotype diversity in northeastern Europe has been confirmed and a zone of secondary contact between these forms localized. In addition, the previous data by Kupriyanova & Melashchenko (2011) predicts more intraspecific karyotype diversity and several new zones of secondary contact in Kaliningrad oblast and in other parts of the southern Baltic Sea seashore. This has been confirmed by a karyological study of *Z. vivipara* where both forms were discovered in Poland for the first time in 2012 by Kupriyanova and Böhme (2012) (see Fig. 1). Moreover, two new zones of their secondary contact with allopatric and in one case with a parapatric distribution in Kaliningrad oblast were found in 2014 by Kupriyanova and Melashchenko (2015).

Thus, the geographically distinct distribution of both forms has been demonstrated and the border of their distribution area in this part of northeastern Europe has been verified. The previous data sets suggest that during the postglacial time, populations of *Z. vivipara* belonging to the western form of *Z. v. vivipara* have been re-colonizing the Kaliningrad region from the west and southwest, and those belonging to the Russian form of *Z. v. vivipara* from the east and south-east. Moreover, the previous data enables us to predict karyotype diversity of *Z. vivipara* and some new zone(s) of secondary contact between the two forms in other parts of the Baltic basin as well as a trend of re-colonization of the Baltic region by *Z. vivipara* during the post-glacial period (Kupriyanova & Melashchenko 2011, 2014, Kupriyanova & Böhme 2012).

To test these predictions, we focused on the diagnostics of morphologically uniform specimens of *Z. vivipara* from populations along the eastern and northern sides of Gulf of Bothnia. We collected 33 specimens of *Z. vivipara*, obtained chromosomes and studied several previously listed karyotype markers to evaluate the karyotype diversity of *Z. vivipara* on the eastern and northern coasts of the Baltic Sea. The data allowed us to 1) define the karyotype of *Z. vivipara* from central and western parts of Finland as well as from a southern part of northern Finland; 2) identify the specimens; 3) verify a border of distribution of different forms of *Z. vivipara* on studied regions and 4) test a hypothesis of re-colonization by specimens of *Z. vivipara* of this part of Fennoscandia.

## Materials and Methods

25 specimens from nine geographically distinct localities from central, western and northern Finland were collected and analyzed in May – June 2011 and 2012 (Table 2; Fig. 1). In addition, we analyzed eight individuals (6 females and 2 males) from an enclosure population originating from several natural populations in central Fin-

land and located at Konnevesi Research Station of the University of Jyväskylä (locality 1 in Table 2).

The chromosomes were obtained according to the scraping and air-drying method from intestinal epithelial and lung cells as well as from the germinal lamina (i.e. the ovarian area where the earliest stages of oogenesis occur) with using 0.05 % colchicines (Odierna *et al.*, 1993). In




Fig. 1. The locations sampled in this study presented as the numbered squares that match the coordinates in table 2. The map also shows wider distributions of Russian (■) and western (●) forms.

Table 2. Number and origin of specimens of *Zootoca vivipara* analyzed in this study.

Locality number	Number of female specimens	Number of male specimens	Locality	
<b>Central Finland</b>				
1	6	2	Central Finland (enclosure population)	–
2	2	0	Konnevesi	62°36.964'N 26°20.746'E
3	5	0	Vesanka	62°30.277'N 25°558'E
4	4	0	Muurame	62°05.159'N 25°36.353'E
<b>Western Finland</b>				
5	2	0	Alaveteli	63°42.351'N 23°17.784'E
6	4	0	Kortesjärvi	63°18.836'N 23°14.682'E
7	2	0	Vaasa	63°6.943'N 22°2.803'E
8	2	0	Närpiö	62°31.892'N 21°15.404'E
9	2	0	Kauhajoki	62°18.166'N 22°33.502'E
<b>Northern Finland (north of the Baltic coast)</b>				
10	2	0	Tornio	65°54.093'N 24°27.957'E

a subset of the samples, we used a different method where metaphase chromosomes were prepared from whole blood and a short term leucocyte culture in Kreavital Lymphocyte Karyotyping Medium with an addition of 0.1 ml 0.1 % phytohaemagglutinin M (Sigma-Aldrich) per 3.5 ml culture for 24 – 48 h and of 0.1 ml 0.002 % colchicine for 30 min (modification of the method of Moritz 1984, 1987). The slides were stained for 10 min with a 5 % Giemsa solution in pH 7 phosphate buffer. Metaphase plates suitable for chromosome analysis were obtained from all samples studied.

## Results and discussion

Chromosomal analysis showed that females of *Z. vivipara* from geographically separate localities 1–10 (Table 2; Fig. 1) have  $2n = 35:34$  acrocentric (A) macrochromosomes and one acrocentric macrochromosome (A), with short arms at rare metaphase plates, sometimes close to subtelocentric (A/ST). Acrocentric macrochromosome (A) is well known as W sex chromosome to *Z. vivipara* (Fig. 2 a–h). Therefore, chromosomal analysis identified these specimens as the Russian form of *Z. v. vivipara*.

A limited number of specimens (2–4 from each locality) does not allow us to assess inter-population or intra-population chromosomal variability (mosaics, polymorphism etc.). Nevertheless, we found for the first time that specimens of the Russian form inhabit the eastern and northern coast of the Baltic Sea (Fig. 1). The data indicates that the Russian form lives in many regions of Finland and in the southern part of northern Sweden. However, an earlier study has identified a western form in the southern and eastern parts of Sweden (Göteborg and Uppsala regions) (Kupriyanova et al. 1995). Moreover, according to molecular (Mt haplotype) data by Surget-Groba et al. (2006) two specimens of a western haplotype (VB haplogroup) have been identified from the south and the central-eastern Sweden (Runsten and Umeå localities) and positioned to western viviparous clade (clade E). At the same time, a specimen west to the border between Sweden and Finland (Kiruna locality) was identified as an eastern haplotype (VU haplogroup) and positioned to an eastern viviparous clade (clade D).

Comparison of molecular phylogenetic trees with karyotype characteristics of different chromosomal forms has demonstrated a good correlation between molecular and chromosomal data (see Kupriyanova 2004, 2013, Kupriyanova et




Fig. 2. Giemsa stained meta-phase plates of females of *Zootoca vivipara* from:

1. Central Finland (localities 1, 2, 4),
2. Western Finland (localities 5, 6, 7, 9) and
3. Northern Finland (locality 10).

Localities refer to table 2.

$2n = 34A + 1A (A/ST)$ . Arrows point to acrocentric- (A) (e, f, h) and acro-/subtelocentric (A/ST) (a, b, c, d, g) W sex chromosomes. According to karyotype markers these females belong to the Russian form of *Z. v. vivipara*.

al. 2006). Therefore, it is clear that chromosomal data supports the presence of main branches of the molecular trees of *Z. vivipara* and shows that their appearance is marked by the chromosomal rearrangements with the forming of several subspecies and separate chromosomal forms. From all these data, we may with confidence say that the specimens belonging to western viviparous clade (clade E, VB haplogroup) should be identified as western chromosomal form of *Z. v. vivipara* whereas those belonging to eastern viviparous clade (clade D, VU haplogroup) should be identified as Russian chromosomal form. Thus all these data points to the direction that the Russian chromosomal form of the subspecies *Z. v. vivipara* inhabits north-eastern part of Sweden.

To conclude, our chromosomal data demonstrates that 1) the Russian form of *Z. v. vivipara*

inhabits the central, western and southern parts of northern Finland; 2) these regions are not characterized by karyotype diversity of *Z. vivipara*; 3) diversity and a zone of secondary contact between two chromosomal forms of *Z. v. vivipara* may be predicted for other northern parts of Finland as well as for those of Sweden and Norway; 4) the border of the distribution area of the Russian form is located in the north-western and northern Baltic Sea seashore (in the northern parts of Sweden, Finland and/or Norway) and 5) the hypothesis of re-colonization of the area of the Baltic Sea by *Z. vivipara* is supported.

Regarding the re-colonization hypothesis, our results suggest that during the postglacial time, populations of *Z. vivipara* belonging to the western form of *Z. v. vivipara* came to the area from south and south-west whereas those belonging

to the Russian form of *Z. v. vivipara* moved into Fennoscandia from east and south-east. Additionally, the present data is consistent with the previous chromosomal results on the presence of the Russian form both in the southern and eastern parts of Finland and in a neighbouring Karelian Russia territory, near the border between Russia and Finland (Kupriyanova et al. 2005).

*Z. vivipara* is distributed all over Finland (e.g. Terhivuo 1993) and it should be stressed that our results regarding the chromosomal characteristics correlate with the data of Voipio (1961, 1968, 1969, 1992) reporting variability in the shield pattern of the *Z. vivipara* in Fennoscandia. Based on these patterns, Voipio pointed out that *Z. vivipara* populations in southern and central Sweden include specimens with western and central European type of shield patterns whereas specimens in northern Sweden, Finland and Russia (north of the 62° N) show patterns of the eastern type.

To summarize, we emphasize that the identification of specimens of *Z. vivipara* based on their morphology is very difficult and misidentifications may occur. Our results demonstrate the value of chromosome diagnostic of *Z. vivipara* from the geographically distant localities of Fennoscandia. Furthermore, intensive chromosomal studies of specimens from the areas of Finland, in particular those from its northern part, could show a presence of both chromosomal forms of the subspecies *Z. v. vivipara*. New chromosomal data could also give additional information about the karyotype diversity of *Z. vivipara* throughout the area of Fennoscandia and clarify the border of the distribution of the two forms in the region.

From the literature and the data obtained in this study, we predict that the western and Russian chromosomal forms of *Z. v. vivipara* occupy the northern and northwestern regions of Fennoscandia. However, more data from a wider set of localities, in particular from the northern part of Finland, Sweden and Norway, is needed to confirm this prediction as any chromosomal data for these territories is still missing. We would also like to stress that a more detailed study of a secondary contact zone and its characteristic (allopatry, sympatry, parapatry and/or hybrid zone) is needed as *Z. vivipara* may represent a group of cryptic taxa. The taxonomic status of the chromosomal forms *Z. v. vivipara* is still unclear and

under discussion in the literature (Kupriyanova 2004, Kupriyanova & Melashchenko 2011). A combination of different types of approaches (chromosomal, molecular, morphometric, life-history and behavioral) would be helpful in evaluating the biodiversity and conservation issues of these unique populations of *Z. vivipara*.

*Acknowledgments.* The study was supported by grant of President of Russian Federation for Support of Leading Scientific Schools No. NSH- 2990.2014.4. We wish to thank Janne Valkonen for the help with sampling, Bryan Maritz for checking and correcting the language and Konnevesi Research Station for providing the facilities.

## References

- Brana, F. & Bea, A. 1987: Bimodalité de reproduction chez *Lacerta vivipara*. — Bull. Soc. Herpetol. Fr. 44: 1–5.
- Cornetti, L., Menegon, M., Giovanni, G., Heulin, B. & Vernes, C. 2014: Mitochondrial and nuclear DNA survey of *Zootoca vivipara* across the eastern Italian Alps: evolutionary relationships, historical demography and conservation implications. — PLoS ONE 9(1): e85912. Doi:10.1371/journal.pone.0085912.
- Heulin, B., Surget-Groba, Y., Guiller, A., Guillaume, C. P. & Deunff, J. 1999: Comparisons of mtDNA sequences (16S rRNA Gene) between oviparous and viviparous strains of *Lacerta vivipara*: a preliminary study. — Mol. Ecol. 8: 1627–1631.
- Heulin B., Surget-Groba Y., Sinervo B., Miles D. & Guiller A. 2011: Dynamica of haplogroup frequencies and survival rates in a contact zone of two mtDNA lineages of the lizard *Lacerta vivipara*. — Ecography 34: 436–447.
- Horváthová, T., Cooney, C.R., Fitze, P.S., Oksanen, T.A., Jelić, D., Ghira, I., Uller, T. & Jandzik, D. 2013: Length of activity season drives geographic variation in body size of a widely distributed lizard. — Ecology and Evolution 3, 8: 2424–2442. Doi: 10.1002/ece3.613.
- Kupriyanova, L. 1990: Cytogenetic studies in lacertid lizard. — In: Olmo, E. (ed.), Cytogenetics of Amphibians and Reptiles, Birkhauser Verlag, Basel, pp. 242–245.
- Kupriyanova, L. 1997: Is the Baltic Sea basin a zone of secondary contact between different chromosomal forms of *Zootoca vivipara*? — Memoranda Soc. Fauna Flora Fennica 73: 115–117.
- Kupriyanova, L. 2004: Cytogenetical approaches to the problem of form-formation and subspeciation in the complex *Lacerta vivipara*. — Tsitologia 46, 7: 643–652 (in Russian).

- Kupriyanova, L. 2013: Modern chromosomal and molecular investigations of the Eurasian species *Zootoca vivipara* (Lichtenstein, 1823) (Lacertidae): results and perspectives. In: Modern herpetology: problems and ways of their solutions. — In: Ananjeva, N., Sromyatnikova, E., Doronin, I. (eds.), The first International Conference of the Young Herpetologists of Russia and Neighboring Countries. Saint-Petersburg, 25–27 Nov. 2013, pp. 25–31.
- Kupriyanova, L. & Rudi, E. 1990: Comparative karyological analysis of *Lacerta vivipara* (Lacertidae, Sauria) populations. — *Zool. Zh.* 69: 93–101 (in Russian).
- Kupriyanova, L., Andrén, C. & Nilson, G. 1995: Distribution of different forms of *Lacerta vivipara* in the Baltic Sea basin. — *Memoranda Soc. Fauna Flora Fennica* 71: 96–97.
- Kupriyanova, L. & Böhme, W. 1997: New data on the distribution of different forms of *Zootoca vivipara* in eastern and western Europe: chromosomal analysis. — *Herpetologia Bonnensis* 13: 199–205.
- Kupriyanova, L., Terhivuo, J. & Pasanen, S. 2005: Contribution to the knowledge of karyotype dispersal of the Common lizard (*Zootoca v. vivipara* Jacq) in eastern Fennoscandia. — *Memoranda Soc. Fauna Flora Fenn.* 81: 97–101.
- Kupriyanova, L., Mayer, W. & Böhme, W. 2006: Karyotype diversity of the Eurasian lizard *Zootoca vivipara* (Jacquin, 1787) from Central Europe and the evolution of viviparity. — In: Vences, M., Köhler, J., Ziegler, T., Böhme, W. (eds.), *Herpetologia Bonnensis II*, Koenig Museum Bonn and Societas Europaea Herpetologica pp. 67–72.
- Kupriyanova, L., Melashchenko, O. & Alekseev, P. 2007: Karyological investigations of populations of the lizard *Zootoca vivipara* from the Baltic Sea basin (western region of Russia). — *Tsitologia* 49: 600–606 (in Russian).
- Kupriyanova, L. & Melashchenko, O. 2011: The common Eurasian lizard *Zootoca vivipara* (Jacquin, 1787) from Russia: sex chromosomes, subspeciation, and colonization. — *Russian J. of Herpetol.* 18, 2: 99–104.
- Kupriyanova, L. & Böhme, W. 2012: Viviparous lizard *Zootoca vivipara*, Lacertidae from northeastern and central Europe: intraspecific karyotypic diversity. — *Zool. Zh.* 91, 11: 1428–1432 (in Russian).
- Kupriyanova, L. & Melashchenko, O. 2015: New data on intraspecific karyotype diversity of viviparous lizard (*Zootoca vivipara*, Lacertidae): distribution and colonization in western Russia. — *Zool. Zh.* 94, 1: 1–5 (in Russian).
- Lindtke, D., Mayer, W. & Böhme, W. 2010: Identification of a contact zone between oviparous and viviparous common lizards (*Zootoca vivipara*) in central Europe: reproductive strategies and natural hybridization. — *Salamandra* 46, 2: 73–82.
- Moritz, C. 1984: The origin and evolution of parthenogenesis in *Heteronotia binoei* (Gekkonidae). Chromosome banding studies. — *Chromosoma* 89:151–162.
- Moritz, C. 1987: Parthenogenesis in the tropical gekkonid lizard, *Nactus arnouxii*, (Sauria: Gekkonidae). — *Evolution* 41, 6: 1252–1266.
- Odierna, G., Kupriyanova, L., Capriglione, T. & Olmo, E. 1993: Evolution of sex chromosomes in lacertid lizards. — *Amphibia-Reptilia* 14: 1–11.
- Odierna, G., Aprea, G., Capriglione, T., Arribas, O., Kupriyanova, L. & Olmo, E. 1998: Progressive differentiation of the W sex-chromosome between oviparous and viviparous populations of *Zootoca vivipara* (Reptilia, Lacertidae). — *Ital. J. Zool.* 65: 295–303.
- Odierna, G., Heulin, B., Guillaume, C., Vogrin, N., Aprea, G., Capriglione, T., Surget-Groba, Y. & Kupriyanova, L. 2001: Evolutionary and biogeographical implications of the karyological variations in the oviparous and viviparous forms of the lizard *Lacerta* (*Zootoca*) *vivipara*. — *Ecography* 24: 332–340.
- Odierna, G., Aprea, G., Capriglione, T. & Puky, O. 2004: Chromosomal evidence for the double origin of viviparity in the European common lizards *Lacerta* (*Zootoca*) *vivipara*. — *Herpetol. J.* 14: 157–160.
- Puky, M., Adam, H., Surget-Groba, Y., Heulin, B. & Odierna, G. 2004: Fajvedelmi programok letjog-sultsaga es feladatai Magyarországon: az eleven-szulo gyik (*Zootoca vivipara* Mayer & Bischoff, 1996) vizsgálata nakeredmenyei – estanulsagai. *Termeszettudom.* — *Kozlemenyek* 11: 411–418.
- Surget-Groba, Y., Heulin, B., Guillaume, C., Thorpe, R., Kupriyanova, L., Vogrin, N., Maslak, R., Mazzotti, S., Venczel, M., Ghira, I., Odierna, G., Leontyeva, O., Monney, J. & Smith, N. 2001: Intraspecific Phylogeography of *Lacerta vivipara* and the evolution of viviparity. — *Molec. Phylogen. Evol.* 18: 449–459.
- Surget-Groba, Y., Heulin, B., Guillaume, C.-P., Puky, M., Semenov, D., Orlova, V., Kupriyanova, L., Ghira, I. & Smajd, B. 2006: Multiple origin of viviparity, or reversal from viviparity to oviparity and the evolution of viviparity. — *Biol. J. Linn. Soc.* 87: 1–11.
- Terhivuo, J. 1993: Provisional atlas and status of populations for the herpetofauna of Finland in 1980–92. — *Ann. Zool. Fennici* 30: 55–69.
- Velekei, B., Lakatos, F., Biro, P., Acs, E. & Puky, M. 2014: The genetic structure of *Zootoca vivipara* (Lichtenstein, 1823) populations did not support the existence of a north-south corridor of the VB haplogroup in eastern Hungary. *North-western Journal of Zoology* 10, 1: 187–189.
- Voipio, P. 1961: Über die Praefrontalia-Konstellationen bei den fennoskandischen Waldeidechen (*Lacerta vivipara* Jacquin). — *Arch. Soc. Vanamo* 16: 115–123.
- Voipio, P. 1968: Variation of the head-shield pattern in *Lacerta vivipara* Jacq. — *Ann. Zool. Fennici* 5: 315–323.
- Voipio, P. 1969: Variation of the postfrontal pileus in *Lacerta vivipara* Jacq. — *Ann. Zool. Fennici* 6: 269–213.
- Voipio, P. 1992: On pileus anomalies in the common lizard *Lacerta vivipara* in Finland – a morphogenetic problem revisited. — *Ann. Zool. Fennici* 28: 83–94.